

HODGE THEORY FOR COMBINATORIAL GEOMETRIES

KARIM ADIPRASITO, JUNE HUH, AND ERIC KATZ

1. INTRODUCTION

The combinatorial theory of matroids starts with Whitney [Whi35], who introduced matroids as models for independence in vector spaces and graphs. By definition, a *matroid* M is given by a closure operator defined on all subsets of a finite set E satisfying the Steinitz-Mac Lane exchange property:

For every subset I of E and every element a not in the closure of I ,

if a is in the closure of $I \cup \{b\}$, then b is in the closure of $I \cup \{a\}$.

The matroid is called *loopless* if the empty subset of E is closed, and is called a *combinatorial geometry* if in addition all single element subsets of E are closed. A closed subset of E is called a *flat* of M , and every subset of E has a well-defined rank and corank in the poset of all flats of M . The notion of matroid played a fundamental role in graph theory, coding theory, combinatorial optimization, and mathematical logic; we refer to [Wel71] and [Oxl11] for general introduction.

As a generalization of the chromatic polynomial of a graph [Bir12, Whi32], Rota defined for an arbitrary matroid M the *characteristic polynomial*

$$\chi_M(\lambda) = \sum_{I \subseteq E} (-1)^{|I|} \lambda^{\text{crk}(I)},$$

where the sum is over all subsets $I \subseteq E$ and $\text{crk}(I)$ is the corank of I in M [Rot64]. Equivalently, the characteristic polynomial of M is

$$\chi_M(\lambda) = \sum_F \mu(\emptyset, F) \lambda^{\text{crk}(F)},$$

where the sum is over all flats F of M and μ is the Möbius function of the poset of flats of M , see Chapters 7 and 8 of [Whi87]. Among the problems that withstood many advances in matroid theory are the following log-concavity conjectures formulated in the 1970s.

Write $r + 1$ for the *rank* of M , that is, the rank of E in the poset of flats of M .

Conjecture 1.1. Let $w_k(M)$ be the absolute value of the coefficient of λ^{r-k+1} in the characteristic polynomial of M . Then the sequence $w_k(M)$ is log-concave:

$$w_{k-1}(M)w_{k+1}(M) \leq w_k(M)^2 \text{ for all } 1 \leq k \leq r.$$

In particular, the sequence $w_k(M)$ is unimodal:

$$w_0(M) \leq w_1(M) \leq \cdots \leq w_l(M) \geq \cdots \geq w_r(M) \geq w_{r+1}(M) \text{ for some index } l.$$

We remark that the positivity of the numbers $w_k(M)$ is used to deduce the unimodality from the log-concavity [Wel76, Chapter 15].

For chromatic polynomials, the unimodality was conjectured by Read, and the log-concavity was conjectured by Hoggar [Rea68, Hog74]. The prediction of Read was then extended to arbitrary matroids by Rota and Heron, and the conjecture in its full generality was given by Welsh [Rot71, Her72, Wel76]. We refer to [Whi87, Chapter 8] and [Oxl11, Chapter 15] for overviews and historical accounts.

A subset $I \subseteq E$ is said to be *independent* in M if no element i in I is in the closure of $I \setminus \{i\}$. A related conjecture of Welsh and Mason concerns the number of independent subsets of E of given cardinality [Wel71, Mas72].

Conjecture 1.2. Let $f_k(M)$ be the number of independent subsets of E with cardinality k . Then the sequence $f_k(M)$ is log-concave:

$$f_{k-1}(M)f_{k+1}(M) \leq f_k(M)^2 \text{ for all } 1 \leq k \leq r.$$

In particular, the sequence $f_k(M)$ is unimodal:

$$f_0(M) \leq f_1(M) \leq \cdots \leq f_l(M) \geq \cdots \geq f_r(M) \geq f_{r+1}(M) \text{ for some index } l.$$

We prove Conjecture 1.1 and Conjecture 1.2 by constructing a “cohomology ring” of M that satisfies the hard Lefschetz theorem and the Hodge-Riemann relations, see Theorem 1.4.

1.1. Matroid theory has experienced a remarkable development in the past century, and has been connected to diverse areas such as topology [GM92], geometric model theory [Pil96], and noncommutative geometry [vN60]. The study of complex hyperplane arrangements provided a particularly strong connection, see for example [Sta07]. Most important for our purposes is the work of de Concini and Procesi on certain “wonderful” compactifications of hyperplane arrangement complements [DP95]. The original work focused only on realizable matroids, but Feichtner and Yuzvinsky [FY04] defined a commutative ring associated to an arbitrary matroid that specializes to the cohomology ring of a wonderful compactification in the realizable case.

Definition 1.3. Let S_M be the polynomial ring

$$S_M := \mathbb{R}[x_F | F \text{ is a nonempty proper flat of } M].$$

The *Chow ring* of M is defined to be the quotient

$$A^*(M)_{\mathbb{R}} := S_M / (I_M + J_M),$$

where I_M is the ideal generated by the quadratic monomials

$$x_{F_1} x_{F_2}, \quad F_1 \text{ and } F_2 \text{ are two incomparable nonempty proper flats of } M,$$

and J_M is the ideal generated by the linear forms

$$\sum_{i_1 \in F} x_{F - i_1} - \sum_{i_2 \in F} x_{F - i_2}, \quad i_1 \text{ and } i_2 \text{ are distinct elements of the ground set } E.$$

Conjecture 1.1 was proved for matroids realizable over \mathbb{C} in [Huh12] by relating $w_k(M)$ to the Milnor numbers of a hyperplane arrangement realizing M over \mathbb{C} . Subsequently in [HK12], using the intersection theory of wonderful compactifications and the Khovanskii-Teissier inequality [Laz04, Section 1.6], the conjecture was verified for matroids that are realizable over some field. Lenz used this result to deduce Conjecture 1.2 for matroids realizable over some field [Len12].

After the completion of [HK12], it was gradually realized that the validity of the Hodge-Riemann relations for the Chow ring of M is a vital ingredient for the proof of the log-concavity conjectures, see Theorem 1.4 below. While the Chow ring of M could be defined for arbitrary M , it was unclear how to formulate and prove the Hodge-Riemann relations. From the point of view of [FY04], the ring $A^*(M)_{\mathbb{R}}$ is the Chow ring of a smooth, but noncompact toric variety $X(\Sigma_M)$, and there is no obvious way to reduce to the classical case of projective varieties. In fact, we will see that $X(\Sigma_M)$ is ‘‘Chow equivalent’’ to a smooth or mildly singular projective variety over \mathbb{K} if and only if the matroid M is realizable over \mathbb{K} , see Theorem 5.12.

1.2. We are nearing a difficult chasm, as there is no reason to expect a working Hodge theory beyond the case of realizable matroids. Nevertheless, there was some evidence on the existence of such a theory for arbitrary matroids. For example, it was proved in [AS14], using the method of concentration of measure, that the log-concavity conjectures hold for c -arrangements in the sense of Goresky and MacPherson [GM92].

We now state the main theorem of this paper. A function c on the set of nonempty proper subsets of E is said to be *strictly submodular* if

$$c_{I_1} + c_{I_2} > c_{I_1 \cap I_2} + c_{I_1 \cup I_2} \quad \text{for any two incomparable subsets } I_1, I_2 \subseteq E,$$

where we replace c_{\emptyset} and c_E by zero whenever they appear in the above inequality. A strictly submodular function c defines an element

$$\ell(c) := \sum_F c_F x_F \in A^1(M)_{\mathbb{R}},$$

where the sum is over all nonempty proper flats of M . Note that the rank function of *any* matroid on E , and more generally any submodular function of the free matroid on E , can, when restricted to the set of nonempty proper subsets of E , be obtained as a *limit* of strictly submodular functions. We write ‘‘deg’’ for the isomorphism $A^r(M)_{\mathbb{R}} \simeq \mathbb{R}$ determined by the property that

$$\deg(x_{F_1} x_{F_2} \cdots x_{F_r}) = 1 \quad \text{for any flag of nonempty proper flats } F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_r.$$

Theorem 1.4. Let ℓ be an element of $A^1(M)_{\mathbb{R}}$ associated to a strictly submodular function.

- (1) (Hard Lefschetz theorem) For every nonnegative integer $q \leq \frac{r}{2}$, the multiplication by ℓ defines an isomorphism

$$L_\ell^q : A^q(M)_\mathbb{R} \longrightarrow A^{r-q}(M)_\mathbb{R}, \quad a \longmapsto \ell^{r-2q} \cdot a.$$

- (2) (Hodge–Riemann relations) For every nonnegative integer $q \leq \frac{r}{2}$, the multiplication by ℓ defines a symmetric bilinear form

$$Q_\ell^q : A^q(M)_\mathbb{R} \times A^q(M)_\mathbb{R} \longrightarrow \mathbb{R}, \quad (a_1, a_2) \longmapsto (-1)^q \deg(a_1 \cdot L_\ell^q a_2)$$

that is positive definite on the kernel of $\ell \cdot L_\ell^q$.

In fact, we will prove that the Chow ring of M satisfies the hard Lefschetz theorem and the Hodge-Riemann relations with respect to any strictly convex piecewise linear function on the tropical linear space Σ_M associated to M , see Theorem 8.8. This implies Theorem 1.4. Our proof of the hard Lefschetz theorem and the Hodge-Riemann relations for general matroids is inspired by an ingenious inductive proof of the analogous facts for simple polytopes given by McMullen [McM93] (compare also [CM02] for related ideas in a different context). To show that this program, with a considerable amount of work, extends beyond polytopes, is our main purpose here.

In Section 9, we show that the Hodge-Riemann relations, which are in fact stronger than the hard Lefschetz theorem, imply Conjecture 1.1 and Conjecture 1.2. We remark that, in the context of projective toric varieties, a similar reasoning leads to the Alexandrov-Fenchel inequality on mixed volumes of convex bodies. In this respect, broadly speaking the approach of the present paper can be viewed as following Rota’s idea that log-concavity conjectures should follow from their relation with the theory of mixed volumes of convex bodies, see [Kun95].

1.3. There are other combinatorial approaches to intersection theory for matroids. Mikhalkin et al. introduced an integral Hodge structure for arbitrary matroids modeled on the cohomology of hyperplane arrangement complements [IKMZ]. Adiprasito and Björner showed that an analogue of the Lefschetz hyperplane section theorem holds for all smooth (i.e. locally matroidal) projective tropical varieties [AB14]. We will discuss the relations with the above perspectives on Hodge theory for matroids in the upcoming paper [AHK15].

Theorem 1.4 should be compared with the counterexample to a version of Hodge conjecture for positive currents in [BH15]: The example used in [BH15] gives a tropical variety that satisfies the Poincaré duality, the hard Lefschetz theorem, but not the Hodge-Riemann relations. We will also discuss this in detail in [AHK15].

Finally, we remark that Zilber and Hrushovski have worked on subjects related to intersection theory for finitary combinatorial geometries, see [Hru92]. At present the relationship between their approach and ours is unclear.

Acknowledgements. The authors thank Patrick Brosnan, Eduardo Cattani, Ben Elias, Ehud Hrushovski, Gil Kalai, and Sam Payne for valuable conversations. Karim Adiprasito was supported by a Minerva Fellowship from the Max Planck Society and NSF Grant DMS-1128155. June Huh was supported by a Clay Research Fellowship and NSF Grant DMS-1128155. Eric Katz was supported by an NSERC Discovery grant.

2. FINITE SETS AND THEIR SUBSETS

2.1. Let E be a nonempty finite set of cardinality $n+1$, say $\{0, 1, \dots, n\}$. We write \mathbb{Z}^E for the free abelian group generated by the standard basis vectors \mathbf{e}_i corresponding to the elements $i \in E$. For an arbitrary subset $I \subseteq E$, we set

$$\mathbf{e}_I := \sum_{i \in I} \mathbf{e}_i.$$

We associate to the set E a dual pair of rank n free abelian groups

$$N_E := \mathbb{Z}^E / \langle \mathbf{e}_E \rangle, \quad M_E := \mathbf{e}_E^\perp \subset \mathbb{Z}^E, \quad \langle -, - \rangle : N_E \times M_E \longrightarrow \mathbb{Z}.$$

The corresponding real vector spaces will be denoted

$$N_{E, \mathbb{R}} := N_E \otimes_{\mathbb{Z}} \mathbb{R}, \quad M_{E, \mathbb{R}} := M_E \otimes_{\mathbb{Z}} \mathbb{R}.$$

We use the same symbols \mathbf{e}_i and \mathbf{e}_I to denote their images in N_E and $N_{E, \mathbb{R}}$.

The groups N and M associated to nonempty finite sets are related to each other in a natural way. For example, if F is a nonempty subset of E , then there is a surjective homomorphism

$$N_E \longrightarrow N_F, \quad \mathbf{e}_I \longmapsto \mathbf{e}_{I \cap F},$$

and an injective homomorphism

$$M_F \longrightarrow M_E, \quad \mathbf{e}_i - \mathbf{e}_j \longmapsto \mathbf{e}_i - \mathbf{e}_j.$$

If F is a nonempty proper subset of E , we have a decomposition

$$(\mathbf{e}_F^\perp \subset M_E) = (\mathbf{e}_{E \setminus F}^\perp \subset M_E) = M_F \oplus M_{E \setminus F}.$$

Dually, we have an isomorphism from the quotient space

$$N_E / \langle \mathbf{e}_F \rangle = N_E / \langle \mathbf{e}_{E \setminus F} \rangle \longrightarrow N_F \oplus N_{E \setminus F}, \quad \mathbf{e}_I \longmapsto \mathbf{e}_{I \cap F} \oplus \mathbf{e}_{I \setminus F}.$$

This isomorphism will be used later to analyze local structure of Bergman fans.

More generally, for any map between nonempty finite sets $\pi : E_1 \rightarrow E_2$, there is an associated homomorphism

$$\pi_N : N_{E_2} \longrightarrow N_{E_1}, \quad \mathbf{e}_I \longmapsto \mathbf{e}_{\pi^{-1}(I)},$$

and the dual homomorphism

$$\pi_M : M_{E_1} \longrightarrow M_{E_2}, \quad \mathbf{e}_i - \mathbf{e}_j \longmapsto \mathbf{e}_{\pi(i)} - \mathbf{e}_{\pi(j)}.$$

When π is surjective, π_N is injective and π_M is surjective.

2.2. Let $\mathcal{P}(E)$ be the poset of nonempty proper subsets of E . Throughout this section the symbol \mathcal{F} will stand for a totally ordered subset of $\mathcal{P}(E)$, that is, a flag of nonempty proper subsets of E :

$$\mathcal{F} = \{F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_l\} \subseteq \mathcal{P}(E).$$

We write $\min \mathcal{F}$ for the intersection of all members of \mathcal{F} . By definition, $\min \emptyset = E$.

Definition 2.1. When I is a proper subset of $\min \mathcal{F}$, we say that I is *compatible* with \mathcal{F} in E , and write $I < \mathcal{F}$.

The set of all compatible pairs in E form a poset under the relation

$$(I_1 < \mathcal{F}_1) \preceq (I_2 < \mathcal{F}_2) \iff I_1 \subseteq I_2 \text{ and } \mathcal{F}_1 \subseteq \mathcal{F}_2.$$

We note that any maximal compatible pair $I < \mathcal{F}$ gives a basis of the group N_E :

$$\{\mathbf{e}_i \text{ and } \mathbf{e}_F \text{ for } i \in I \text{ and } F \in \mathcal{F}\} \subseteq N_E.$$

If 0 is the unique element of E not in I and not in any member of \mathcal{F} , then the above basis of N_E is related to the basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ by an invertible upper triangular matrix.

Definition 2.2. For each compatible pair $I < \mathcal{F}$ in E , we define two polyhedra

$$\Delta_{I < \mathcal{F}} := \text{conv}\{\mathbf{e}_i \text{ and } \mathbf{e}_F \text{ for } i \in I \text{ and } F \in \mathcal{F}\} \subseteq N_{E, \mathbb{R}},$$

$$\sigma_{I < \mathcal{F}} := \text{cone}\{\mathbf{e}_i \text{ and } \mathbf{e}_F \text{ for } i \in I \text{ and } F \in \mathcal{F}\} \subseteq N_{E, \mathbb{R}}.$$

Since maximal compatible pairs give bases of N_E , the polytope $\Delta_{I < \mathcal{F}}$ is a simplex, and the cone $\sigma_{I < \mathcal{F}}$ is unimodular. Any proper subset of E is compatible with the empty flag in $\mathcal{P}(E)$, and the empty subset of E is compatible with any flag in $\mathcal{P}(E)$. Therefore we may write

$$\Delta_{I < \mathcal{F}} = \Delta_{I < \emptyset} * \Delta_{\emptyset < \mathcal{F}} \quad \text{and} \quad \sigma_{I < \mathcal{F}} = \sigma_{I < \emptyset} + \sigma_{\emptyset < \mathcal{F}}.$$

The set of all simplices of the form $\Delta_{I < \mathcal{F}}$ is in fact a simplicial complex, that is,

$$\Delta_{I_1 < \mathcal{F}_1} \cap \Delta_{I_2 < \mathcal{F}_2} = \Delta_{I_1 \cap I_2 < \mathcal{F}_1 \cap \mathcal{F}_2}.$$

This gives a geometric realization of the poset of compatible pairs in E .

2.3. An *order filter* \mathcal{P} of $\mathcal{P}(E)$ is a collection of nonempty proper subsets of E with the following property:

If $F_1 \subseteq F_2$ are nonempty proper subsets of E , then $F_1 \in \mathcal{P}$ implies $F_2 \in \mathcal{P}$.

Any such order filter cuts out a simplicial sphere in the simplicial complex of compatible pairs.

Definition 2.3. The *Bergman complex* of an order filter $\mathcal{P} \subseteq \mathcal{P}(E)$ is the collection of simplices

$$\Delta_{\mathcal{P}} := \{\Delta_{I < \mathcal{F}} \text{ for } I \notin \mathcal{P} \text{ and } \mathcal{F} \subseteq \mathcal{P}\}.$$

The *Bergman fan* of an order filter $\mathcal{P} \subseteq \mathcal{P}(E)$ is the collection of simplicial cones

$$\Sigma_{\mathcal{P}} := \{\sigma_{I < \mathcal{F}} \text{ for } I \notin \mathcal{P} \text{ and } \mathcal{F} \subseteq \mathcal{P}\}.$$

The Bergman complex $\Delta_{\mathcal{P}}$ is a simplicial complex because \mathcal{P} is an order filter. We will see below that the Bergman fan $\Sigma_{\mathcal{P}}$ indeed is a fan, that is,

$$\sigma_{I_1 < \mathcal{F}_1} \cap \sigma_{I_2 < \mathcal{F}_2} = \sigma_{I_1 \cap I_2 < \mathcal{F}_1 \cap \mathcal{F}_2} \text{ for } \sigma_{I_1 < \mathcal{F}_1}, \sigma_{I_2 < \mathcal{F}_2} \in \Sigma_{\mathcal{P}}.$$

The extreme cases $\mathcal{P} = \emptyset$ and $\mathcal{P} = \mathcal{P}(E)$ correspond to familiar geometric objects. When \mathcal{P} is empty, the collection $\Sigma_{\mathcal{P}}$ is the normal fan of the standard n -dimensional simplex

$$\Delta_n := \text{conv}\{\mathbf{e}_0, \mathbf{e}_1, \dots, \mathbf{e}_n\} \subseteq \mathbb{R}^E.$$

When \mathcal{P} contains all nonempty proper subsets of E , the collection $\Sigma_{\mathcal{P}}$ is the normal fan of the n -dimensional permutohedron

$$\Pi_n := \text{conv}\{(x_0, x_1, \dots, x_n) \mid x_0, x_1, \dots, x_n \text{ is a permutation of } 0, 1, \dots, n\} \subseteq \mathbb{R}^E.$$

Proposition 2.4 below shows that, in general, the Bergman complex $\Delta_{\mathcal{P}}$ is a simplicial sphere and $\Sigma_{\mathcal{P}}$ is a complete unimodular fan.

Proposition 2.4. For any order filter $\mathcal{P} \subseteq \mathcal{P}(E)$, the collection $\Sigma_{\mathcal{P}}$ is the normal fan of a polytope.

Proof. We show that $\Sigma_{\mathcal{P}}$ can be obtained from Σ_{\emptyset} by performing a sequence of stellar subdivisions. In other words, a polytope corresponding to $\Sigma_{\mathcal{P}}$ can be obtained by repeatedly truncating faces of the standard simplex Δ_n .

For this we choose a sequence of order filters obtained by adding a single subset in \mathcal{P} at a time:

$$\emptyset, \dots, \mathcal{P}_-, \mathcal{P}_+, \dots, \mathcal{P} \text{ with } \mathcal{P}_+ = \mathcal{P}_- \cup \{Z\}.$$

The corresponding sequence of Σ interpolates between the collections Σ_{\emptyset} and $\Sigma_{\mathcal{P}}$:

$$\Sigma_{\emptyset} \rightsquigarrow \dots \rightsquigarrow \Sigma_{\mathcal{P}_-} \rightsquigarrow \Sigma_{\mathcal{P}_+} \rightsquigarrow \dots \rightsquigarrow \Sigma_{\mathcal{P}}.$$

The modification in the middle replaces the cones of the form $\sigma_{Z < \mathcal{F}}$ with the sums of the form

$$\sigma_{\emptyset < \{Z\}} + \sigma_{I < \mathcal{F}},$$

where I is any proper subset of Z . In other words, the modification is the stellar subdivision of $\Sigma_{\mathcal{P}_-}$ relative to the cone $\sigma_{Z < \emptyset}$. Since a stellar subdivision of the normal fan of a polytope is the normal fan of a polytope, by induction we know that the collection $\Sigma_{\mathcal{P}}$ is the normal fan of a polytope. \square

3. MATROIDS AND THEIR FLATS

3.1. Let M be a loopless matroid of rank $r + 1$ on the ground set E . We denote rk_M , crk_M , and cl_M for the rank function, the corank function, and the closure operator of M respectively. We

omit the subscripts when M is understood from the context. If F is a nonempty proper flat of M , we write

$$\begin{aligned} M^F &:= \text{the restriction of } M \text{ to } F, \text{ a loopless matroid on } F \text{ of rank } = \text{rk}_M(F), \\ M_F &:= \text{the contraction of } M \text{ by } F, \text{ a loopless matroid on } E \setminus F \text{ of rank } = \text{crk}_M(F). \end{aligned}$$

We refer to [Oxl11] and [Wel76] for basic notions of matroid theory.

Let $\mathcal{P}(M)$ be the poset of nonempty proper flats of M . There is an injective map from the poset of the restriction

$$\iota^F : \mathcal{P}(M^F) \longrightarrow \mathcal{P}(M), \quad G \longmapsto G,$$

and an injective map from the poset of the contraction

$$\iota_F : \mathcal{P}(M_F) \longrightarrow \mathcal{P}(M), \quad G \longmapsto G \cup F.$$

We identify the flats of M_F with the flats of M containing F using ι_F . If \mathcal{P} is a subset of $\mathcal{P}(M)$, we set

$$\mathcal{P}^F := (\iota^F)^{-1} \mathcal{P} \quad \text{and} \quad \mathcal{P}_F := (\iota_F)^{-1} \mathcal{P}.$$

3.2. Throughout this section the symbol \mathcal{F} will stand for a totally ordered subset of $\mathcal{P}(M)$, that is, a flag of nonempty proper flats of M :

$$\mathcal{F} = \{F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_l\} \subseteq \mathcal{P}(M).$$

As before, we write $\min \mathcal{F}$ for the intersection of all members of \mathcal{F} inside E . We extend the notion of compatibility in Definition 2.1 to the case when the matroid M is not Boolean.

Definition 3.1. When I is a subset of $\min \mathcal{F}$ of cardinality less than $\text{rk}_M(\min \mathcal{F})$, we say that I is *compatible* with \mathcal{F} in M , and write $I <_M \mathcal{F}$.

Since any flag of nonempty proper flats of M has length at most r , any cone

$$\sigma_{I <_M \mathcal{F}} = \text{cone} \left\{ \mathbf{e}_i \text{ and } \mathbf{e}_F \text{ for } i \in I \text{ and } F \in \mathcal{F} \right\}$$

associated to a compatible pair in M has dimension at most r . Conversely, any such cone is contained in an r -dimensional cone of the same type: For this one may take

$$\begin{aligned} I' &= \text{a subset that is maximal among those containing } I \text{ and compatible with } \mathcal{F} \text{ in } M, \\ \mathcal{F}' &= \text{a flag of flats maximal among those containing } \mathcal{F} \text{ and compatible with } I' \text{ in } M, \end{aligned}$$

or alternatively take

$$\begin{aligned} \mathcal{F}' &= \text{a flag of flats maximal among those containing } \mathcal{F} \text{ and compatible with } I \text{ in } M, \\ I' &= \text{a subset that is maximal among those containing } I \text{ and compatible with } \mathcal{F}' \text{ in } M. \end{aligned}$$

We note that any subset of E with cardinality at most r is compatible in M with the empty flag of flats, and the empty subset of E is compatible in M with any flag of nonempty proper flats of M . Therefore we may write

$$\Delta_{I <_M \mathcal{F}} = \Delta_{I <_M \emptyset} * \Delta_{\emptyset <_M \mathcal{F}} \quad \text{and} \quad \sigma_{I <_M \mathcal{F}} = \sigma_{I <_M \emptyset} + \sigma_{\emptyset <_M \mathcal{F}}.$$

The set of all simplices associated to compatible pairs in M form a simplicial complex, that is,

$$\Delta_{I_1 <_M \mathcal{F}_1} \cap \Delta_{I_2 <_M \mathcal{F}_2} = \Delta_{I_1 \cap I_2 <_M \mathcal{F}_1 \cap \mathcal{F}_2}.$$

3.3. An *order filter* \mathcal{P} of $\mathcal{P}(M)$ is a collection of nonempty proper flats of M with the following property:

If $F_1 \subseteq F_2$ are nonempty proper flats of M , then $F_1 \in \mathcal{P}$ implies $F_2 \in \mathcal{P}$.

We write $\widehat{\mathcal{P}} := \mathcal{P} \cup \{E\}$ for the order filter of the lattice of flats of M generated by \mathcal{P} .

Definition 3.2. The *Bergman fan* of an order filter $\mathcal{P} \subseteq \mathcal{P}(M)$ is the set of simplicial cones

$$\Sigma_{M, \mathcal{P}} := \left\{ \sigma_{I < \mathcal{F}} \text{ for } \text{cl}_M(I) \notin \widehat{\mathcal{P}} \text{ and } \mathcal{F} \subseteq \mathcal{P} \right\}.$$

The *reduced Bergman fan* of \mathcal{P} is the subset of the Bergman fan

$$\widetilde{\Sigma}_{M, \mathcal{P}} := \left\{ \sigma_{I <_M \mathcal{F}} \text{ for } \text{cl}_M(I) \notin \widehat{\mathcal{P}} \text{ and } \mathcal{F} \subseteq \mathcal{P} \right\}.$$

When $\mathcal{P} = \mathcal{P}(M)$, we omit \mathcal{P} from the notation and write the Bergman fan by Σ_M .

We note that the *Bergman complex* and the *reduced Bergman complex* $\widetilde{\Delta}_{M, \mathcal{P}} \subseteq \Delta_{M, \mathcal{P}}$, defined in analogous ways using the simplices $\Delta_{I < \mathcal{F}}$ and $\Delta_{I <_M \mathcal{F}}$, share the same set of vertices.

Two extreme cases give familiar geometric objects. When \mathcal{P} is the set of all nonempty proper flats of M , we have

$$\Sigma_M = \Sigma_{M, \mathcal{P}} = \widetilde{\Sigma}_{M, \mathcal{P}} = \text{the fine subdivision of the tropical linear space of } M \text{ [AK06].}$$

When \mathcal{P} is empty, we have

$$\widetilde{\Sigma}_{M, \emptyset} = \text{the } r\text{-dimensional skeleton of the normal fan of the simplex } \Delta_n,$$

and $\Sigma_{M, \emptyset}$ is the fan whose maximal cones are $\sigma_{F < \emptyset}$ for rank r flats F of M . We remark that

$$\Delta_{M, \emptyset} = \text{the Alexander dual of the matroid complex } \text{IN}(M^*) \text{ of the dual matroid } M^*.$$

See [Bjo92] for basic facts on the matroid complexes and [MS05, Chapter 5] for the Alexander dual of a simplicial complex.

We show that, in general, the Bergman fan and the reduced Bergman fan are indeed fans, and the reduced Bergman fan is pure of dimension r .

Proposition 3.3. The collection $\Sigma_{M, \mathcal{P}}$ is a subfan of the normal fan of a polytope.

Proof. Since \mathcal{P} is an order filter, any face of a cone in $\Sigma_{M, \mathcal{P}}$ is in $\Sigma_{M, \mathcal{P}}$. Therefore it is enough to show that there is a normal fan of a polytope that contains $\Sigma_{M, \mathcal{P}}$ as a subset.

For this we consider the order filter of $\mathcal{P}(E)$ generated by \mathcal{P} , that is, the collection of sets

$$\widetilde{\mathcal{P}} := \{\text{nonempty proper subset of } E \text{ containing a flat in } \mathcal{P}\} \subseteq \mathcal{P}(E).$$

If the closure of $I \subseteq E$ in M is not in $\widetilde{\mathcal{P}}$, then I does not contain any flat in \mathcal{P} , and hence

$$\Sigma_{M, \mathcal{P}} \subseteq \Sigma_{\widetilde{\mathcal{P}}}.$$

The latter collection is the normal fan of a polytope by Proposition 2.4. \square

Since \mathcal{P} is an order filter, any face of a cone in $\widetilde{\Sigma}_{M, \mathcal{P}}$ is in $\widetilde{\Sigma}_{M, \mathcal{P}}$, and hence $\widetilde{\Sigma}_{M, \mathcal{P}}$ is a subfan of $\Sigma_{M, \mathcal{P}}$. It follows that the reduced Bergman fan also is a subfan of the normal fan of a polytope.

Proposition 3.4. The reduced Bergman fan $\widetilde{\Sigma}_{M, \mathcal{P}}$ is pure of dimension r .

Proof. Let I be a subset of E whose closure is not in \mathcal{P} , and let \mathcal{F} be a flag of flats in \mathcal{P} compatible with I in M . We show that there are I' containing I and \mathcal{F}' containing \mathcal{F} such that

$$I' <_M \mathcal{F}', \quad \text{cl}_M(I') \notin \widetilde{\mathcal{P}}, \quad \mathcal{F}' \subseteq \mathcal{P}, \quad \text{and} \quad |I'| + |\mathcal{F}'| = r.$$

First choose any flag of flats \mathcal{F}' that is maximal among those containing \mathcal{F} , contained in \mathcal{P} , and compatible with I in M . Next choose any flat F of M that is maximal among those containing I and strictly contained in $\min \mathcal{F}'$.

We note that, by the maximality of F and the maximality of \mathcal{F}' respectively,

$$\text{rk}_M(F) = \text{rk}_M(\min \mathcal{F}') - 1 = r - |\mathcal{F}'|.$$

Since the rank of a set is at most its cardinality, the above implies

$$|I| \leq r - |\mathcal{F}'| \leq |F|.$$

This shows that there is I' containing I , contained in F , and with cardinality exactly $r - |\mathcal{F}'|$. Any such I' is automatically compatible with \mathcal{F}' in M .

We show that the closure of I' is not in \mathcal{P} by showing that the flat F is not in \mathcal{P} . If otherwise, by the maximality of \mathcal{F}' , the set I cannot be compatible in M with the flag $\{F\}$, meaning

$$|I| \geq \text{rk}_M(F).$$

The above implies that the closure of I in M , which is not in \mathcal{P} , is equal to F . This gives the desired contradiction. \square

Our inductive approach to the hard Lefschetz theorem and the Hodge-Riemann relations for matroids is modeled on the observation that any facet of a permutohedron is the product of two smaller permutohedrons. We note below that the Bergman fan $\Sigma_{M, \mathcal{P}}$ has an analogous local structure when M has no parallel elements.

Recall that the *star* of a cone σ in a fan Σ in a latticed vector space $N_{\mathbb{R}}$ is the fan

$$\text{star}(\sigma, \Sigma) := \{\overline{\sigma'} \mid \overline{\sigma'} \text{ is the image in } N_{\mathbb{R}}/\langle\sigma\rangle \text{ of a cone } \sigma' \text{ in } \Sigma \text{ containing } \sigma\}.$$

When σ is a ray generated by its primitive generator \mathbf{e} , we write $\text{star}(\mathbf{e}, \Sigma)$ for the star of σ in Σ .

Proposition 3.5. Let M be a loopless matroid on E , and let \mathcal{P} be an order filter of $\mathcal{P}(M)$.

(1) If F is a flat in \mathcal{P} , then the isomorphism $N_E/\langle\mathbf{e}_F\rangle \rightarrow N_F \oplus N_{E \setminus F}$ induces a bijection

$$\text{star}(\mathbf{e}_F, \Sigma_{M, \mathcal{P}}) \longrightarrow \Sigma_{M^F, \mathcal{P}^F} \times \Sigma_{M^c}.$$

(2) If $\{i\}$ is a proper flat of M , then the isomorphism $N_E/\langle\mathbf{e}_i\rangle \rightarrow N_{E \setminus \{i\}}$ induces a bijection

$$\text{star}(\mathbf{e}_i, \Sigma_{M, \mathcal{P}}) \longrightarrow \Sigma_{M \setminus \{i\}, \mathcal{P} \setminus \{i\}}.$$

Under the same assumptions, the stars of \mathbf{e}_F and \mathbf{e}_i in the reduced Bergman fan $\widetilde{\Sigma}_{M, \mathcal{P}}$ admit analogous descriptions.

When M is not a combinatorial geometry, the star of \mathbf{e}_i in $\Sigma_{M, \mathcal{P}}$ is not necessarily a product of smaller Bergman fans. However, when M is a combinatorial geometry, Proposition 3.5 shows that the star of every ray in $\Sigma_{M, \mathcal{P}}$ is a product of at most two Bergman fans.

4. PIECEWISE LINEAR FUNCTIONS AND THEIR CONVEXITY

4.1. Piecewise linear functions on possibly incomplete fans will play an important role throughout the paper. In this section, we prove several general properties concerning convexity of such functions, working with a dual pair free abelian groups

$$\langle -, - \rangle : N \times M \longrightarrow \mathbb{Z}, \quad N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}, \quad M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R},$$

and a unimodular fan Σ in the latticed vector space $N_{\mathbb{R}}$. The set of primitive ray generators of Σ will be denoted V_{Σ} .

We say that a function $\ell : |\Sigma| \rightarrow \mathbb{R}$ is *piecewise linear* if it is continuous and the restriction of ℓ to any cone in Σ is the restriction of a linear function on $N_{\mathbb{R}}$. The function ℓ is said to be *integral* if

$$\ell(|\Sigma| \cap N) \subseteq \mathbb{Z},$$

and the function ℓ is said to be *positive* if

$$\ell(|\Sigma| \setminus \{0\}) \subseteq \mathbb{R}_{>0}.$$

An important example of a piecewise linear function on Σ is the *Courant function* $x_{\mathbf{e}}$ associated to a primitive ray generator \mathbf{e} of Σ , whose values at V_{Σ} are given by the Kronecker delta function. Since Σ is unimodular, the Courant functions are integral, and they form a basis of the group of integral piecewise linear functions on Σ :

$$\text{PL}(\Sigma) = \left\{ \sum_{\mathbf{e} \in V_{\Sigma}} c_{\mathbf{e}} x_{\mathbf{e}} \mid c_{\mathbf{e}} \in \mathbb{Z} \right\} \simeq \mathbb{Z}^{V_{\Sigma}}.$$

An integral linear function on $N_{\mathbb{R}}$ restricts to an integral piecewise linear function on Σ , giving a homomorphism

$$\text{res}_{\Sigma} : M \longrightarrow \text{PL}(\Sigma), \quad m \longmapsto \sum_{\mathbf{e} \in V_{\Sigma}} \langle \mathbf{e}, m \rangle x_{\mathbf{e}}.$$

We denote the cokernel of the restriction map by

$$A^1(\Sigma) := \text{PL}(\Sigma)/M.$$

In general, this group may have torsion, even under our assumption that Σ is unimodular. When integral piecewise linear functions ℓ and ℓ' on Σ differ by the restriction of an integral linear function on $N_{\mathbb{R}}$, we say that ℓ and ℓ' are *equivalent* over \mathbb{Z} .

Note that the group of piecewise linear functions modulo linear functions on Σ can be identified with the tensor product

$$A^1(\Sigma)_{\mathbb{R}} := A^1(\Sigma) \otimes_{\mathbb{Z}} \mathbb{R}.$$

When piecewise linear functions ℓ and ℓ' on Σ differ by the restriction of a linear function on $N_{\mathbb{R}}$, we say that ℓ and ℓ' are *equivalent*.

We describe three basic pullback homomorphisms between the groups A^1 . Let Σ' be a subfan of Σ , and let σ be a cone in Σ .

- (1) The restriction of functions from Σ to Σ' defines a surjective homomorphism

$$\text{PL}(\Sigma) \longrightarrow \text{PL}(\Sigma'),$$

and this descends to a surjective homomorphism

$$\text{p}_{\Sigma' \subseteq \Sigma} : A^1(\Sigma) \longrightarrow A^1(\Sigma').$$

In terms of Courant functions, $\text{p}_{\Sigma' \subseteq \Sigma}$ is uniquely determined by its values

$$x_{\mathbf{e}} \longmapsto \begin{cases} x_{\mathbf{e}} & \text{if } \mathbf{e} \text{ is in } V_{\Sigma'}, \\ 0 & \text{if otherwise.} \end{cases}$$

- (2) Any integral piecewise linear function ℓ on Σ is equivalent over \mathbb{Z} to an integral ℓ' that is zero on σ , and the choice of such ℓ' is unique up to an integral linear function on $N_{\mathbb{R}}/\langle \sigma \rangle$. Therefore we have a surjective homomorphism

$$\text{p}_{\sigma \in \Sigma} : A^1(\Sigma) \longrightarrow A^1(\text{star}(\sigma, \Sigma)),$$

uniquely determined by its values on $x_{\mathbf{e}}$ for primitive ray generators \mathbf{e} not contained in σ :

$$x_{\mathbf{e}} \longmapsto \begin{cases} x_{\bar{\mathbf{e}}} & \text{if there is a cone in } \Sigma \text{ containing } \mathbf{e} \text{ and } \sigma, \\ 0 & \text{if otherwise.} \end{cases}$$

Here $\bar{\mathbf{e}}$ is the image of \mathbf{e} in the quotient space $N_{\mathbb{R}}/\langle \sigma \rangle$.

- (3) A piecewise linear function on the product of two fans $\Sigma_1 \times \Sigma_2$ is the sum of its restrictions to the subfans

$$\Sigma_1 \times \{0\} \subseteq \Sigma_1 \times \Sigma_2 \quad \text{and} \quad \{0\} \times \Sigma_2 \subseteq \Sigma_1 \times \Sigma_2.$$

Therefore we have an isomorphism

$$\text{PL}(\Sigma_1 \times \Sigma_2) \simeq \text{PL}(\Sigma_1) \oplus \text{PL}(\Sigma_2),$$

and this descends to an isomorphism

$$\text{p}_{\Sigma_1, \Sigma_2} : A^1(\Sigma_1 \times \Sigma_2) \simeq A^1(\Sigma_1) \oplus A^1(\Sigma_2).$$

4.2. We define the *link* of a cone σ in Σ to be the collection

$$\text{link}(\sigma, \Sigma) := \{\sigma' \in \Sigma \mid \sigma' \text{ is contained in a cone in } \Sigma \text{ containing } \sigma, \text{ and } \sigma \cap \sigma' = \{0\}\}.$$

Note that the link of σ in Σ is a subfan of Σ .

Definition 4.1. Let ℓ be a piecewise linear function on Σ , and let σ be a cone in Σ .

- (1) The function ℓ is *convex* around σ if it is equivalent to a piecewise linear function that is zero on σ and nonnegative on the link of σ .
- (2) The function ℓ is *strictly convex* around σ if it is equivalent to a piecewise linear function that is zero on σ and positive on the link of σ .

The function ℓ is *convex* if it is convex around every cone in Σ , and *strictly convex* if it is strictly convex around every cone in Σ .

When Σ is complete, the function ℓ is convex in the sense of Definition 4.1 if and only if it is convex in the usual sense:

$$\ell(\mathbf{u}_1 + \mathbf{u}_2) \leq \ell(\mathbf{u}_1) + \ell(\mathbf{u}_2) \quad \text{for } \mathbf{u}_1, \mathbf{u}_2 \in N_{\mathbb{R}}.$$

In general, writing ι for the inclusion of the torus orbit closure corresponding to σ in the toric variety of Σ , we have

$$\ell \text{ is convex around } \sigma \iff \iota^* \text{ of the class of the divisor associated to } \ell \text{ is effective.}$$

For a detailed discussion and related notions of convexity from the point of view of toric geometry, see [GM12].

Definition 4.2. The *ample cone* of Σ is the open convex cone

$$\mathcal{K}_{\Sigma} := \{\text{classes of strictly convex piecewise linear functions on } \Sigma\} \subseteq A^1(\Sigma)_{\mathbb{R}}.$$

The *nef cone* of Σ is the closed convex cone

$$\mathcal{N}_{\Sigma} := \{\text{classes of convex piecewise linear functions on } \Sigma\} \subseteq A^1(\Sigma)_{\mathbb{R}}.$$

Note that the closure of the ample cone \mathcal{K}_{Σ} is contained in the nef cone \mathcal{N}_{Σ} . In many interesting cases, the reverse inclusion also holds.

Proposition 4.3. If \mathcal{K}_Σ is nonempty, then \mathcal{N}_Σ is the closure of \mathcal{K}_Σ .

Proof. If ℓ_1 is a convex piecewise linear function and ℓ_2 is strictly convex piecewise linear function on Σ , then the sum $\ell_1 + \epsilon \ell_2$ is strictly convex for every positive number ϵ . This shows that the nef cone of Σ is in the closure of the ample cone of Σ . \square

We record here that the various pullbacks of an ample class are ample. The proof is straightforward from Definition 4.1.

Proposition 4.4. Let Σ' be a subfan of Σ , σ be a cone in Σ , and let $\Sigma_1 \times \Sigma_2$ be a product fan.

(1) The pullback homomorphism $p_{\Sigma' \subseteq \Sigma}$ induces a map between the ample cones

$$\mathcal{K}_\Sigma \longrightarrow \mathcal{K}_{\Sigma'}.$$

(2) The pullback homomorphism $p_{\sigma \in \Sigma}$ induces a map between the ample cones

$$\mathcal{K}_\Sigma \longrightarrow \mathcal{K}_{\text{star}(\sigma, \Sigma)}.$$

(3) The isomorphism p_{Σ_1, Σ_2} induces a bijective map between the ample cones

$$\mathcal{K}_{\Sigma_1 \times \Sigma_2} \longrightarrow \mathcal{K}_{\Sigma_1} \times \mathcal{K}_{\Sigma_2}.$$

It follows from the first item that any subfan of the normal fan of a polytope has a nonempty ample cone. In particular, by Proposition 3.3, the Bergman fan $\Sigma_{M, \mathcal{P}}$ has a nonempty ample cone.

Strictly convex piecewise linear functions on the normal fan of the permutohedron can be described in a particularly nice way: A piecewise linear function on $\Sigma_{\mathcal{P}(E)}$ is strictly convex if and only if it is of the form

$$\sum_{F \in \mathcal{P}(E)} c_F x_F, \quad c_{F_1} + c_{F_2} > c_{F_1 \cap F_2} + c_{F_1 \cup F_2} \text{ for any incomparable } F_1, F_2, \text{ with } c_\emptyset = c_E = 0.$$

For this and related results, see [BB11]. The restriction of any such *strictly submodular function* gives a strictly convex function on the Bergman fan Σ_M , and defines an ample class on Σ_M .

4.3. We specialize to the case of matroids and prove basic properties of convex piecewise linear functions on the Bergman fan $\Sigma_{M, \mathcal{P}}$. We write $\mathcal{K}_{M, \mathcal{P}}$ for the ample cone of $\Sigma_{M, \mathcal{P}}$, and $\mathcal{N}_{M, \mathcal{P}}$ for the nef cone of $\Sigma_{M, \mathcal{P}}$.

Proposition 4.5. Let M be a loopless matroid on E , and let \mathcal{P} be an order filter of $\mathcal{P}(M)$.

(1) The nef cone of $\Sigma_{M, \mathcal{P}}$ is equal to the closure of the ample cone of $\Sigma_{M, \mathcal{P}}$:

$$\overline{\mathcal{K}_{M, \mathcal{P}}} = \mathcal{N}_{M, \mathcal{P}}.$$

(2) The ample cone of $\Sigma_{M, \mathcal{P}}$ is equal to the interior of the nef cone of $\Sigma_{M, \mathcal{P}}$:

$$\mathcal{K}_{M, \mathcal{P}} = \mathcal{N}_{M, \mathcal{P}}^\circ.$$

Proof. Propositions 3.3 shows that the ample cone $\mathcal{K}_{M, \mathcal{P}}$ is nonempty. Therefore, by Proposition 4.3, the nef cone $\mathcal{N}_{M, \mathcal{P}}$ is equal to the closure of $\mathcal{K}_{M, \mathcal{P}}$.

The second assertion can be deduced from the first using the following general property of convex sets: An open convex set is equal to the interior of its closure. \square

The main result here is that the ample cone and its ambient vector space

$$\mathcal{K}_{M, \mathcal{P}} \subseteq A^1(\Sigma_{M, \mathcal{P}})_{\mathbb{R}}$$

depend only on \mathcal{P} and the combinatorial geometry of M , see Proposition 4.8 below. We set

$$\bar{E} := \{A \mid A \text{ is a rank 1 flat of } M\}.$$

Definition 4.6. The *combinatorial geometry* of M is the simple matroid \bar{M} on \bar{E} determined by its poset of nonempty proper flats $\mathcal{P}(\bar{M}) = \mathcal{P}(M)$.

The set of primitive ray generators of $\Sigma_{M, \mathcal{P}}$ is the disjoint union

$$\{e_i \mid \text{the closure of } i \text{ in } M \text{ is not in } \mathcal{P}\} \cup \{e_F \mid F \text{ is a flat in } \mathcal{P}\} \subseteq N_{E, \mathbb{R}},$$

and the set of primitive ray generators of $\Sigma_{\bar{M}, \mathcal{P}}$ is the disjoint union

$$\{e_A \mid A \text{ is a rank 1 flat of } M \text{ not in } \mathcal{P}\} \cup \{e_F \mid F \text{ is a flat in } \mathcal{P}\} \subseteq N_{\bar{E}, \mathbb{R}}.$$

The corresponding Courant functions on the Bergman fans will be denoted x_i , x_F , and x_A , x_F respectively.

Let π be the surjective map between the ground sets of M and \bar{M} given by the closure operator of M . We fix an arbitrary section ι of π by choosing an element from each rank 1 flat:

$$\pi : E \longrightarrow \bar{E}, \quad \iota : \bar{E} \longrightarrow E, \quad \pi \circ \iota = \text{id}.$$

The maps π and ι induce the horizontal homomorphisms in the diagram

$$\begin{array}{ccc} \text{PL}(\Sigma_{M, \mathcal{P}}) & \begin{array}{c} \xleftarrow{\pi_{\text{PL}}} \\ \xrightarrow{\iota_{\text{PL}}} \end{array} & \text{PL}(\Sigma_{\bar{M}, \mathcal{P}}) \\ \text{res} \uparrow & & \uparrow \text{res} \\ M_E & \begin{array}{c} \xleftarrow{\pi_M} \\ \xrightarrow{\iota_M} \end{array} & M_{\bar{E}} \end{array}$$

where the homomorphism π_{PL} is obtained by setting

$$x_i \longmapsto x_{\pi(i)}, \quad x_F \longmapsto x_F, \quad \text{for elements } i \text{ whose closure is not in } \mathcal{P}, \text{ and for flats } F \text{ in } \mathcal{P},$$

and the homomorphism ι_{PL} is obtained by setting

$$x_A \longmapsto x_{\iota(A)}, \quad x_F \longmapsto x_F, \quad \text{for rank 1 flats } A \text{ not in } \mathcal{P}, \text{ and for flats } F \text{ in } \mathcal{P}.$$

In the diagram above, we have

$$\pi_{\text{PL}} \circ \text{res} = \text{res} \circ \pi_M, \quad \iota_{\text{PL}} \circ \text{res} = \text{res} \circ \iota_M, \quad \pi_{\text{PL}} \circ \iota_{\text{PL}} = \text{id}, \quad \pi_M \circ \iota_M = \text{id}.$$

Proposition 4.7. The homomorphism π_{PL} induces an isomorphism

$$\pi_{\text{PL}} : A^1(\Sigma_{M, \mathcal{P}}) \longrightarrow A^1(\Sigma_{\overline{M}, \mathcal{P}}).$$

The homomorphism ι_{PL} induces the inverse isomorphism

$$\iota_{\text{PL}} : A^1(\Sigma_{\overline{M}, \mathcal{P}}) \longrightarrow A^1(\Sigma_{M, \mathcal{P}}).$$

We use the same symbols to denote the isomorphisms $A^1(\Sigma_{M, \mathcal{P}})_{\mathbb{R}} \xleftrightarrow{\cong} A^1(\Sigma_{\overline{M}, \mathcal{P}})_{\mathbb{R}}$.

Proof. It is enough to check that the composition $\iota_{\text{PL}} \circ \pi_{\text{PL}}$ is the identity. Let i and j be elements whose closures are not in \mathcal{P} . Consider the linear function on $N_{E, \mathbb{R}}$ given by the integral vector

$$\mathbf{e}_i - \mathbf{e}_j \in M_E.$$

The restriction of this linear function to $\Sigma_{M, \mathcal{P}}$ is the linear combination

$$\text{res}(\mathbf{e}_i - \mathbf{e}_j) = \left(x_i + \sum_{i \in F \in \mathcal{P}} x_F \right) - \left(x_j + \sum_{j \in F \in \mathcal{P}} x_F \right).$$

If i and j have the same closure, then a flat contains i if and only if it contains j , and hence the linear function witnesses that the piecewise linear functions x_i and x_j are equivalent over \mathbb{Z} . It follows that $\iota_{\text{PL}} \circ \pi_{\text{PL}} = \text{id}$. \square

The maps π and ι induce simplicial maps between the Bergman complexes

$$\Delta_{M, \mathcal{P}} \begin{array}{c} \xrightarrow{\pi_{\Delta}} \\ \xleftarrow{\iota_{\Delta}} \end{array} \Delta_{\overline{M}, \mathcal{P}}, \quad \Delta_{I < \mathcal{F}} \mapsto \Delta_{\pi(I) < \mathcal{F}}, \quad \Delta_{\mathcal{F} < \mathcal{F}} \mapsto \Delta_{\iota(\mathcal{F}) < \mathcal{F}}.$$

The simplicial map π_{Δ} collapses those simplices containing vectors of parallel elements, and

$$\pi_{\Delta} \circ \iota_{\Delta} = \text{id}.$$

The other composition $\iota_{\Delta} \circ \pi_{\Delta}$ is a deformation retraction. For this note that

$$\Delta_{I < \mathcal{F}} \in \Delta_{M, \mathcal{P}} \implies \iota_{\Delta} \circ \pi_{\Delta}(\Delta_{I < \mathcal{F}}) \cup \Delta_{I < \mathcal{F}} \subseteq \Delta_{\pi^{-1}\pi I < \mathcal{F}}.$$

The simplex $\Delta_{\pi^{-1}\pi I < \mathcal{F}}$ is in $\Delta_{M, \mathcal{P}}$, and hence we can find a homotopy $\iota_{\Delta} \circ \pi_{\Delta} \simeq \text{id}$.

Proposition 4.8. The isomorphism π_{PL} restricts to a bijective map between the ample cones

$$\mathcal{H}_{M, \mathcal{P}} \longrightarrow \mathcal{H}_{\overline{M}, \mathcal{P}}.$$

Proof. By Proposition 4.5, it is enough to show that π_{PL} restricts to a bijective map

$$\mathcal{N}_{M, \mathcal{P}} \longrightarrow \mathcal{N}_{\overline{M}, \mathcal{P}}.$$

We use the following maps corresponding to π_{Δ} and ι_{Δ} :

$$\Sigma_{M, \mathcal{P}} \begin{array}{c} \xrightarrow{\pi_{\Sigma}} \\ \xleftarrow{\iota_{\Sigma}} \end{array} \Sigma_{\overline{M}, \mathcal{P}}, \quad \sigma_{I < \mathcal{F}} \mapsto \sigma_{\pi(I) < \mathcal{F}}, \quad \sigma_{\mathcal{F} < \mathcal{F}} \mapsto \sigma_{\iota(\mathcal{F}) < \mathcal{F}}.$$

One direction is more direct: The homomorphism ι_{PL} maps a convex piecewise linear function $\bar{\ell}$ to a convex piecewise linear function $\iota_{\text{PL}}(\bar{\ell})$. Indeed, for any cone $\sigma_{I < \mathcal{F}}$ in $\Sigma_{\mathbb{M}, \mathcal{F}}$,

$$\begin{aligned} \left(\bar{\ell} \text{ is zero on } \sigma_{\pi(I) < \mathcal{F}} \text{ and nonnegative on the link of } \sigma_{\pi(I) < \mathcal{F}} \text{ in } \Sigma_{\bar{\mathbb{M}}, \mathcal{F}} \right) &\implies \\ \left(\iota_{\text{PL}}(\bar{\ell}) \text{ is zero on } \sigma_{\pi^{-1}\pi(I) < \mathcal{F}} \text{ and nonnegative on the link of } \sigma_{\pi^{-1}\pi(I) < \mathcal{F}} \text{ in } \Sigma_{\mathbb{M}, \mathcal{F}} \right) & \\ \implies \left(\iota_{\text{PL}}(\bar{\ell}) \text{ is zero on } \sigma_{I < \mathcal{F}} \text{ and nonnegative on the link of } \sigma_{I < \mathcal{F}} \text{ in } \Sigma_{\mathbb{M}, \mathcal{F}} \right). & \end{aligned}$$

Next we show the other direction: The homomorphism π_{PL} maps a convex piecewise linear function ℓ to a convex piecewise linear function $\pi_{\text{PL}}(\ell)$. The main claim is that, for any cone $\sigma_{\mathcal{J} < \mathcal{F}}$ in $\Sigma_{\bar{\mathbb{M}}, \mathcal{F}}$,

$$\ell \text{ is convex around } \sigma_{\pi^{-1}(\mathcal{J}) < \mathcal{F}} \implies \pi_{\text{PL}}(\ell) \text{ is convex around } \sigma_{\mathcal{J} < \mathcal{F}}.$$

This can be deduced from the following identities between the subfans of $\Sigma_{\mathbb{M}, \mathcal{F}}$:

$$\begin{aligned} \pi_{\Sigma}^{-1} \left(\text{the set of all faces of } \sigma_{\mathcal{J} < \mathcal{F}} \right) &= \left(\text{the set of all faces of } \sigma_{\pi^{-1}(\mathcal{J}) < \mathcal{F}} \right), \\ \pi_{\Sigma}^{-1} \left(\text{the link of } \sigma_{\mathcal{J} < \mathcal{F}} \text{ in } \Sigma_{\bar{\mathbb{M}}, \mathcal{F}} \right) &= \left(\text{the link of } \sigma_{\pi^{-1}(\mathcal{J}) < \mathcal{F}} \text{ in } \Sigma_{\mathbb{M}, \mathcal{F}} \right). \end{aligned}$$

It is straightforward to check the two equalities from the definitions of $\Sigma_{\mathbb{M}, \mathcal{F}}$ and $\Sigma_{\bar{\mathbb{M}}, \mathcal{F}}$. \square

Remark 4.9. Note that a Bergman fan and the corresponding reduced Bergman fan share the same set of primitive ray generators. Therefore we have isomorphisms

$$\begin{array}{ccc} A^1(\Sigma_{\mathbb{M}, \mathcal{F}}) & \xleftrightarrow{\quad} & A^1(\Sigma_{\bar{\mathbb{M}}, \mathcal{F}}) \\ \updownarrow & & \updownarrow \\ A^1(\tilde{\Sigma}_{\mathbb{M}, \mathcal{F}}) & \xleftrightarrow{\quad} & A^1(\tilde{\Sigma}_{\bar{\mathbb{M}}, \mathcal{F}}). \end{array}$$

We remark that there are inclusion maps between the corresponding ample cones

$$\begin{array}{ccc} \mathcal{H}_{\mathbb{M}, \mathcal{F}} & \xlongequal{\quad} & \mathcal{H}_{\bar{\mathbb{M}}, \mathcal{F}} \\ \downarrow & & \downarrow \\ \tilde{\mathcal{H}}_{\mathbb{M}, \mathcal{F}} & \longleftarrow & \tilde{\mathcal{H}}_{\bar{\mathbb{M}}, \mathcal{F}}. \end{array}$$

In general, all three inclusions shown above may be strict.

5. HOMOLOGY AND COHOMOLOGY

5.1. Let Σ be a unimodular fan in an n -dimensional latticed vector space $N_{\mathbb{R}}$, and let Σ_k be the set of k -dimensional cones in Σ . If τ is a codimension 1 face of a unimodular cone σ , we write

$$\mathbf{e}_{\sigma/\tau} := \text{the primitive generator of the unique 1-dimensional face of } \sigma \text{ not in } \tau.$$

Definition 5.1. A k -dimensional Minkowski weight on Σ is a function

$$\omega : \Sigma_k \longrightarrow \mathbb{Z}$$

which satisfies the *balancing condition*: For every $(k-1)$ -dimensional cone τ in Σ ,

$$\sum_{\tau \subset \sigma} \omega(\sigma) \mathbf{e}_{\sigma/\tau} \text{ is contained in the subspace generated by } \tau.$$

The group of Minkowski weights on Σ is the group

$$\text{MW}_*(\Sigma) := \bigoplus_{k \in \mathbb{Z}} \text{MW}_k(\Sigma),$$

where $\text{MW}_k(\Sigma) := \{k\text{-dimensional Minkowski weights on } \Sigma\} \subseteq \mathbb{Z}^{\Sigma_k}$.

The group of Minkowski weights was studied by Fulton and Sturmfels in the context of toric geometry [FS97]. An equivalent notion of stress space was independently pursued by Lee in [Lee96]. We record here some immediate properties of the group of Minkowski weights on Σ .

(1) The group $\text{MW}_0(\Sigma)$ is canonically isomorphic to the group of integers:

$$\text{MW}_0(\Sigma) = \mathbb{Z}^{\Sigma_0} \simeq \mathbb{Z}.$$

(2) The group $\text{MW}_1(\Sigma)$ is perpendicular to the image of the restriction map from M :

$$\text{MW}_1(\Sigma) = \text{im}(\text{res}_\Sigma)^\perp \subseteq \mathbb{Z}^{\Sigma_1}.$$

(3) The group $\text{MW}_k(\Sigma)$ is trivial for k negative or k larger than the dimension of Σ .

If Σ is in addition complete, then an n -dimensional weight on Σ satisfies the balancing condition if and only if it is constant. Therefore, in this case, there is a canonical isomorphism

$$\text{MW}_n(\Sigma) \simeq \mathbb{Z}.$$

We show that the Bergman fan Σ_M has the same property with respect to its dimension r .

Proposition 5.2. An r -dimensional weight on Σ_M satisfies the balancing condition if and only if it is constant.

It follows that there is a canonical isomorphism $\text{MW}_r(\Sigma_M) \simeq \mathbb{Z}$.

Lemma 5.3. The Bergman fan Σ_M is connected in codimension 1.

We remark that Lemma 5.3 is a direct consequence of the shellability of Δ_M , see [Bjo92].

Proof. The claim is that, for any two r -dimensional cones $\sigma_{\mathcal{F}}, \sigma_{\mathcal{G}}$ in Σ_M , there is a sequence

$$\sigma_{\mathcal{F}} = \sigma_0 \supset \tau_1 \subset \sigma_1 \supset \cdots \subset \sigma_{l-1} \supset \tau_l \subset \sigma_l = \sigma_{\mathcal{G}},$$

where τ_i is a common facet of σ_{i-1} and σ_i in Σ_M . We express this by writing $\sigma_{\mathcal{F}} \sim \sigma_{\mathcal{G}}$.

We prove by induction on the rank of M . If $\min \mathcal{F} = \min \mathcal{G}$, then the induction hypothesis applied to $M_{\min \mathcal{F}}$ shows that

$$\sigma_{\mathcal{F}} \sim \sigma_{\mathcal{G}}.$$

If otherwise, we choose a flag of nonempty proper flats \mathcal{H} maximal among those satisfying $\min \mathcal{F} \cup \min \mathcal{G} < \mathcal{H}$. By the induction hypothesis applied to $M_{\min \mathcal{F}}$, we have

$$\sigma_{\mathcal{F}} \sim \sigma_{\{\min \mathcal{F}\} \cup \mathcal{H}}.$$

Similarly, by the induction hypothesis applied to $M_{\min \mathcal{G}}$, we have

$$\sigma_{\mathcal{G}} \sim \sigma_{\{\min \mathcal{G}\} \cup \mathcal{H}}.$$

Since any 1-dimensional fan is connected in codimension 1, this complete the induction. \square

Proof of Proposition 5.2. The proof is based on the *flat partition property* for matroids M on E :

If F is a flat of M , then the flats of M that cover F partition $E \setminus F$.

Let $\tau_{\mathcal{G}}$ be a codimension 1 cone in the Bergman fan Σ_M , and set

$$V_{\text{star}(\mathcal{G})} := \text{the set of primitive ray generators of the star of } \tau_{\mathcal{G}} \text{ in } \Sigma_M \subseteq N_{E, \mathbb{R}} / \langle \tau_{\mathcal{G}} \rangle.$$

The flat partition property applied to the restrictions of M shows that, first, the sum of all the vectors in $V_{\text{star}(\mathcal{G})}$ is zero and, second, any proper subset of $V_{\text{star}(\mathcal{G})}$ is linearly independent. Therefore, for an r -dimensional weight ω on Σ_M ,

$$\omega \text{ satisfies the balancing condition at } \tau_{\mathcal{G}} \iff \omega \text{ is constant on cones containing } \tau_{\mathcal{G}}.$$

By the connectedness of Lemma 5.3, the latter condition for every $\tau_{\mathcal{G}}$ implies that ω is constant. \square

5.2. We continue to work with a unimodular fan Σ in $N_{\mathbb{R}}$. As before, we write V_{Σ} for the set of primitive ray generators of Σ . Let S_{Σ} be the polynomial ring over \mathbb{Z} with variables indexed by V_{Σ} :

$$S_{\Sigma} := \mathbb{Z}[x_{\mathbf{e}}]_{\mathbf{e} \in V_{\Sigma}}.$$

For each k -dimensional cone σ in Σ , we associate a degree k square-free monomial

$$x_{\sigma} := \prod_{\mathbf{e} \in \sigma} x_{\mathbf{e}}.$$

The subgroup of S_{Σ} generated by all such monomials x_{σ} will be denoted

$$Z^k(\Sigma) := \bigoplus_{\sigma \in \Sigma_k} \mathbb{Z}x_{\sigma}.$$

Let $Z^*(\Sigma)$ be the sum of $Z^k(\Sigma)$ over all nonnegative integers k .

Definition 5.4. The *Chow ring* of Σ is the commutative graded algebra

$$A^*(\Sigma) := S_\Sigma / (I_\Sigma + J_\Sigma),$$

where I_Σ and J_Σ are the ideals of S_Σ defined by

$$\begin{aligned} I_\Sigma &:= \text{the ideal generated by the square-free monomials not in } Z^*(\Sigma), \\ J_\Sigma &:= \text{the ideal generated by the linear forms } \sum_{\mathbf{e} \in V_\Sigma} \langle \mathbf{e}, m \rangle x_{\mathbf{e}} \text{ for } m \in M. \end{aligned}$$

We write $A^k(\Sigma)$ for the degree k component of $A^*(\Sigma)$, and set

$$A^*(\Sigma)_{\mathbb{R}} := A^*(\Sigma) \otimes_{\mathbb{Z}} \mathbb{R} \quad \text{and} \quad A^k(\Sigma)_{\mathbb{R}} := A^k(\Sigma) \otimes_{\mathbb{Z}} \mathbb{R}.$$

If we identify the variables of S_Σ with the Courant functions on Σ , then the degree 1 component of $A^*(\Sigma)$ agrees with the group introduced in Section 4:

$$A^1(\Sigma) = \text{PL}(\Sigma)/M.$$

Note that the pullback homomorphisms between A^1 introduced in that section uniquely extend to graded ring homomorphisms between A^* :

(1) The homomorphism $p_{\Sigma' \subseteq \Sigma}$ uniquely extends to a surjective graded ring homomorphism

$$p_{\Sigma' \subseteq \Sigma} : A^*(\Sigma) \longrightarrow A^*(\Sigma').$$

(2) The homomorphism $p_{\sigma \in \Sigma}$ uniquely extends to a surjective graded ring homomorphism

$$p_{\sigma \in \Sigma} : A^*(\Sigma) \longrightarrow A^*(\text{star}(\sigma, \Sigma)).$$

(3) The isomorphism p_{Σ_1, Σ_2} uniquely extends to a graded ring isomorphism

$$p_{\Sigma_1, \Sigma_2} : A^*(\Sigma_1 \times \Sigma_2) \longrightarrow A^*(\Sigma_1) \otimes_{\mathbb{Z}} A^*(\Sigma_2).$$

We remark that the Chow ring $A^*(\Sigma)_{\mathbb{R}}$ can be identified with the ring of piecewise polynomial functions on Σ modulo linear functions on $N_{\mathbb{R}}$, see [Bil89].

Proposition 5.5. The group $A^k(\Sigma)$ is generated by $Z^k(\Sigma)$ for each nonnegative integer k .

In particular, if k larger than the dimension of Σ , then $A^k(\Sigma) = 0$.

Proof. Let σ be a cone in Σ , let $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_l$ be its primitive ray generators. and consider a degree k monomial of the form

$$x_{\mathbf{e}_1}^{k_1} x_{\mathbf{e}_2}^{k_2} \cdots x_{\mathbf{e}_l}^{k_l}, \quad k_1 \geq k_2 \geq \cdots \geq k_l \geq 1.$$

We show that the image of this monomial in $A^k(\Sigma)$ is in the span of $Z^k(\Sigma)$.

We do this by descending induction on the dimension of σ . If $\dim \sigma = k$, there is nothing to prove. If otherwise, we use the unimodularity of σ to choose $m \in M$ such that

$$\langle \mathbf{e}_1, m \rangle = -1 \quad \text{and} \quad \langle \mathbf{e}_2, m \rangle = \cdots = \langle \mathbf{e}_l, m \rangle = 0.$$

This shows that, modulo the relations given by I_Σ and J_Σ , we have

$$x_{\mathbf{e}_1}^{k_1} x_{\mathbf{e}_2}^{k_2} \cdots x_{\mathbf{e}_k}^{k_l} = x_{\mathbf{e}_1}^{k_1-1} x_{\mathbf{e}_2}^{k_2} \cdots x_{\mathbf{e}_l}^{k_l} \sum_{\mathbf{e} \in \text{link}(\sigma)} \langle \mathbf{e}, m \rangle x_{\mathbf{e}},$$

where the sum is over the set of primitive ray generators of the link of σ in Σ . The induction hypothesis applies to each of the terms in the expansion of the right-hand side. \square

The group of k -dimensional weights on Σ can be identified with the dual of $Z^k(\Sigma)$ under the tautological isomorphism

$$t_\Sigma : \mathbb{Z}^{\Sigma^k} \longrightarrow \text{Hom}_{\mathbb{Z}}(Z^k(\Sigma), \mathbb{Z}), \quad \omega \longmapsto \left(x_\sigma \longmapsto \omega(\sigma) \right).$$

By Proposition 5.5, the target of t_Σ contains $\text{Hom}_{\mathbb{Z}}(A^k(\Sigma), \mathbb{Z})$ as a subgroup.

Proposition 5.6. The isomorphism t_Σ restricts to the bijection between the subgroups

$$\text{MW}_k(\Sigma) \longrightarrow \text{Hom}_{\mathbb{Z}}(A^k(\Sigma), \mathbb{Z}).$$

The bijection in Proposition 5.6 is an analogue of the Kronecker duality homomorphism in algebraic topology. We use it to define the *cap product*

$$A^l(\Sigma) \times \text{MW}_k(\Sigma) \longrightarrow \text{MW}_{k-l}(\Sigma), \quad \xi \cap \omega(\sigma) := t_\Sigma \omega(\xi \cdot x_\sigma).$$

This makes the group $\text{MW}_*(\Sigma)$ a graded module over the Chow ring $A^*(\Sigma)$.

Proof. The homomorphisms from $A^k(\Sigma)$ to \mathbb{Z} bijectively correspond to the homomorphisms from $Z^k(\Sigma)$ to \mathbb{Z} which vanish on the subgroup

$$Z^k(\Sigma) \cap (I_\Sigma + J_\Sigma) \subseteq Z^k(\Sigma).$$

The main point is that this subgroup is generated by polynomials of the form

$$\left(\sum_{\mathbf{e} \in \text{link}(\tau)} \langle \mathbf{e}, m \rangle x_{\mathbf{e}} \right) x_\tau,$$

where τ is a $(k-1)$ -dimensional cone of Σ and m is an element perpendicular to $\langle \tau \rangle$. It follows that a k -dimensional weight ω corresponds to a homomorphism $A^k(\Sigma) \rightarrow \mathbb{Z}$ if and only if

$$\sum_{\tau \subset \sigma} \omega(\sigma) \langle \mathbf{e}_{\sigma/\tau}, m \rangle = 0 \text{ for all } m \in \langle \tau \rangle^\perp,$$

where the sum is over all k -dimensional cones σ in Σ containing τ . Since $\langle \tau \rangle^{\perp\perp} = \langle \tau \rangle$, the latter condition is equivalent to the balancing condition on ω at τ . \square

5.3. The ideals I_Σ and J_Σ have a particularly simple description when $\Sigma = \Sigma_M$. In this case, we label the variables of S_Σ by the nonempty proper flats of M , and write

$$S_\Sigma = \mathbb{Z}[x_F]_{F \in \mathcal{P}(M)}.$$

For a flag of nonempty proper flats \mathcal{F} , we set $x_{\mathcal{F}} = \prod_{F \in \mathcal{F}} x_F$.

(Incomparability relations) The ideal I_Σ is generated by the quadratic monomials

$$x_{F_1} x_{F_2},$$

where F_1 and F_2 are two incomparable nonempty proper flats of M .

(Linear relations) The ideal J_Σ is generated by the linear forms

$$\sum_{i_1 \in F} x_F - \sum_{i_2 \in F} x_F,$$

where i_1 and i_2 are distinct elements of the ground set E .

The quotient ring $A^*(\Sigma_M)$ and its generalizations were studied by Feichtner and Yuzvinsky in [FY04].

Definition 5.7. To an element i in E , we associate linear forms

$$\alpha_{M,i} := \sum_{i \in F} x_F, \quad \beta_{M,i} := \sum_{i \notin F} x_F.$$

Their classes in $A^*(\Sigma_M)$, which are independent of i , will be written α_M and β_M respectively.

We show that $A^r(\Sigma_M)$ is generated by the element α_M^r , where r is the dimension of Σ_M .

Proposition 5.8. Let $F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_k$ be any flag of nonempty proper flats of M .

(1) If the rank of F_m is not m for some $m \leq k$, then

$$x_{F_1} x_{F_2} \cdots x_{F_k} \alpha_M^{r-k} = 0 \in A^r(\Sigma_M).$$

(2) If the rank of F_m is m for all $m \leq k$, then

$$x_{F_1} x_{F_2} \cdots x_{F_k} \alpha_M^{r-k} = \alpha_M^r \in A^r(\Sigma_M).$$

In particular, for any two maximal flags of nonempty proper flats \mathcal{F}_1 and \mathcal{F}_2 of M ,

$$x_{\mathcal{F}_1} = x_{\mathcal{F}_2} \in A^r(\Sigma_M).$$

Since $\text{MW}_r(\Sigma_M)$ is isomorphic to \mathbb{Z} , this implies that $A^r(\Sigma_M)$ is isomorphic to \mathbb{Z} , see Proposition 5.10.

Proof. As a general observation, we note that for any element i not in a nonempty proper flat F ,

$$x_F \alpha_M = x_F \left(\sum_G x_G \right) \in A^*(\Sigma_M),$$

where the sum is over all proper flats containing F and $\{i\}$. In particular, if the rank of F is r , then the product is zero.

We prove the first assertion by descending induction on k , which is necessarily less than r . If $k = r - 1$, then the rank of F_k should be r , and hence the product is zero. For general k , we choose an element i not in F_k . By the observation made above, we have

$$x_{F_1} x_{F_2} \cdots x_{F_k} \alpha_M^{r-k} = x_{F_1} x_{F_2} \cdots x_{F_k} \left(\sum_G x_G \right) \alpha_M^{r-k-1},$$

where the sum is over all proper flats containing F and $\{i\}$. The right-hand side is zero by the induction hypothesis for $k + 1$ applied to each of the terms in the expansion.

We prove the second assertion by ascending induction on k . When $k = 1$, we choose an element i in F_k . By the first part of the proposition for $k = 1$, we have

$$x_{F_1} \alpha_M^{r-1} = \alpha_M^r.$$

For general k , we choose an element i in $F_k \setminus F_{k-1}$. By the first part of the proposition for k , we have

$$x_{F_1} x_{F_2} \cdots x_{F_{k-1}} x_{F_k} \alpha_M^{r-k} = x_{F_1} x_{F_2} \cdots x_{F_{k-1}} \left(\sum_G x_G \right) \alpha_M^{r-k},$$

where the sum is over all proper flats containing F_{k-1} and $\{i\}$. The right-hand side is α_M^r by the induction hypothesis for $k - 1$. \square

When Σ is complete, Fulton and Sturmfels showed in [FS97] that there is an isomorphism

$$A^k(\Sigma) \longrightarrow \text{MW}_{n-k}(\Sigma), \quad \xi \longmapsto (\sigma \longmapsto \deg \xi \cdot x_\sigma),$$

where n is the dimension of Σ and “deg” is the degree map of the complete toric variety of Σ . In Theorem 6.19, we show that there is an isomorphism for the Bergman fan

$$A^k(\Sigma_M) \longrightarrow \text{MW}_{r-k}(\Sigma_M), \quad \xi \longmapsto (\sigma_{\mathcal{F}} \longmapsto \deg \xi \cdot x_{\mathcal{F}}),$$

where r is the dimension of Σ_M and “deg” is a homomorphism constructed from M . These isomorphisms are analogues of the Poincaré duality homomorphism in algebraic topology.

Definition 5.9. The *degree map* of M is the homomorphism obtained by taking the cap product

$$\deg : A^r(\Sigma_M) \longrightarrow \mathbb{Z}, \quad \xi \longmapsto \xi \cap 1_M,$$

where $1_M = 1$ is the constant r -dimensional Minkowski weight on Σ_M .

By Proposition 5.5, the homomorphism \deg is uniquely determined by its property

$$\deg(x_{\mathcal{F}}) = 1 \text{ for all monomials } x_{\mathcal{F}} \text{ corresponding to an } r\text{-dimensional cone in } \Sigma_M.$$

Proposition 5.10. The degree map of M is an isomorphism.

Proof. The second part of Proposition 5.8 shows that $A^r(\Sigma_M)$ is generated by the element α_M^r , and that $\deg(\alpha_M^r) = \deg(x_{\mathcal{F}}) = 1$. \square

5.4. We remark on algebraic geometric properties of Bergman fans, working over a fixed field \mathbb{K} . For basics on toric varieties, we refer to [Ful93]. The results of this subsection will be independent from the remainder of the paper.

The main object is the smooth toric variety $X(\Sigma)$ over \mathbb{K} associated to a unimodular fan Σ in $N_{\mathbb{R}}$:

$$X(\Sigma) := \bigcup_{\sigma \in \Sigma} \text{Spec } \mathbb{K}[\sigma^{\vee} \cap M].$$

It is known that the Chow ring of Σ is naturally isomorphic to the Chow ring of $X(\Sigma)$:

$$A^*(\Sigma) \longrightarrow A^*(X(\Sigma)), \quad x_{\sigma} \longmapsto [X(\text{star}(\sigma))].$$

See [Dan78, Section 10] for the proof when Σ is complete, and see [BDP90] and [Bri96] for the general case.

Definition 5.11. A morphism between smooth algebraic varieties $X_1 \rightarrow X_2$ is a *Chow equivalence* if the induced homomorphism between the Chow rings $A^*(X_2) \rightarrow A^*(X_1)$ is an isomorphism.

In fact, the results of this subsection will be valid for any variety that is locally a quotient of a manifold by a finite group so that $A^*(X) \otimes_{\mathbb{Z}} \mathbb{Q}$ has the structure of a graded algebra over \mathbb{Q} . Matroids provide nontrivial examples of Chow equivalences. For example, consider the subfan $\tilde{\Sigma}_{M, \mathcal{P}} \subseteq \Sigma_{M, \mathcal{P}}$ and the corresponding open subset

$$X(\tilde{\Sigma}_{M, \mathcal{P}}) \subseteq X(\Sigma_{M, \mathcal{P}}).$$

In Proposition 6.2, we show that the above inclusion is a Chow equivalence for any M and \mathcal{P} .

We remark that, when $\mathbb{K} = \mathbb{C}$, a Chow equivalence need not induce an isomorphism between singular cohomology rings. For example, consider any line in a projective plane minus two points

$$\mathbb{C}P^1 \subseteq \mathbb{C}P^2 \setminus \{p_1, p_2\}.$$

The inclusion is a Chow equivalence for any two distinct points p_1, p_2 outside $\mathbb{C}P^1$, but the two spaces have different singular cohomology rings.

We show that the notion of Chow equivalence can be used to characterize the realizability of matroids.

Theorem 5.12. There is a Chow equivalence from a smooth projective variety over \mathbb{K} to $X(\Sigma_M)$ if and only if the matroid M is realizable over \mathbb{K} .

Proof. This is a classical variant of the tropical characterization of the realizability of matroids in [KP11]. We write r for the dimension of Σ_M , and n for the dimension of $X(\Sigma_M)$. As before, the ground set of M will be $E = \{0, 1, \dots, n\}$.

The “if” direction follows from the construction of De Concini-Procesi wonderful models [DP95]. Suppose that the loopless matroid M is realized by a spanning set of nonzero vectors

$$\mathcal{R} = \{f_0, f_1, \dots, f_n\} \subseteq V/\mathbb{K}.$$

The realization \mathcal{R} gives an injective linear map between two projective spaces

$$L_{\mathcal{R}} : \mathbb{P}(V^{\vee}) \longrightarrow X(\Sigma_{\emptyset}), \quad L_{\mathcal{R}} = [f_0 : f_1 : \cdots : f_n],$$

where Σ_{\emptyset} is the complete fan in $N_{E, \mathbb{R}}$ corresponding to the empty order filter of $\mathcal{P}(E)$. Note that the normal fan of the n -dimensional permutohedron $\Sigma_{\mathcal{P}(E)}$ can be obtained from the normal fan of the n -dimensional simplex Σ_{\emptyset} by performing a sequence of stellar subdivisions. In other words, there is a morphism between toric varieties

$$\pi : X(\Sigma_{\mathcal{P}(E)}) \longrightarrow X(\Sigma_{\emptyset}),$$

which is the composition of blowups of torus-invariant subvarieties. To be explicit, consider a sequence of order filters of $\mathcal{P}(E)$ obtained by adding a single subset at a time:

$$\emptyset, \dots, \mathcal{P}_-, \mathcal{P}_+, \dots, \mathcal{P}(E) \quad \text{with} \quad \mathcal{P}_+ = \mathcal{P}_- \cup \{Z\}.$$

The corresponding sequence of Σ interpolates between the collections Σ_{\emptyset} and $\Sigma_{\mathcal{P}(E)}$:

$$\Sigma_{\emptyset} \rightsquigarrow \dots \rightsquigarrow \Sigma_{\mathcal{P}_-} \rightsquigarrow \Sigma_{\mathcal{P}_+} \rightsquigarrow \dots \rightsquigarrow \Sigma_{\mathcal{P}(E)}.$$

The modification in the middle replaces the cones of the form $\sigma_{Z < \mathcal{F}}$ with the sums of the form

$$\sigma_{\emptyset < \{Z\}} + \sigma_{I < \mathcal{F}},$$

where I is any proper subset of Z . The wonderful model $Y_{\mathcal{R}}$ associated to \mathcal{R} is by definition the strict transform of $\mathbb{P}(V^{\vee})$ under the composition of toric blowups π . The torus-invariant prime divisors of $X(\Sigma_{\mathcal{P}(E)})$ correspond to nonempty proper subsets of E , and those divisors intersecting $Y_{\mathcal{R}}$ exactly correspond to nonempty proper flats of M . Therefore, the smooth projective variety $Y_{\mathcal{R}}$ is contained in the open subset

$$X(\Sigma_M) \subseteq X(\Sigma_{\mathcal{P}(E)}).$$

The inclusion $Y_{\mathcal{R}} \subseteq X(\Sigma_M)$ is a Chow equivalence [FY04, Corollary 2].

The “only if” direction follows from computations in $A^*(\Sigma_M)$ made in the previous subsection. Suppose that there is a Chow equivalence from a smooth projective variety

$$f : Y \longrightarrow X(\Sigma_M).$$

Proposition 5.5 and Proposition 5.10 show that

$$A^r(Y) \simeq A^r(\Sigma_M) \simeq \mathbb{Z} \quad \text{and} \quad A^k(Y) \simeq A^k(\Sigma_M) \simeq 0 \quad \text{for all } k \text{ larger than } r.$$

Since Y is complete, the above implies that the dimension of Y is r . Let g be the composition

$$Y \xrightarrow{f} X(\Sigma_M) \xrightarrow{\pi_M} X(\Sigma_{\emptyset}) \simeq \mathbb{P}^n,$$

where π_M is the restriction of the composition of toric blowups π . We use Proposition 5.8 to compute the degree of the image $g(Y) \subseteq \mathbb{P}^n$.

For this we note that, for any element $i \in E$, we have

$$\pi_M^{-1}\{z_i = 0\} = \bigcup_{i \in F} D_F,$$

where z_i is the homogeneous coordinate of \mathbb{P}^n corresponding to i and D_F is the torus-invariant prime divisor of $X(\Sigma_M)$ corresponding to a nonempty proper flat F . All the components of $\pi_M^{-1}\{z_i = 0\}$ appear with multiplicity 1, and hence

$$\pi_M^* \mathcal{O}_{\mathbb{P}^n}(1) = \alpha_M \in A^1(\Sigma_M).$$

Hence, under the isomorphism f^* between the Chow rings, the 0-dimensional cycle $(g^* \mathcal{O}_{\mathbb{P}^n}(1))^r$ is the image of the generator

$$(\pi_M^* \mathcal{O}_{\mathbb{P}^n}(1))^r = \alpha_M^r \in A^r(\Sigma_M) \simeq \mathbb{Z}.$$

By the projection formula, the above implies that the degree of the image of Y in \mathbb{P}^n is 1. In other words, $g(Y) \subseteq \mathbb{P}^n$ is an r -dimensional linear subspace defined over \mathbb{K} . We express the inclusion in the form

$$L_{\mathcal{R}} : \mathbb{P}(V^\vee) \longrightarrow \mathbb{P}^n, \quad L_{\mathcal{R}} = [f_0 : f_1 : \cdots : f_n].$$

Let M' be the loopless matroid on E defined by the set of nonzero vectors $\mathcal{R} \subseteq V/\mathbb{K}$. The image of Y in $X(\Sigma_M)$ is the wonderful model $Y_{\mathcal{R}}$, and hence

$$X(\Sigma_{M'}) \subseteq X(\Sigma_M).$$

Observe that none of the torus-invariant prime divisors of $X(\Sigma_M)$ are rationally equivalent to zero. Since f is a Chow equivalence, the observation implies that the torus-invariant prime divisors of $X(\Sigma_{M'})$ and $X(\Sigma_M)$ bijectively correspond to each other. Since a matroid is determined by its set of nonempty proper flats, this shows that $M = M'$. \square

6. POINCARÉ DUALITY FOR MATROIDS

6.1. The principal result of this section is an analogue of Poincaré duality for $A^*(\Sigma_{M, \mathcal{P}})$, see Theorem 6.19. We give an alternative description of the Chow ring suitable for this purpose.

Definition 6.1. Let $S_{E \cup \mathcal{P}}$ be the polynomial ring over \mathbb{Z} with variables indexed by $E \cup \mathcal{P}$:

$$S_{E \cup \mathcal{P}} := \mathbb{Z}[x_i, x_F]_{i \in E, F \in \mathcal{P}}.$$

The *Chow ring* of (M, \mathcal{P}) is the commutative graded algebra

$$A^*(M, \mathcal{P}) := S_{E \cup \mathcal{P}} / (\mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3 + \mathcal{I}_4),$$

where $\mathcal{I}_1, \mathcal{I}_2, \mathcal{I}_3, \mathcal{I}_4$ are the ideals of $S_{E \cup \mathcal{P}}$ defined below.

(Incomparability relations) The ideal \mathcal{I}_1 is generated by the quadratic monomials

$$x_{F_1} x_{F_2},$$

where F_1 and F_2 are two incomparable flats in the order filter \mathcal{P} .

(Complement relations) The ideal \mathcal{I}_2 is generated by the quadratic monomials

$$x_i x_F,$$

where F is a flat in the order filter \mathcal{P} and i is an element in the complement $E \setminus F$.

(Closure relations) The ideal \mathcal{S}_3 is generated by the monomials

$$\prod_{i \in I} x_i,$$

where I is an independent set of M whose closure is in $\mathcal{P} \cup \{E\}$.

(Linear relations) The ideal \mathcal{S}_4 is generated by the linear forms

$$\left(x_i + \sum_{i \in F} x_F\right) - \left(x_j + \sum_{j \in F} x_F\right),$$

where i and j are distinct elements of E and the sums are over flats F in \mathcal{P} .

When $\mathcal{P} = \mathcal{P}(M)$, we omit \mathcal{P} from the notation and write the Chow ring by $A^*(M)$.

When \mathcal{P} is empty, the relations in \mathcal{S}_4 show that all x_i are equal in the Chow ring, and hence

$$A^*(M, \emptyset) \simeq \mathbb{Z}[x]/(x^{r+1}).$$

When \mathcal{P} is $\mathcal{P}(M)$, the relations in \mathcal{S}_3 show that all x_i are zero in the Chow ring, and hence

$$A^*(M) \simeq A^*(\Sigma_M).$$

In general, if i is an element whose closure is in \mathcal{P} , then x_i is zero in the Chow ring. The square-free monomial relations in the remaining set of variables bijectively correspond to the non-faces of the Bergman complex $\Delta_{M, \mathcal{P}}$, and hence

$$A^*(M, \mathcal{P}) \simeq A^*(\Sigma_{M, \mathcal{P}}).$$

We show that the Chow ring of (M, \mathcal{P}) is also isomorphic to the Chow ring of the reduced Bergman fan $\tilde{\Sigma}_{M, \mathcal{P}}$.

Proposition 6.2. Let I be a subset of E , and let F be a flat in an order filter \mathcal{P} of $\mathcal{P}(M)$.

(1) If I has cardinality at least the rank of F , then

$$\left(\prod_{i \in I} x_i\right) x_F = 0 \in A^*(M, \mathcal{P}).$$

(2) If I has cardinality at least $r + 1$, then

$$\prod_{i \in I} x_i = 0 \in A^*(M, \mathcal{P}).$$

In other words, the inclusion of the open subset $X(\tilde{\Sigma}_{M, \mathcal{P}}) \subseteq X(\Sigma_{M, \mathcal{P}})$ is a Chow equivalence. Since the reduced Bergman fan has dimension r , this implies that

$$A^k(M, \mathcal{P}) = 0 \text{ for } k > r.$$

Proof. For the first assertion, we use complement relations in \mathcal{S}_2 to reduce to the case when $I \subseteq F$. We prove by induction on the difference between the rank of F and the rank of I .

When the difference is zero, I contains a basis of F , and the desired vanishing follows from a closure relation in \mathcal{S}_3 . When the difference is positive, we choose a subset $J \subseteq F$ with

$$\mathrm{rk}(J) = \mathrm{rk}(I) + 1, \quad I \setminus J = \{i\} \quad \text{and} \quad J \setminus I = \{j\}.$$

From the linear relation in \mathcal{S}_4 for i and j , we deduce that

$$x_i + \sum_{\substack{i \in G \\ j \notin G}} x_G = x_j + \sum_{\substack{j \in G \\ i \notin G}} x_G,$$

where the sums are over flats G in \mathcal{P} . Multiplying both sides by $\left(\prod_{i \in I \cap J} x_i\right) x_F$, we get

$$\left(\prod_{i \in I} x_i\right) x_F = \left(\prod_{j \in J} x_j\right) x_F.$$

Indeed, a term involving x_G in the expansions of the products is zero in the Chow ring by

- (1) an incomparability relation in \mathcal{S}_1 , if $G \not\subseteq F$,
- (2) a complement relation in \mathcal{S}_2 , if $I \cap J \not\subseteq G$,
- (3) the induction hypothesis for $I \cap J \subseteq G$, if otherwise.

The right-hand side of the equality is zero by the induction hypothesis for $J \subseteq F$.

The second assertion can be proved in the same way, by descending induction on the rank of I , using the first part of the proposition. \square

We record here that the isomorphism of Proposition 4.7 uniquely extends to an isomorphism between the corresponding Chow rings.

Proposition 6.3. The homomorphism π_{PL} induces an isomorphism of graded rings

$$\pi_{\mathrm{PL}} : A^*(M, \mathcal{P}) \longrightarrow A^*(\overline{M}, \mathcal{P}).$$

The homomorphism ι_{PL} induces the inverse isomorphism of graded rings

$$\iota_{\mathrm{PL}} : A^*(\overline{M}, \mathcal{P}) \longrightarrow A^*(M, \mathcal{P}).$$

Proof. Consider the extensions of π_{PL} and ι_{PL} to the polynomial rings

$$S_{E \cup \mathcal{P}} \begin{array}{c} \xrightarrow{\tilde{\pi}_{\mathrm{PL}}} \\ \xleftarrow{\tilde{\iota}_{\mathrm{PL}}} \end{array} S_{\overline{E} \cup \mathcal{P}}.$$

The result follows from the observation that $\tilde{\pi}_{\mathrm{PL}}$ and $\tilde{\iota}_{\mathrm{PL}}$ preserve the monomial relations in \mathcal{S}_1 , \mathcal{S}_2 , and \mathcal{S}_3 . \square

6.2. Let \mathcal{P}_- be an order filter of $\mathcal{P}(M)$, and let Z be a flat maximal in $\mathcal{P}(M) \setminus \mathcal{P}_-$. We set

$$\mathcal{P}_+ := \mathcal{P}_- \cup \{Z\} \subseteq \mathcal{P}(M).$$

The collection \mathcal{P}_+ is an order filter of $\mathcal{P}(M)$.

Definition 6.4. The *matroidal flip* from \mathcal{P}_- to \mathcal{P}_+ is the modification of fans $\Sigma_{M, \mathcal{P}_-} \rightsquigarrow \Sigma_{M, \mathcal{P}_+}$.

The flat Z will be called the *center* of the matroidal flip. The matroidal flip removes the cones

$$\sigma_{I < \mathcal{F}} \text{ with } \text{cl}_M(I) = Z \text{ and } \min \mathcal{F} \neq Z,$$

and replaces them with the cones

$$\sigma_{I < \mathcal{F}} \text{ with } \text{cl}_M(I) \neq Z \text{ and } \min \mathcal{F} = Z.$$

The center Z is necessarily minimal in \mathcal{P}_+ , and we have

$$\begin{aligned} \text{star}(\sigma_{Z < \emptyset}, \Sigma_{M, \mathcal{P}_-}) &\simeq \Sigma_{M_Z}, \\ \text{star}(\sigma_{\emptyset < \{Z\}}, \Sigma_{M, \mathcal{P}_+}) &\simeq \Sigma_{M^Z, \emptyset} \times \Sigma_{M_Z}. \end{aligned}$$

Remark 6.5. The matroidal flip preserves the homotopy type of the underlying simplicial complexes $\Delta_{M, \mathcal{P}_-}$ and $\Delta_{M, \mathcal{P}_+}$. To see this, consider the inclusion

$$\Delta_{M, \mathcal{P}_+} \subseteq \Delta_{M, \mathcal{P}_-}^* := \text{the stellar subdivision of } \Delta_{M, \mathcal{P}_-} \text{ relative to } \Delta_{Z < \emptyset}.$$

We claim that the left-hand side is a deformation retract of the right-hand side. More precisely, there is a sequence of compositions of elementary collapses

$$\begin{aligned} \Delta_{M, \mathcal{P}_+}^* &= \Delta_{M, \mathcal{P}_-}^{1,1} \rightsquigarrow \Delta_{M, \mathcal{P}_-}^{1,2} \rightsquigarrow \dots \rightsquigarrow \Delta_{M, \mathcal{P}_-}^{1, \text{crk}(Z)-1} \rightsquigarrow \\ &\Delta_{M, \mathcal{P}_-}^{1, \text{crk}(Z)} = \Delta_{M, \mathcal{P}_-}^{2,1} \rightsquigarrow \Delta_{M, \mathcal{P}_-}^{2,2} \rightsquigarrow \dots \rightsquigarrow \Delta_{M, \mathcal{P}_-}^{2, \text{crk}(Z)-1} \rightsquigarrow \\ &\Delta_{M, \mathcal{P}_-}^{2, \text{crk}(Z)} = \Delta_{M, \mathcal{P}_-}^{3,1} \rightsquigarrow \Delta_{M, \mathcal{P}_-}^{3,2} \rightsquigarrow \dots \rightsquigarrow \Delta_{M, \mathcal{P}_-}^{3, \text{crk}(Z)-1} \rightsquigarrow \dots \rightsquigarrow \Delta_{M, \mathcal{P}_+}, \end{aligned}$$

where $\Delta_{M, \mathcal{P}_-}^{m, k+1}$ is the subcomplex of $\Delta_{M, \mathcal{P}_-}^{m, k}$ obtained by collapsing all the faces $\Delta_{I < \mathcal{F}}$ with

$$\text{cl}_M(I) = Z, \quad \min \mathcal{F} \neq Z, \quad |Z \setminus I| = m, \quad |\mathcal{F}| = \text{crk}_M(Z) - k.$$

The faces $\Delta_{I < \mathcal{F}}$ satisfying the above conditions can be collapsed in $\Delta_{M, \mathcal{P}_-}^{m, k}$ because

$$\text{link}(\Delta_{I < \mathcal{F}}, \Delta_{M, \mathcal{P}_-}^{m, k}) = \{\mathbf{e}_Z\}.$$

It follows that the homotopy type of the Bergman complex $\Delta_{M, \mathcal{P}}$ is independent of \mathcal{P} . For basics of elementary collapses of simplicial complexes, see [Koz08, Chapter 6]. The special case that $\Delta_{M, \emptyset}$ is homotopic to Δ_M is known in combinatorial topology as the crosscut theorem, see for example [Koz08, Chapter 13].

We construct homomorphisms associated to the matroidal flip, the *pullback homomorphism* and the *Gysin homomorphism*.

Proposition 6.6. There is a graded ring homomorphism between the Chow rings

$$\Phi_Z : A^*(M, \mathcal{P}_-) \longrightarrow A^*(M, \mathcal{P}_+)$$

uniquely determined by the property

$$x_F \longmapsto x_F \quad \text{and} \quad x_i \longmapsto \begin{cases} x_i + x_Z & \text{if } i \in Z, \\ x_i & \text{if } i \notin Z. \end{cases}$$

We call this map the *pullback homomorphism* associated to the matroidal flip from \mathcal{P}_- to \mathcal{P}_+ . The pullback homomorphism will later shown to be injective, see Theorem 6.18.

Proof. Consider the homomorphism between the polynomial rings

$$\phi_Z : S_{E \cup \mathcal{P}_-} \longrightarrow S_{E \cup \mathcal{P}_+}$$

defined by the same rule determining Φ_Z . We claim that

$$\phi_Z(\mathcal{I}_1) \subseteq \mathcal{I}_1, \quad \phi_Z(\mathcal{I}_2) \subseteq \mathcal{I}_1 + \mathcal{I}_2, \quad \phi_Z(\mathcal{I}_3) \subseteq \mathcal{I}_2 + \mathcal{I}_3, \quad \phi_Z(\mathcal{I}_4) \subseteq \mathcal{I}_4.$$

The first and the last inclusions are straightforward to verify.

We check the second inclusion. For an element i in $E \setminus F$, we have

$$\phi_Z(x_i x_F) = \begin{cases} x_i x_F + x_Z x_F & \text{if } i \in Z, \\ x_i x_F & \text{if } i \notin Z. \end{cases}$$

If i is in $Z \setminus F$, then the monomial $x_Z x_F$ is in \mathcal{I}_1 because Z is minimal in \mathcal{P}_+ .

We check the third inclusion. For an independent set I whose closure is in $\mathcal{P}_- \cup \{E\}$,

$$\phi_Z\left(\prod_{i \in I} x_i\right) = \prod_{i \in I \setminus Z} x_i \prod_{i \in I \cap Z} (x_i + x_Z).$$

The term $\prod_{i \in I} x_i$ in the expansion of the right-hand side is in \mathcal{I}_3 . Since Z is minimal in \mathcal{P}_+ , there is an element in $I \setminus Z$, and hence all the remaining terms in the expansion are in \mathcal{I}_2 . \square

Proposition 6.7. The pullback homomorphism Φ_Z is an isomorphism when $\text{rk}_M(Z) = 1$.

Proof. Let j_1 and j_2 be distinct elements of Z . If Z has rank 1, then a flat contains j_1 if and only if it contains j_2 . It follows from the linear relation in $S_{E \cup \mathcal{P}_-}$ for j_1 and j_2 that

$$x_{j_1} = x_{j_2} \in A^*(M, \mathcal{P}_-).$$

We choose an element $j \in Z$, and construct the inverse Φ'_Z of Φ_Z by setting

$$x_Z \longmapsto x_j, \quad x_F \longmapsto x_F, \quad \text{and} \quad x_i \longmapsto \begin{cases} 0 & \text{if } i \in Z, \\ x_i & \text{if } i \notin Z. \end{cases}$$

It is straightforward to check that Φ'_Z is well-defined, and that $\Phi'_Z = \Phi_Z^{-1}$. \square

As before, we identify the flats of M_Z with the flats of M containing Z , and identify the flats of M^Z with the flats of M contained in Z .

Proposition 6.8. Let p and q be positive integers.

(1) There is a group homomorphism

$$\Psi_Z^{p,q} : A^{q-p}(M_Z) \longrightarrow A^q(M, \mathcal{P}_+)$$

uniquely determined by the property $x_{\mathcal{F}} \mapsto x_Z^p x_{\mathcal{F}}$.

(2) There is a group homomorphism

$$\Gamma_Z^{p,q} : A^{q-p}(M^Z) \longrightarrow A^q(M)$$

uniquely determined by the property $x_{\mathcal{F}} \mapsto x_Z^p x_{\mathcal{F}}$.

We call the map $\Psi_Z^{p,q}$ the *Gysin homomorphism* of type p, q associated to the matroidal flip from \mathcal{P}_- to \mathcal{P}_+ . The Gysin homomorphism will later shown to be injective when $p < \text{rk}_M(Z)$, see Theorem 6.18.

Proof. It is clear that the Gysin homomorphism $\Psi_Z^{p,q}$ respects the incomparability relations. We check that $\Psi_Z^{p,q}$ respects the linear relations.

Let i_1 and i_2 be elements in $E \setminus Z$, and consider the linear relation in $S_{E \cup \mathcal{P}_+}$ for i_1 and i_2 :

$$\left(x_{i_1} + \sum_{i_1 \in F} x_F \right) - \left(x_{i_2} + \sum_{i_2 \in F} x_F \right) \in \mathcal{I}_4.$$

Since i_1 and i_2 are not in Z , multiplying the linear relation with x_Z^p gives

$$x_Z^p \left(\sum_{Z \cup \{i_1\} \subseteq F} x_F - \sum_{Z \cup \{i_2\} \subseteq F} x_F \right) \in \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_4.$$

The second statement on $\Gamma_Z^{p,q}$ can be proved in the same way, using i_1 and i_2 in Z . \square

Let \mathcal{P} be any order filter of $\mathcal{P}(M)$. We choose a sequence of order filters of the form

$$\emptyset, \mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}, \dots, \mathcal{P}(M),$$

where an order filter in the sequence is obtained from the preceding one by adding a single flat. The corresponding sequence of matroidal flips interpolates between $\Sigma_{M, \emptyset}$ and Σ_M :

$$\Sigma_{M, \emptyset} \rightsquigarrow \Sigma_{M, \mathcal{P}_1} \rightsquigarrow \dots \rightsquigarrow \Sigma_{M, \mathcal{P}} \rightsquigarrow \dots \rightsquigarrow \Sigma_M.$$

Definition 6.9. We write $\Phi_{\mathcal{P}}$ and $\Phi_{\mathcal{P}^c}$ for the compositions of pullback homomorphisms

$$\Phi_{\mathcal{P}} : A^*(M, \emptyset) \longrightarrow A^*(M, \mathcal{P}) \quad \text{and} \quad \Phi_{\mathcal{P}^c} : A^*(M, \mathcal{P}) \longrightarrow A^*(M).$$

Note that $\Phi_{\mathcal{P}}$ and $\Phi_{\mathcal{P}^c}$ depend only on \mathcal{P} and not on the chosen sequence of matroidal flips. The composition of all the pullback homomorphisms $\Phi_{\mathcal{P}^c} \circ \Phi_{\mathcal{P}}$ is uniquely determined by its property

$$\Phi_{\mathcal{P}^c} \circ \Phi_{\mathcal{P}}(x_i) = \alpha_M.$$

6.3. Let \mathcal{P}_- and \mathcal{P}_+ be as before, and let Z be the center of the matroidal flip from \mathcal{P}_- to \mathcal{P}_+ . For positive integers p and q , we consider the pullback homomorphism in degree q

$$\Phi_Z^q : A^q(M, \mathcal{P}_-) \longrightarrow A^q(M, \mathcal{P}_+)$$

and the Gysin homomorphism of type p, q

$$\Psi_Z^{p,q} : A^{q-p}(M_Z) \longrightarrow A^q(M, \mathcal{P}_+).$$

Proposition 6.10. For any positive integer q , the sum of the pullback homomorphism and Gysin homomorphisms

$$\Phi_Z^q \oplus \bigoplus_{p=1}^{\text{rk}(Z)-1} \Psi_Z^{p,q}$$

is a surjective group homomorphism.

The proof is given below Lemma 6.16. In Theorem 6.18, we will show that the sum is in fact an isomorphism.

Corollary 6.11. The pullback homomorphism Φ_Z is an isomorphism in degree r :

$$\Phi_Z^r : A^r(M, \mathcal{P}_-) \simeq A^r(M, \mathcal{P}_+).$$

Repeated application of the corollary shows that, for any order filter \mathcal{P} , the homomorphisms $\Phi_{\mathcal{P}}$ and $\Phi_{\mathcal{P}^c}$ are isomorphisms in degree r :

$$\Phi_{\mathcal{P}}^r : A^r(M, \emptyset) \simeq A^r(M, \mathcal{P}) \quad \text{and} \quad \Phi_{\mathcal{P}^c}^r : A^r(M, \mathcal{P}) \simeq A^r(M).$$

Proof of Corollary 6.11. The contracted matroid M_Z has rank $\text{crk}_M(Z)$, and hence

$$\Psi_Z^{p,q} = 0 \quad \text{when } p < \text{rk}_M(Z) \text{ and } q = r.$$

Therefore, Proposition 6.10 for $q = r$ says that the homomorphism Φ_Z is surjective in degree r .

Choose a sequence of matroidal flips

$$\Sigma_{M, \emptyset} \rightsquigarrow \dots \rightsquigarrow \Sigma_{M, \mathcal{P}_-} \rightsquigarrow \Sigma_{M, \mathcal{P}_+} \rightsquigarrow \dots \rightsquigarrow \Sigma_M,$$

and consider the corresponding group homomorphisms

$$A^r(M, \emptyset) \xrightarrow{\Phi_{\mathcal{P}_-}} A^r(M, \mathcal{P}_-) \xrightarrow{\Phi_{\mathcal{P}_Z}} A^r(M, \mathcal{P}_+) \xrightarrow{\Phi_{\mathcal{P}_+^c}} A^r(M).$$

Proposition 6.10 applied to each matroidal flip in the sequence shows that all three homomorphisms are surjective. The first group is clearly isomorphic to \mathbb{Z} , and by Proposition 5.10, the last group is also isomorphic to \mathbb{Z} . It follows that all three homomorphisms are isomorphisms. \square

Let β_{M_Z} be the element β in Definition 5.7 for the contracted matroid M_Z . The first part of Proposition 6.8 shows that the expression $x_Z \beta_{M_Z}$ defines an element in $A^*(M, \mathcal{P}_+)$.

Lemma 6.12. For any element i in Z , we have

$$x_i x_Z + x_Z^2 + x_Z \beta_{M_Z} = 0 \in A^*(M, \mathcal{P}_+).$$

Proof. We choose an element j in $E \setminus Z$, and consider the linear relation in $S_{E \cup \mathcal{P}_+}$ for i and j :

$$\left(x_i + \sum_{\substack{i \in F \\ j \notin F}} x_F \right) - \left(x_j + \sum_{\substack{j \in F \\ i \notin F}} x_F \right) \in \mathcal{I}_4.$$

Since i is in Z and Z is minimal in \mathcal{P}_+ , multiplying the linear relation with x_Z gives

$$x_Z x_i + x_Z^2 + \left(\sum_{Z \subsetneq F \subsetneq F \cup \{j\}} x_Z x_F \right) \in \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_4.$$

The sum in the parenthesis is the image of β_{M_Z} under the homomorphism $\Psi_Z^{1,2}$. \square

Let α_{M_Z} be the element α in Definition 5.7 for the restricted matroid M^Z . The second part of Proposition 6.8 shows that the expression $x_Z \alpha_{M_Z}$ defines an element in $A^*(M)$.

Lemma 6.13. If Z is maximal among flats strictly contained in a proper flat \tilde{Z} , then

$$x_Z x_{\tilde{Z}} (x_Z + \alpha_{M_Z}) = 0 \in A^*(M).$$

If Z is maximal among flats strictly contained in the flat E , then

$$x_Z (x_Z + \alpha_{M_Z}) = 0 \in A^*(M).$$

Proof. We justify the first statement; the second statement can be proved in the same way.

Choose an element i in Z and an element j in $\tilde{Z} \setminus Z$. The linear relation for i and j shows that

$$\sum_{\substack{i \in F \\ j \notin F}} x_F = \sum_{\substack{j \in F \\ i \notin F}} x_F \in A^*(M).$$

Multiplying both sides by the monomial $x_Z x_{\tilde{Z}}$, the incomparability relations give

$$x_Z^2 x_{\tilde{Z}} + \left(\sum_{i \in F \subsetneq Z} x_F x_Z \right) x_{\tilde{Z}} = 0 \in A^*(M).$$

The sum in the parenthesis is the image of α_{M_Z} under the homomorphism $\Gamma_Z^{1,2}$. \square

Lemma 6.14. The sum of the images of Gysin homomorphisms is the ideal generated by x_Z :

$$\sum_{p>0} \sum_{q>0} \text{im } \Psi_Z^{p,q} = x_Z A^*(M, \mathcal{P}_+).$$

Proof. It is enough to prove that the right-hand side is contained in the left-hand side. Since Z is minimal in \mathcal{P}_+ , the incomparability relations in \mathcal{I}_1 and the complement relations in \mathcal{I}_2 show that any nonzero degree q monomial in the ideal generated by x_Z is of the form

$$x_Z^k \prod_{F \in \mathcal{F}} x_F^{k_F} \prod_{i \in I} x_i^{k_i}, \quad I \subseteq Z < \mathcal{F},$$

where the sum of the exponents is q . Since the exponent k of x_Z is positive, Lemma 6.12 shows that this monomial is in the sum

$$\operatorname{im} \Psi_Z^{k,q} + \operatorname{im} \Psi_Z^{k+1,q} + \cdots + \operatorname{im} \Psi_Z^{q,q}. \quad \square$$

Lemma 6.15. For positive integers p and q , we have

$$x_Z \operatorname{im} \Phi_Z^q \subseteq \operatorname{im} \Psi_Z^{1,q+1} \quad \text{and} \quad x_Z \operatorname{im} \Psi_Z^{p,q} \subseteq \operatorname{im} \Psi_Z^{p+1,q+1}.$$

If F is a proper flat strictly containing Z , then

$$x_F \operatorname{im} \Phi_Z^q \subseteq \operatorname{im} \Phi_Z^{q+1} \quad \text{and} \quad x_F \operatorname{im} \Psi_Z^{p,q} \subseteq \operatorname{im} \Psi_Z^{p,q+1}.$$

Proof. Only the first inclusion is nontrivial. Note that the left-hand side is generated by elements of the form

$$\xi = x_Z \prod_{F \in \mathcal{F}} x_F^{k_F} \prod_{i \in I \setminus Z} x_i^{k_i} \prod_{i \in I \cap Z} (x_i + x_Z)^{k_i},$$

where I is a subset of E and \mathcal{F} is a flag in \mathcal{P}_- . When I is contained in Z , Lemma 6.12 shows that

$$\xi = x_Z \prod_{F \in \mathcal{F}} x_F^{k_F} \prod_{i \in I} (-\beta_{M_Z})^{k_i} \in \operatorname{im} \Psi_Z^{1,q+1}.$$

When I is not contained in Z , a complement relation in $S_{E \cup \mathcal{P}_+}$ shows that $\xi = 0$. \square

Lemma 6.16. For any integers $k \geq \operatorname{rk}_M(Z)$ and $q \geq k$, we have

$$\operatorname{im} \Psi_Z^{k,q} \subseteq \operatorname{im} \Phi_Z^q + \sum_{p=1}^{k-1} \operatorname{im} \Psi_Z^{p,q}.$$

Proof. By the second statement of Lemma 6.15, it is enough to prove the assertion when $q = k$: The general case can be deduced by multiplying both sides of the inclusion by $x_{\mathcal{F}}$ for $Z < \mathcal{F}$.

By the first statement of Lemma 6.15, it is enough to justify the above when $k = \operatorname{rk}_M(Z)$: The general case can be deduced by multiplying both sides of the inclusion by powers of x_Z .

We prove the assertion when $k = q = \operatorname{rk}_M(Z)$. For this we choose a basis I of Z , and expand the product

$$\prod_{i \in I} (x_i + x_Z) \in \operatorname{im} \Phi_Z^k.$$

The closure relation for I shows that the term $\prod_{i \in I} x_i$ in the expansion is zero, and hence, by Lemma 6.12,

$$\prod_{i \in I} (x_i + x_Z) = (-\beta_{M_Z})^k - (-x_Z - \beta_{M_Z})^k \in \operatorname{im} \Phi_Z^k.$$

Expanding the right-hand side, we see that

$$x_Z^k \in \operatorname{im} \Phi_Z^k + \sum_{p=1}^{k-1} \operatorname{im} \Psi_Z^{p,k}.$$

Since $\operatorname{im} \Psi_Z^{k,k}$ is generated by x_Z^k , this implies the asserted inclusion. \square

Proof of Proposition 6.10. By Lemma 6.16, it is enough to show that the sum $\Phi_Z^q \oplus \bigoplus_{p=1}^q \Psi_Z^{p,q}$ is surjective. By Lemma 6.14, the image of the second summand is the degree q part of the ideal generated by x_Z .

We show that any monomial is in the image of the pullback homomorphism Φ_Z modulo the ideal generated by x_Z . Note that any degree q monomial not in the ideal generated by x_Z is of the form

$$\prod_{F \in \mathcal{F}} x_F^{k_F} \prod_{i \in I} x_i^{k_i}, \quad Z \notin \mathcal{F}.$$

Modulo the ideal generated by x_Z , this monomial is equal to

$$\Phi_Z \left(\prod_{F \in \mathcal{F}} x_F^{k_F} \prod_{i \in I} x_i^{k_i} \right) = \prod_{F \in \mathcal{F}} x_F^{k_F} \prod_{i \in I \setminus Z} x_i^{k_i} \prod_{i \in I \cap Z} (x_i + x_Z)^{k_i}. \quad \square$$

We use Proposition 6.10 to show that the Gysin homomorphism between top degrees is an isomorphism.

Proposition 6.17. The Gysin homomorphism $\Psi_Z^{p,q}$ is an isomorphism when $p = \text{rk}(Z)$ and $q = r$:

$$\Psi_Z^{p,q} : A^{\text{crk}(Z)-1}(M_Z) \simeq A^r(M, \mathcal{P}_+).$$

Proof. We consider the composition

$$A^{\text{crk}(Z)-1}(M_Z) \xrightarrow{\Psi_Z^{p,q}} A^r(M, \mathcal{P}_+) \xrightarrow{\Phi_{\mathcal{P}_+}} A^r(M), \quad x_{\mathcal{F}} \mapsto x_Z^{\text{rk}(Z)} x_{\mathcal{F}}.$$

The second map is an isomorphism by Corollary 6.11, and therefore it is enough to show that the composition is an isomorphism.

For this we choose two flags of nonempty proper flats of M :

$$\mathcal{L}_1 = \text{a flag of flats strictly contained in } Z \text{ with } |\mathcal{L}_1| = \text{rk}(Z) - 1,$$

$$\mathcal{L}_2 = \text{a flag of flats strictly containing } Z \text{ with } |\mathcal{L}_2| = \text{crk}(Z) - 1.$$

We claim that the composition maps a generator to a generator:

$$(-1)^{\text{rk}(Z)-1} x_Z^{\text{rk}(Z)} x_{\mathcal{L}_2} = x_{\mathcal{L}_1} x_Z x_{\mathcal{L}_2} \in A^*(M).$$

Indeed, the map $\Gamma_Z^{1, \text{rk}(Z)}$ applied to the second formula of Proposition 5.8 for M^Z gives

$$x_{\mathcal{L}_1} x_Z x_{\mathcal{L}_2} = (\alpha_{M^Z})^{\text{rk}(Z)-1} x_Z x_{\mathcal{L}_2} \in A^*(M),$$

and, by Lemma 6.13, the right-hand side of the above is equal to

$$(-1)^{\text{rk}(Z)-1} x_Z^{\text{rk}(Z)} x_{\mathcal{L}_2} \in A^*(M). \quad \square$$

6.4. Let \mathcal{P}_- , \mathcal{P}_+ , and Z be as before, and let \mathcal{P} be any order filter of $\mathcal{P}(M)$.

Theorem 6.18 (Decomposition). For any positive integer q , the sum of the pullback homomorphism and the Gysin homomorphisms

$$\Phi_Z^q \oplus \bigoplus_{p=1}^{\text{rk}(Z)-1} \Psi_Z^{p,q}$$

is an isomorphism.

Theorem 6.19 (Poincaré Duality). For any nonnegative integer $q \leq r$, the multiplication map

$$A^q(M, \mathcal{P}) \times A^{r-q}(M, \mathcal{P}) \longrightarrow A^r(M, \mathcal{P})$$

defines an isomorphism between groups

$$A^{r-q}(M, \mathcal{P}) \simeq \text{Hom}_{\mathbb{Z}}(A^q(M, \mathcal{P}), A^r(M, \mathcal{P})).$$

In particular, the groups $A^q(M, \mathcal{P})$ are torsion free. We simultaneously prove Theorem 6.18 (Decomposition) and Theorem 6.19 (Poincaré Duality) by lexicographic induction on the rank of matroids and the cardinality of the order filters. The proof is given below Lemma 6.21.

Lemma 6.20. Let q_1 and q_2 be positive integers.

(1) For any positive integer p , we have

$$\text{im } \Psi_Z^{p,q_1} \cdot \text{im } \Phi_Z^{q_2} \subseteq \text{im } \Psi_Z^{p,q_1+q_2}$$

(2) For any positive integers p_1 and p_2 , we have

$$\text{im } \Psi_Z^{p_1,q_1} \cdot \text{im } \Psi_Z^{p_2,q_2} \subseteq \text{im } \Psi_Z^{p_1+p_2,q_1+q_2}.$$

The first inclusion shows that, when $q_1 + q_2 = r$ and p is less than $\text{rk}(Z)$,

$$\text{im } \Psi_Z^{p,q_1} \cdot \text{im } \Phi_Z^{q_2} = 0.$$

The second inclusion shows that, when $q_1 + q_2 = r$ and $p_1 + p_2$ is less than $\text{rk}(Z)$,

$$\text{im } \Psi_Z^{p_1,q_1} \cdot \text{im } \Psi_Z^{p_2,q_2} = 0.$$

Proof. The assertions are direct consequences of Lemma 6.15. \square

Lemma 6.21. Let q be a positive integer, and p_1, p_2 be distinct positive integers less than $\text{rk}(Z)$.

(1) If Poincaré Duality holds for $A^*(M, \mathcal{P}_-)$, then

$$\ker \Phi_Z^q = 0 \quad \text{and} \quad \text{im } \Phi_Z^q \cap \sum_{p=1}^{\text{rk}(Z)-1} \text{im } \Psi_Z^{p,q} = 0.$$

(2) If Poincaré Duality holds for $A^*(M_Z)$, then

$$\ker \Psi_Z^{p_1,q} = \ker \Psi_Z^{p_2,q} = 0 \quad \text{and} \quad \text{im } \Psi_Z^{p_1,q} \cap \text{im } \Psi_Z^{p_2,q} = 0$$

Proof. Let ξ be a nonzero element in the domain of Φ_Z^q . Since Φ_Z is an isomorphism between top degrees, Poincaré Duality for (M, \mathcal{P}_-) implies that

$$\Phi_Z(\xi) \cdot \text{im } \Phi_Z^{r-q} \neq 0.$$

This shows that Φ_Z^q is injective. On the other hand, Lemma 6.20 shows that

$$\left(\sum_{p=1}^{\text{rk}(Z)-1} \text{im } \Psi_Z^{p,q} \right) \cdot \text{im } \Phi_Z^{r-q} = 0.$$

This shows that the image of Φ_Z^q intersects the image of $\bigoplus_{p=1}^{\text{rk}(Z)-1} \Psi_Z^{p,q}$ trivially.

Let ξ be a nonzero element in the domain of $\Psi_Z^{p,q}$, where $p = p_1$ or $p = p_2$. Since Ψ_Z is an isomorphism between top degrees, Poincaré Duality for M_Z implies that

$$\Psi_Z^{p,q}(\xi) \cdot \text{im } \Psi_Z^{\text{rk}(Z)-p,r-q} \neq 0.$$

This shows that $\Psi_Z^{p,q}$ is injective. For this assertion on the intersection, we assume that $p = p_1 > p_2$. Under the assumption Lemma 6.20 shows

$$\text{im } \Psi_Z^{p_2,q} \cdot \text{im } \Psi_Z^{\text{rk}(Z)-p,r-q} = 0.$$

This shows that the image of $\Psi_Z^{p_2,q}$ intersects the image of $\Psi_Z^{p_1,q}$ trivially. \square

Proofs of Theorem 6.18 and Theorem 6.19. We simultaneously prove Decomposition and Poincaré Duality by lexicographic induction on the rank of M and the cardinality of \mathcal{P} and \mathcal{P}_- . Note that both statements are valid when $r = 1$, and Poincaré Duality holds when $q = 0$ or $q = r$. Assuming that Poincaré Duality holds for $A^*(M_Z)$, we show the implications

$$\begin{aligned} \left(\text{Poincaré Duality holds for } A^*(M, \mathcal{P}_-) \right) &\implies \\ \left(\text{Poincaré Duality holds for } A^*(M, \mathcal{P}_-) \text{ and Decomposition holds for } \mathcal{P}_- \subseteq \mathcal{P}_+ \right) & \\ \implies \left(\text{Poincaré Duality holds for } A^*(M, \mathcal{P}_+) \right). & \end{aligned}$$

The base case of the induction is provided by the isomorphism

$$A^*(M, \emptyset) \simeq \mathbb{Z}[x]/(x^{r+1}).$$

The first implication follows from Proposition 6.10 and Lemma 6.21.

We prove the second implication. Decomposition for $\mathcal{P}_- \subseteq \mathcal{P}_+$ shows that, for any positive integer $q < r$, we have

$$A^q(M, \mathcal{P}_+) = \text{im } \Phi_Z^q \oplus \text{im } \Psi_Z^{1,q} \oplus \text{im } \Psi_Z^{2,q} \oplus \dots \oplus \text{im } \Psi_Z^{\text{rk}(Z)-1,q}, \text{ and}$$

$$A^{r-q}(M, \mathcal{P}_+) = \text{im } \Phi_Z^{r-q} \oplus \text{im } \Psi_Z^{\text{rk}(Z)-1,r-q} \oplus \text{im } \Psi_Z^{\text{rk}(Z)-2,r-q} \oplus \dots \oplus \text{im } \Psi_Z^{1,r-q}.$$

By Poincaré Duality for (M, \mathcal{P}_-) and Poincaré Duality for M_Z , all the summands above are torsion free. We construct bases of the sums by choosing bases of their summands.

We use Corollary 6.11 and Proposition 6.17 to obtain isomorphisms

$$A^r(M, \mathcal{P}_-) \simeq A^r(M, \mathcal{P}_+) \simeq A^{\text{crk}(Z)-1}(M_Z) \simeq \mathbb{Z}.$$

For a positive integer $q < r$, consider the matrices of multiplications

$$\mathcal{M}_+ := \left(A^q(M, \mathcal{P}_+) \times A^{r-q}(M, \mathcal{P}_+) \longrightarrow \mathbb{Z} \right),$$

$$\mathcal{M}_- := \left(A^q(M, \mathcal{P}_-) \times A^{r-q}(M, \mathcal{P}_-) \longrightarrow \mathbb{Z} \right),$$

and, for positive integers $p < \text{rk}(Z)$,

$$\mathcal{M}_p := \left(A^{q-p}(M_Z) \times A^{r-q-\text{rk}(Z)+p}(M_Z) \longrightarrow \mathbb{Z} \right).$$

By Lemma 6.20, under the chosen bases ordered as shown above, \mathcal{M}_+ is a block upper triangular matrix with block diagonals \mathcal{M}_- and \mathcal{M}_p , up to signs. It follows from Poincaré Duality for (M, \mathcal{P}_-) and Poincaré Duality for M_Z that

$$\det \mathcal{M}_+ = \pm \det \mathcal{M}_- \times \prod_{p=1}^{\text{rk}(Z)-1} \det \mathcal{M}_p = \pm 1.$$

This proves the second implication, completing the lexicographic induction. \square

7. HARD LEFSCHETZ PROPERTY AND HODGE-RIEMANN RELATIONS

7.1. Let r be a nonnegative integer. We record basic algebraic facts concerning the Poincaré duality, the hard Lefschetz property, and the Hodge-Riemann relations.

Definition 7.1. A graded Artinian ring R^* satisfies the *Poincaré duality of dimension r* if

- (1) there are isomorphisms $R^0 \simeq \mathbb{R}$ and $R^r \simeq \mathbb{R}$,
- (2) for every integer $q > r$, we have $R^q \simeq 0$, and,
- (3) for every integer $q \leq r$, the multiplication defines an isomorphism

$$R^{r-q} \longrightarrow \text{Hom}_{\mathbb{R}}(R^q, R^r).$$

In this case, we say that R^* is a *Poincaré duality algebra of dimension r* .

In the remainder of this subsection, we suppose that R^* is a Poincaré duality algebra of dimension r . We fix an isomorphism, called the *degree map* for R^* ,

$$\text{deg} : R^r \longrightarrow \mathbb{R}.$$

Proposition 7.2. For any nonzero element x in R^d , the quotient ring

$$R^*/\text{ann}(x), \text{ where } \text{ann}(x) := \{a \in R^* \mid x \cdot a = 0\},$$

is a Poincaré duality algebra of dimension $r - d$.

By definition, the degree map for $R^*/\text{ann}(x)$ induced by x is the homomorphism

$$\deg(x \cdot -) : R^{r-d}/\text{ann}(x) \longrightarrow \mathbb{R}, \quad a + \text{ann}(x) \longmapsto \deg(x \cdot a).$$

The Poincaré duality for R^* shows that the degree map for $R^*/\text{ann}(x)$ is an isomorphism.

Proof. This is straightforward to check, see for example [MS05, Corollary I.2.3]. \square

Definition 7.3. Let ℓ be an element of R^1 , and let q be a nonnegative integer $\leq \frac{r}{2}$.

(1) The *Lefschetz operator* on R^q associated to ℓ is the linear map

$$L_\ell^q : R^q \longrightarrow R^{r-q}, \quad a \longmapsto \ell^{r-2q} a.$$

(2) The *Hodge-Riemann form* on R^q associated to ℓ is the symmetric bilinear form

$$Q_\ell^q : R^q \times R^q \longrightarrow \mathbb{R}, \quad (a_1, a_2) \longmapsto (-1)^q \deg(a_1 \cdot L_\ell^q(a_2)).$$

(3) The *primitive subspace* of R^q associated to ℓ is the subspace

$$P_\ell^q := \{a \in R^q \mid \ell \cdot L_\ell^q(a) = 0\} \subseteq R^q.$$

Definition 7.4 (Hard Lefschetz property and Hodge-Riemann relations). We say that

(1) R^* satisfies HL(ℓ) if the Lefschetz operator L_ℓ^q is an isomorphism on R^q for all $q \leq \frac{r}{2}$, and

(2) R^* satisfies HR(ℓ) if the Hodge-Riemann form Q_ℓ^q is positive definite on P_ℓ^q for all $q \leq \frac{r}{2}$.

If the Lefschetz operator L_ℓ^q is an isomorphism, then there is a decomposition

$$R^{q+1} = P_\ell^{q+1} \oplus \ell R^q.$$

Consequently, when R^* satisfies HL(ℓ), we have the *Lefschetz decomposition* of R^q for $q \leq \frac{r}{2}$:

$$R^q = P_\ell^q \oplus \ell P_\ell^{q-1} \oplus \cdots \oplus \ell^q P_\ell^0.$$

An important basic fact is that the Lefschetz decomposition of R^q is orthogonal with respect to the Hodge-Riemann form Q_ℓ^q : For nonnegative integers $q_1 < q_2 \leq q$, we have

$$Q_\ell^q(\ell^{q_1} a_1, \ell^{q_2} a_2) = (-1)^q \deg(\ell^{q_2-q_1} \ell^{r-2q_2} a_1 a_2) = 0, \quad a_1 \in P_\ell^{q-q_1}, \quad a_2 \in P_\ell^{q-q_2}.$$

Proposition 7.5. The following conditions are equivalent for $\ell \in R^1$:

(1) R^* satisfies HL(ℓ).

(2) The Hodge-Riemann form Q_ℓ^q on R^q is nondegenerate for all $q \leq \frac{r}{2}$.

Proof. The Hodge-Riemann form Q_ℓ^q on R^q is nondegenerate if and only if the composition

$$R^q \xrightarrow{L_\ell^q} R^{r-q} \longrightarrow \text{Hom}_{\mathbb{R}}(R^q, R^r)$$

is an isomorphism, where the second map is given by the multiplication in R^* . Since R^* satisfies Poincaré duality, the composition is an isomorphism if and only if L_ℓ^q is an isomorphism. \square

If $L_\ell^q(a) = 0$, then $Q_\ell^q(a, a) = 0$ and $a \in P_\ell^q$. Thus the property $\text{HR}(\ell)$ implies the property $\text{HL}(\ell)$.

Proposition 7.6. The following conditions are equivalent for $\ell \in R^1$:

- (1) R^* satisfies $\text{HR}(\ell)$.
- (2) The Hodge-Riemann form Q_ℓ^q on R^q is nondegenerate and has signature

$$\sum_{p=0}^q (-1)^{q-p} (\dim_{\mathbb{R}} R^p - \dim_{\mathbb{R}} R^{p-1}) \text{ for all } q \leq \frac{r}{2}.$$

Proof. If R^* satisfies $\text{HR}(\ell)$, then R^* satisfies $\text{HL}(\ell)$, and therefore we have the Lefschetz decomposition

$$R^q = P_\ell^q \oplus \ell P_\ell^{q-1} \oplus \cdots \oplus \ell^q P_\ell^0.$$

Note that the Lefschetz decomposition of R^q is orthogonal with respect to Q_ℓ^q , and that there is an isometry

$$(P_\ell^p, Q_\ell^p) \simeq (\ell^{q-p} P_\ell^p, (-1)^{q-p} Q_\ell^q) \text{ for every nonnegative integer } p \leq q.$$

Therefore, the condition $\text{HR}(\ell)$ implies that

$$\begin{aligned} \left(\text{signature of } Q_\ell^q \text{ on } R^q \right) &= \sum_{p=0}^q (-1)^{q-p} \left(\text{signature of } Q_\ell^p \text{ on } P_\ell^p \right) \\ &= \sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} R^p - \dim_{\mathbb{R}} R^{p-1} \right). \end{aligned}$$

Conversely, suppose that the Hodge-Riemann forms Q_ℓ^q are nondegenerate and their signatures are given by the stated formula. Proposition 7.5 shows that R^* satisfies $\text{HL}(\ell)$, and hence

$$R^q = P_\ell^q \oplus \ell P_\ell^{q-1} \oplus \cdots \oplus \ell^q P_\ell^0.$$

The Lefschetz decomposition of R^q is orthogonal with respect to Q_ℓ^q , and therefore

$$\left(\text{signature of } Q_\ell^q \text{ on } P_\ell^q \right) = \left(\text{signature of } Q_\ell^q \text{ on } R^q \right) - \left(\text{signature of } Q_\ell^{q-1} \text{ on } R^{q-1} \right).$$

The assumptions on the signatures of Q_ℓ^q and Q_ℓ^{q-1} show that the right-hand side is

$$\dim_{\mathbb{R}} R^q - \dim_{\mathbb{R}} R^{q-1} = \dim_{\mathbb{R}} P_\ell^q.$$

Since Q_ℓ^q is nondegenerate on P_ℓ^q , this means that Q_ℓ^q is positive definite on P_ℓ^q . □

7.2. In this subsection, we show that the properties HL and HR are preserved under the tensor product of Poincaré duality algebras.

Let R_1^* and R_2^* be Poincaré duality algebras of dimensions r_1 and r_2 respectively. We choose degree maps for R_1^* and for R_2^* , denoted

$$\deg_1 : R_1^{r_1} \longrightarrow \mathbb{R}, \quad \deg_2 : R_2^{r_2} \longrightarrow \mathbb{R}.$$

We note that $R_1 \otimes_{\mathbb{R}} R_2$ is a Poincaré duality algebra of dimension $r_1 + r_2$: For any two graded components of the tensor product with complementary degrees

$$\begin{aligned} & \left(R_1^p \otimes_{\mathbb{R}} R_2^0 \right) \oplus \left(R_1^{p-1} \otimes_{\mathbb{R}} R_2^1 \right) \oplus \cdots \oplus \left(R_1^0 \otimes_{\mathbb{R}} R_2^p \right), \\ & \left(R_1^q \otimes_{\mathbb{R}} R_2^0 \right) \oplus \left(R_1^{q-1} \otimes_{\mathbb{R}} R_2^1 \right) \oplus \cdots \oplus \left(R_1^0 \otimes_{\mathbb{R}} R_2^q \right), \end{aligned}$$

the multiplication of the two can be represented by a block diagonal matrix with diagonals

$$\left(R_1^{p-k} \otimes_{\mathbb{R}} R_2^k \right) \times \left(R_1^{q-r_2+k} \otimes_{\mathbb{R}} R_2^{r_2-k} \right) \longrightarrow R_1^{r_1} \otimes_{\mathbb{R}} R_2^{r_2}.$$

By definition, the *induced degree map* for the tensor product is the isomorphism

$$\deg_1 \otimes_{\mathbb{R}} \deg_2 : R_1^{r_1} \otimes_{\mathbb{R}} R_2^{r_2} \longrightarrow \mathbb{R}.$$

We use the induced degree map whenever we discuss the property HR for tensor products.

Proposition 7.7. Let ℓ_1 be an element of R_1^1 , and let ℓ_2 be an element of R_2^1 .

- (1) If R_1^* satisfies HL(ℓ_1) and R_2^* satisfies HL(ℓ_2), then $R_1^* \otimes_{\mathbb{R}} R_2^*$ satisfies HL($\ell_1 \otimes 1 + 1 \otimes \ell_2$).
- (2) If R_1^* satisfies HR(ℓ_1) and R_2^* satisfies HR(ℓ_2), then $R_1^* \otimes_{\mathbb{R}} R_2^*$ satisfies HR($\ell_1 \otimes 1 + 1 \otimes \ell_2$).

We begin the proof with the following special case.

Lemma 7.8. Let $r_1 \leq r_2$ be nonnegative integers, and consider the Poincaré duality algebras

$$R_1^* = \mathbb{R}[x_1]/(x_1^{r_1+1}) \quad \text{and} \quad R_2^* = \mathbb{R}[x_2]/(x_2^{r_2+1})$$

equipped with the degree maps

$$\begin{aligned} \deg_1 : R_1^{r_1} &\longrightarrow \mathbb{R}, & x_1^{r_1} &\longmapsto 1, \\ \deg_2 : R_2^{r_2} &\longrightarrow \mathbb{R}, & x_2^{r_2} &\longmapsto 1. \end{aligned}$$

Then R_1^* satisfies HR(x_1), R_2^* satisfies HR(x_2), and $R_1^* \otimes_{\mathbb{R}} R_2^*$ satisfies HR($x_1 \otimes 1 + 1 \otimes x_2$).

The first two assertions are easy to check, and the third assertion follows from the Hodge-Riemann relations for the cohomology of the compact Kähler manifold $\mathbb{C}P^{r_1} \times \mathbb{C}P^{r_2}$. Below we give a combinatorial proof using the Lindström-Gessel-Viennot lemma.

Proof. For the third assertion, we identify the tensor product with

$$R^* := \mathbb{R}[x_1, x_2]/(x_1^{r_1+1}, x_2^{r_2+1}), \quad \text{and set } \ell := x_1 + x_2.$$

The induced degree map for the tensor product will be written

$$\deg : R^{r_1+r_2} \longrightarrow \mathbb{R}, \quad x_1^{r_1} x_2^{r_2} \longmapsto 1.$$

Claim. For some (equivalently any) choice of basis of R^q , we have

$$(-1)^{\frac{q(q+1)}{2}} \det(Q_\ell^q) > 0 \text{ for all nonnegative integers } q \leq r_1.$$

We show that it is enough to prove the claim. The inequality of the claim implies that Q_ℓ^q is nondegenerate for $q \leq r_1$, and hence L_ℓ^q is an isomorphism for $q \leq r_1$. The Hilbert function of R^* forces the dimensions of the primitive subspaces to satisfy

$$\dim_{\mathbb{R}} P_\ell^q = \begin{cases} 1 & \text{for } q \leq r_1, \\ 0 & \text{for } q > r_1, \end{cases}$$

and that there is a decomposition

$$R^q = P_\ell^q \oplus \ell P_\ell^{q-1} \oplus \cdots \oplus \ell^q P_\ell^0 \text{ for } q \leq r_1.$$

Every summand of the above decomposition is 1-dimensional, and hence

$$\left(\text{signature of } Q_\ell^q \text{ on } R^q \right) = \pm 1 - \left(\text{signature of } Q_\ell^{q-1} \text{ on } R^{q-1} \right).$$

The claim on the determinant of Q_ℓ^q determines the sign of ± 1 in the above equality:

$$\left(\text{signature of } Q_\ell^q \right) = 1 - \left(\text{signature of } Q_\ell^{q-1} \right).$$

It follows that the signature of Q_ℓ^q on P_ℓ^q is 1 for $q \leq r_1$, and thus R satisfies $\text{HR}(\ell)$.

We now prove the claim on $\det(Q_\ell^q) = \det((-1)^q Q_\ell^q)$ for $q \leq r_1$. We use the monomial basis

$$\left\{ x_1^i x_2^{q-i} \mid i = 0, 1, \dots, q \right\} \subseteq R^q.$$

The matrix $[a_{ij}]$ which represents $(-1)^q Q_\ell^q$ has binomial coefficients as its entries:

$$[a_{ij}] := \left[\deg \left((x_1 + x_2)^{r_1+r_2-2q} x_1^{i+j} x_2^{q-i+q-j} \right) \right] = \left[\binom{r_1+r_2-2q}{r_1-i-j} \right].$$

We determine the sign of the determinant of $[a_{ij}]$ using the Lindström-Gessel-Viennot lemma, see [Aig07, Section 5.4] for an exposition and similar examples.

Consider the grid graph in the plane with vertices \mathbb{Z}^2 and edges directed in the positive x -directions and the positive y -directions. We place the starting points $\mathcal{P}_0, \dots, \mathcal{P}_q$ and the ending points $\mathcal{Q}_0, \dots, \mathcal{Q}_q$ on two parallel diagonal lines, \mathcal{P}_i from northwest to southeast and \mathcal{Q}_j from southeast to northwest:

$$\begin{aligned} \mathcal{P}_0 &= (-r_1, q), & \mathcal{P}_1 &= (1-r_1, q-1), & \dots & \mathcal{P}_q &= (q-r_1, 0), \\ \mathcal{Q}_q &= (-q, r_2), & \mathcal{Q}_{q-1} &= (1-q, r_2-1), & \dots & \mathcal{Q}_0 &= (0, r_2-q). \end{aligned}$$

Note that there are exactly a_{ij} distinct lattice paths from \mathcal{P}_i to \mathcal{Q}_j .

The Lindström–Gessel–Viennot lemma says that

$$\det [a_{ij}] = \sum_{\sigma} \text{sign}(\sigma),$$

where the sum is over tuples of non-intersecting lattice paths $\sigma : \{\mathcal{P}_0, \dots, \mathcal{P}_q\} \rightarrow \{\mathcal{Q}_0, \dots, \mathcal{Q}_q\}$ and $\text{sign}(\sigma)$ is the sign of the induced permutation of $\{0, 1, \dots, q\}$. In our case, any tuple of non-intersecting lattice paths σ as above should go from \mathcal{P}_i to \mathcal{Q}_{q-i} , and hence

$$\text{sign}(\sigma) = (-1)^{q(q+1)/2}.$$

It is clear that there is at least one such tuple of non-intersecting lattice paths; for example, for all \mathcal{P}_i one may first go east $r_1 - q$ times and then go north $r_2 - q$ times to arrive at \mathcal{Q}_{q-i} . This gives

$$(-1)^{q(q+1)/2} \det [a_{ij}] > 0. \quad \square$$

Now we reduce Proposition 7.7 to the case of Lemma 7.8. We first introduce some useful notions to be used in the remaining part of the proof.

Let R^* be a Poincaré duality algebra of dimension r , and let ℓ be an element of R^1 .

Definition 7.9. Let V^* be a graded subspace of R^* . We say that

- (1) V^* satisfies HL(ℓ) if Q_{ℓ}^q restricted to V^q is nondegenerate for all nonnegative $q \leq \frac{r}{2}$.
- (2) V^* satisfies HR(ℓ) if Q_{ℓ}^q restricted to V^q is nondegenerate and has signature

$$\sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} V^p - \dim_{\mathbb{R}} V^{p-1} \right) \text{ for all nonnegative } q \leq \frac{r}{2}.$$

Propositions 7.5 and 7.6 show that this agrees with the previous definition when $V^* = R^*$.

Definition 7.10. Let V_1^* and V_2^* be graded subspaces of R^* . We write

$$V_1^* \perp_{\text{PD}} V_2^*$$

to mean that $V_1^* \cap V_2^* = 0$ and $V_1^{r-q} V_2^q = 0$ for all nonnegative integers $q \leq r$, and write

$$V_1^* \perp_{Q_{\ell}^*} V_2^*$$

to mean that $V_1^* \cap V_2^* = 0$ and $Q_{\ell}^q(V_1^q, V_2^q) = 0$ for all nonnegative integers $q \leq \frac{r}{2}$.

We record here basic properties of the two notions of orthogonality. Let S^* be another Poincaré duality algebra of dimension s .

Lemma 7.11. Let $V_1^*, V_2^* \subseteq R^*$ and $W_1^*, W_2^* \subseteq S^*$ be graded subspaces.

- (1) If $V_1^* \perp_{Q_{\ell}^*} V_2^*$ and if both V_1^*, V_2^* satisfy HL(ℓ), then $V_1^* \oplus V_2^*$ satisfy HL(ℓ).
- (2) If $V_1^* \perp_{Q_{\ell}^*} V_2^*$ and if both V_1^*, V_2^* satisfy HR(ℓ), then $V_1^* \oplus V_2^*$ satisfy HR(ℓ).
- (3) If $V_1^* \perp_{\text{PD}} V_2^*$ and if $\ell V_1^* \subseteq V_1^*$, then $V_1^* \perp_{Q_{\ell}^*} V_2^*$.

(4) If $V_1^* \perp_{\text{PD}} V_2^*$, then $(V_1^* \otimes_{\mathbb{R}} W_1^*) \perp_{\text{PD}} (V_2^* \otimes_{\mathbb{R}} W_2^*)$.

Proof. The first two assertions are straightforward. We justify the third assertion: For any non-negative integer $q \leq \frac{r}{2}$, the assumption on V_1^* implies $L_\ell^q V_1^q \subseteq V_1^{r-q}$, and hence

$$Q_\ell^q(V_1^q, V_2^q) \subseteq \text{deg}(V_1^{r-q} V_2^q) = 0.$$

For the fourth assertion, we check that, for any nonnegative integers p_1, p_2, q_1, q_2 whose sum is $r + s$,

$$V_1^{p_1} V_2^{p_2} \otimes_{\mathbb{R}} W_1^{q_1} W_2^{q_2} = 0.$$

The assumption on V_1^* and V_2^* shows that the first factor is trivial if $p_1 + p_2 \geq r$, and the second factor is trivial if otherwise. \square

Proof of Proposition 7.7. Suppose that R_1^* satisfies $\text{HR}(\ell_1)$ and that R_2^* satisfies $\text{HR}(\ell_2)$. We set

$$R^* := R_1^* \otimes_{\mathbb{R}} R_2^*, \quad \ell := \ell_1 \otimes 1 + 1 \otimes \ell_2.$$

We show that R^* satisfy $\text{HR}(\ell)$. The assertion on HL can be proved in the same way.

For every $p \leq \frac{r_1}{2}$, choose an orthogonal basis of $P_{\ell_1}^p \subseteq R_1^p$ with respect to $Q_{\ell_1}^p$:

$$\{v_1^p, v_2^p, \dots, v_{m(p)}^p\} \subseteq P_{\ell_1}^p.$$

Similarly, for every $q \leq \frac{r_2}{2}$, choose an orthogonal basis of $P_{\ell_2}^q \subseteq R_2^q$ with respect to $Q_{\ell_2}^q$:

$$\{w_1^q, w_2^q, \dots, w_{n(q)}^q\} \subseteq P_{\ell_2}^q.$$

Here we use the upper indices to indicate the degrees of basis elements. To each pair of v_i^p and w_j^q , we associate a graded subspace of R^* :

$$B^*(v_i^p, w_j^q) := B^*(v_i^p) \otimes_{\mathbb{R}} B^*(w_j^q), \quad \text{where}$$

$$B^*(v_i^p) := \langle v_i^p \rangle \oplus \ell_1 \langle v_i^p \rangle \oplus \dots \oplus \ell_1^{r_1-2p} \langle v_i^p \rangle \subseteq R_1^*,$$

$$B^*(w_j^q) := \langle w_j^q \rangle \oplus \ell_2 \langle w_j^q \rangle \oplus \dots \oplus \ell_2^{r_2-2q} \langle w_j^q \rangle \subseteq R_2^*,$$

Let us compare the tensor product $B^*(v_i^p, w_j^q)$ with the truncated polynomial ring

$$S_{p,q}^* := \mathbb{R}[x_1, x_2] / (x_1^{r_1-2p+1}, x_2^{r_2-2q+1}).$$

The properties $\text{HR}(\ell_1)$ and $\text{HR}(\ell_2)$ show that, for every nonnegative integer $k \leq \frac{r_1+r_2-2p-2q}{2}$, there is an isometry

$$\left(B^{k+p+q}(v_i^p, w_j^q), Q_\ell^{k+p+q} \right) \simeq \left(S_{p,q}^k, (-1)^{p+q} Q_{x_1+x_2}^k \right).$$

Therefore, by Lemma 7.8, the graded subspace $B^*(v_i^p, w_j^q) \subseteq R^*$ satisfies $\text{HR}(\ell)$.

The properties $\text{HL}(\ell_1)$ and $\text{HL}(\ell_2)$ imply that there is a direct sum decomposition

$$R^* = \bigoplus_{p,q,i,j} B^*(v_i^p, w_j^q).$$

It is enough to prove that the above decomposition is orthogonal with respect to Q_ℓ^* :

Claim. Any two distinct summands of R^* satisfy $B^*(v, w) \perp_{Q_\ell^*} B^*(v', w')$.

For the proof of the claim, we may suppose that $w \neq w'$. The orthogonality of the Lefschetz decomposition for R_2^* with respect to $Q_{\ell_2}^*$ shows that

$$B(w) \perp_{\text{PD}} B(w').$$

By the fourth assertion of Lemma 7.11, the above implies

$$B^*(v, w) \perp_{\text{PD}} B^*(v', w').$$

By the third assertion of Lemma 7.11, this gives the claimed statement. \square

7.3. Let Σ be a unimodular fan in $N_{\mathbb{R}}$. For our purposes, it will be enough to assume that Σ is simplicial.

Definition 7.12. We say that Σ satisfies the *Poincaré duality of dimension r* if $A^*(\Sigma)_{\mathbb{R}}$ is a Poincaré duality algebra of dimension r .

In the remainder of this subsection, we suppose that Σ satisfies the Poincaré duality of dimension r . We fix an isomorphism, called the *degree map* for Σ ,

$$\text{deg} : A^r(\Sigma)_{\mathbb{R}} \longrightarrow \mathbb{R}.$$

As before, we write V_{Σ} for the set of primitive ray generators of Σ .

Note that for any nonnegative integer q and $\mathbf{e} \in V_{\Sigma}$ there is a commutative diagram

$$\begin{array}{ccc} A^q(\Sigma) & \xrightarrow{p_{\mathbf{e}}} & A^q(\text{star}(\mathbf{e}, \Sigma)) \\ & \searrow^{x_{\mathbf{e}} \cdot -} & \downarrow^{x_{\mathbf{e}} \cdot -} \\ & & A^{q+1}(\Sigma), \end{array}$$

where $p_{\mathbf{e}}$ is the pullback homomorphism $p_{\mathbf{e} \in \Sigma}$ and $x_{\mathbf{e}} \cdot -$ are the multiplications by $x_{\mathbf{e}}$. It follows that there is a surjective graded ring homomorphism

$$\pi_{\mathbf{e}} : A^*(\text{star}(\mathbf{e}, \Sigma)) \longrightarrow A^*(\Sigma)/\text{ann}(x_{\mathbf{e}}).$$

Proposition 7.13. The star of \mathbf{e} in Σ satisfies the Poincaré duality of dimension $r - 1$ if and only if $\pi_{\mathbf{e}}$ is an isomorphism:

$$A^*(\text{star}(\mathbf{e}, \Sigma)) \simeq A^*(\Sigma)/\text{ann}(x_{\mathbf{e}}).$$

Proof. The “if” direction follows from Proposition 7.2: The quotient $A^*(\Sigma)/\text{ann}(x_{\mathbf{e}})$ is a Poincaré duality algebra of dimension $r - 1$.

The “only if” direction follows from the observation that any surjective graded ring homomorphism between Poincaré duality algebras of the same dimension is an isomorphism. \square

Definition 7.14. Let Σ be a fan that satisfies Poincaré duality of dimension r . We say that

- (1) Σ satisfies the *hard Lefschetz property* if $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HL}(\ell)$ for all $\ell \in \mathcal{K}_{\Sigma}$,
- (2) Σ satisfies the *Hodge-Riemann relations* if $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HR}(\ell)$ for all $\ell \in \mathcal{K}_{\Sigma}$, and
- (3) Σ satisfies the *local Hodge-Riemann relations* if the Poincaré duality algebra

$$A^*(\Sigma)_{\mathbb{R}}/\text{ann}(x_{\mathbf{e}})$$

satisfies $\text{HR}(\ell_{\mathbf{e}})$ with respect to the degree map induced by $x_{\mathbf{e}}$ for all $\ell \in \mathcal{K}_{\Sigma}$ and $\mathbf{e} \in V_{\Sigma}$.

Hereafter we write $\ell_{\mathbf{e}}$ for the image of ℓ in the quotient $A^*(\Sigma)_{\mathbb{R}}/\text{ann}(x_{\mathbf{e}})$.

Proposition 7.15. If Σ satisfies the local Hodge-Riemann relations, then Σ satisfies the hard Lefschetz property.

Proof. By definition, for $\ell \in \mathcal{K}_{\Sigma}$ there are positive real numbers $c_{\mathbf{e}}$ such that

$$\ell = \sum_{\mathbf{e} \in V_{\Sigma}} c_{\mathbf{e}} x_{\mathbf{e}} \in A^1(\Sigma)_{\mathbb{R}}.$$

We need to show that the Lefschetz operator L_{ℓ}^q on $A^q(\Sigma)_{\mathbb{R}}$ is injective for all $q \leq \frac{r}{2}$. Nothing is claimed when $r = 2q$, so we may assume that $r - 2q$ is positive.

Let f be an element in the kernel of L_{ℓ}^q , and write $f_{\mathbf{e}}$ for the image of f in the quotient $A^q(\Sigma)_{\mathbb{R}}/\text{ann}(x_{\mathbf{e}})$. Note that the element f has the following properties:

- (1) For all $\mathbf{e} \in V_{\Sigma}$, the image $f_{\mathbf{e}}$ belongs to the primitive subspace $P_{\ell_{\mathbf{e}}}^q$, and
- (2) for the positive real numbers $c_{\mathbf{e}}$ as above, we have

$$\sum_{\mathbf{e} \in V_{\Sigma}} c_{\mathbf{e}} Q_{\ell_{\mathbf{e}}}^q(f_{\mathbf{e}}, f_{\mathbf{e}}) = Q_{\ell}^q(f, f) = 0.$$

By the local Hodge-Riemann relations, the two properties above show that all the $f_{\mathbf{e}}$ are zero:

$$x_{\mathbf{e}} \cdot f = 0 \in A^*(\Sigma)_{\mathbb{R}} \text{ for all } \mathbf{e} \in V_{\Sigma}.$$

Since the elements $x_{\mathbf{e}}$ generate the Poincaré duality algebra $A^*(\Sigma)_{\mathbb{R}}$, this implies that $f = 0$. \square

Proposition 7.16. If Σ satisfies the hard Lefschetz property, then the following are equivalent:

- (1) $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HR}(\ell)$ for some $\ell \in \mathcal{K}_{\Sigma}$.
- (2) $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HR}(\ell)$ for all $\ell \in \mathcal{K}_{\Sigma}$.

Proof. Let ℓ_0 and ℓ_1 be elements of \mathcal{K}_{Σ} , and suppose that $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HR}(\ell_0)$. Consider the parametrized family

$$\ell_t := (1-t)\ell_0 + t\ell_1, \quad 0 \leq t \leq 1.$$

Since \mathcal{K}_{Σ} is convex, the elements ℓ_t are ample for all t .

Note that $Q_{\ell_t}^q$ are nondegenerate on $A^q(\Sigma)_{\mathbb{R}}$ for all t and $q \leq \frac{r}{2}$ because Σ satisfies the hard Lefschetz property. It follows that the signatures of $Q_{\ell_t}^q$ should be independent of t for all $q \leq \frac{r}{2}$.

Since $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HR}(\ell_0)$, the common signature should be

$$\sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} A^p(\Sigma)_{\mathbb{R}} - \dim_{\mathbb{R}} A^{p-1}(\Sigma)_{\mathbb{R}} \right).$$

We conclude by Proposition 7.6 that $A^*(\Sigma)_{\mathbb{R}}$ satisfies $\text{HR}(\ell_1)$. \square

8. PROOF OF THE MAIN THEOREM

8.1. As a final preparation for the proof of the main theorem, we show that the property HR is preserved by a matroidal flip for particular choices of ample classes.

Let M be as before, and consider the matroidal flip from \mathcal{P}_- to \mathcal{P}_+ with center Z . We will use the following homomorphisms:

- (1) The pullback homomorphism $\Phi_Z : A^*(M, \mathcal{P}_-) \rightarrow A^*(M, \mathcal{P}_+)$.
- (2) The Gysin homomorphisms $\Psi_Z^{p,q} : A^{q-p}(M_Z) \rightarrow A^q(M, \mathcal{P}_+)$.
- (3) The pullback homomorphism $p_Z : A^*(M, \mathcal{P}_-) \rightarrow A^*(M_Z)$.

The homomorphism p_Z is the graded ring homomorphism $p_{\sigma_{Z<\emptyset} \in \Sigma_{M, \mathcal{P}_-}}$ obtained from the identification

$$\text{star}(\sigma_{Z<\emptyset}, \Sigma_{M, \mathcal{P}_-}) \simeq \Sigma_{M_Z}.$$

In the remainder of this section, we fix a strictly convex piecewise linear function ℓ_- on $\Sigma_{M, \mathcal{P}_-}$. For nonnegative real numbers t , we set

$$\ell_+(t) := \Phi_Z(\ell_-) - tx_Z \in A^1(M, \mathcal{P}_+) \otimes_{\mathbb{Z}} \mathbb{R}.$$

We write ℓ_Z for the pullback of ℓ_- to the star of the cone $\sigma_{Z<\emptyset}$ in the Bergman fan $\Sigma_{M, \mathcal{P}_-}$:

$$\ell_Z := p_Z(\ell_-) \in A^1(M_Z) \otimes_{\mathbb{Z}} \mathbb{R}.$$

Proposition 4.4 shows that ℓ_Z is the class of a strictly convex piecewise linear function on Σ_{M_Z} .

Lemma 8.1. $\ell_+(t)$ is strictly convex for all sufficiently small positive t .

Proof. It is enough to show that $\ell_+(t)$ is strictly convex around a given cone $\sigma_{I<\mathcal{F}}$ in $\Sigma_{M, \mathcal{P}_+}$.

When $Z \notin \mathcal{F}$, the cone $\sigma_{I<\mathcal{F}}$ is in the fan $\Sigma_{M, \mathcal{P}_-}$, and hence we may suppose that

$$\ell_- \text{ is zero on } \sigma_{I<\mathcal{F}} \text{ and positive on the link of } \sigma_{I<\mathcal{F}} \text{ in } \Sigma_{M, \mathcal{P}_-}.$$

It is straightforward to deduce from the above that, for all sufficiently small positive t ,

$$\ell_+(t) \text{ is zero on } \sigma_{I<\mathcal{F}} \text{ and positive on the link of } \sigma_{I<\mathcal{F}} \text{ in } \Sigma_{M, \mathcal{P}_+}.$$

More precisely, the statement is valid for all t that satisfies the inequalities

$$0 < t < \sum_{i \in Z \setminus I} \ell_-(\mathbf{e}_i).$$

Note that $Z \setminus I$ is nonempty and each of the summands in the right-hand side is positive.

When $Z \in \mathcal{F}$, the cone $\sigma_{Z < \mathcal{F} \setminus \{Z\}}$ is in the fan $\Sigma_{M, \mathcal{P}_-}$, and hence we may suppose that

ℓ_- is zero on $\sigma_{Z < \mathcal{F} \setminus \{Z\}}$ and positive on the link of $\sigma_{Z < \mathcal{F} \setminus \{Z\}}$ in $\Sigma_{M, \mathcal{P}_-}$.

Let J be the flat min $\mathcal{F} \setminus \{Z\}$, and let $m(t)$ be the linear function on N_E defined by setting

$$\mathbf{e}_i \mapsto \begin{cases} \frac{t}{|Z \setminus I|} & \text{if } i \in Z \setminus I, \\ \frac{-t}{|J \setminus Z|} & \text{if } i \in J \setminus Z, \\ 0 & \text{if otherwise.} \end{cases}$$

It is straightforward to deduce from the above that, for all sufficiently small positive t ,

$\ell_+(t) + m(t)$ is zero on $\sigma_{I < \mathcal{F}}$ and positive on the link of $\sigma_{I < \mathcal{F}}$ in $\Sigma_{M, \mathcal{P}_+}$.

More precisely, the latter statement is valid for all t that satisfies the inequalities

$$0 < t < \min \left\{ \ell_-(\mathbf{e}_F), \mathbf{e}_F \text{ is in the link of } \sigma_{Z < \mathcal{F} \setminus \{Z\}} \text{ in } \Sigma_{M, \mathcal{P}_-} \right\}.$$

Here the minimum of the empty set is defined to be ∞ . □

We write “deg” for the degree map of M and of M_Z , and fix the degree maps

$$\begin{aligned} \deg_+ : A^r(M, \mathcal{P}_+) &\longrightarrow \mathbb{Z}, & a &\longmapsto \deg(\Phi_{\mathcal{P}_+^c}(a)), \\ \deg_- : A^r(M, \mathcal{P}_-) &\longrightarrow \mathbb{Z}, & a &\longmapsto \deg(\Phi_{\mathcal{P}_-^c}(a)), \end{aligned}$$

see Definition 6.9. We omit the subscripts $+$ and $-$ from the notation when there is no danger of confusion. The goal of this subsection is to prove the following.

Proposition 8.2. Let ℓ_- , ℓ_Z , and $\ell_+(t)$ be as above, and suppose that

- (1) the Chow ring of $\Sigma_{M, \mathcal{P}_-}$ satisfies $\text{HR}(\ell_-)$, and
- (2) the Chow ring of Σ_{M_Z} satisfies $\text{HR}(\ell_Z)$.

Then the Chow ring of $\Sigma_{M, \mathcal{P}_+}$ satisfies $\text{HR}(\ell_+(t))$ for all sufficiently small positive t .

Hereafter we suppose $\text{HR}(\ell_-)$ and $\text{HR}(\ell_Z)$. We introduce the main characters appearing in the proof of Proposition 8.2:

- (1) A Poincaré duality algebra of dimension r :

$$A_+^* := \bigoplus_{q=0}^r A_+^q, \quad A_+^q := A^q(M, \mathcal{P}_+) \otimes_{\mathbb{Z}} \mathbb{R}.$$

- (2) A Poincaré duality algebra of dimension r :

$$A_-^* := \bigoplus_{q=0}^r A_-^q, \quad A_-^q := (\text{im } \Phi_Z^q) \otimes_{\mathbb{Z}} \mathbb{R}.$$

- (3) A Poincaré duality algebra of dimension $r - 2$:

$$T_Z^* := \bigoplus_{q=0}^{r-2} T_Z^q, \quad T_Z^q := \left(\mathbb{Z}[x_Z] / (x_Z^{\text{rk}(Z)-1}) \otimes_{\mathbb{Z}} A^*(M_Z) \right)^q \otimes_{\mathbb{Z}} \mathbb{R}.$$

(4) A graded subspace of A_+^* , the sum of the images of the Gysin homomorphisms:

$$G_Z^* := \bigoplus_{q=1}^{r-1} G_Z^q, \quad G_Z^q := \bigoplus_{p=1}^{\text{rk}(Z)-1} (\text{im } \Psi_Z^{p,q}) \otimes_{\mathbb{Z}} \mathbb{R}.$$

The truncated polynomial ring in the definition of T_Z^* is given the degree map

$$(-x_Z)^{\text{rk}(Z)-2} \mapsto 1,$$

so that the truncated polynomial ring satisfies $\text{HR}(-x_Z)$. The tensor product T_Z^* is given the induced degree map

$$(-x_Z)^{\text{rk}(Z)-2} x_{\mathcal{F}} \mapsto 1.$$

It follows from Proposition 7.7 that the tensor product satisfies $\text{HR}(1 \otimes \ell_Z - x_Z \otimes 1)$.

Definition 8.3. For nonnegative $q \leq \frac{r}{2}$, we write the Poincaré duality pairings for A_-^* and T_Z^* by

$$\begin{aligned} \langle -, - \rangle_{A_-^*}^q &: A_-^q \times A_-^{r-q} \longrightarrow \mathbb{R}, \\ \langle -, - \rangle_{T_Z^*}^{q-1} &: T_Z^{q-1} \times T_Z^{r-q-1} \longrightarrow \mathbb{R}. \end{aligned}$$

We omit the superscripts q and $q-1$ from the notation when there is no danger of confusion.

Theorem 6.18 shows that Φ_Z defines an isomorphism between the graded rings

$$A^*(M, \mathcal{P}_-) \otimes_{\mathbb{Z}} \mathbb{R} \simeq A_-^*,$$

and that there is a decomposition into a direct sum

$$A_+^* = A_-^* \oplus G_Z^*.$$

In addition, it shows that $x_Z \cdot -$ is an isomorphism between the graded vector spaces

$$T_Z^* \simeq G_Z^{*+1}.$$

The inverse of the isomorphism $x_Z \cdot -$ will be denoted $x_Z^{-1} \cdot -$.

We equip the above graded vector spaces with the following symmetric bilinear forms.

Definition 8.4. Let q be a nonnegative integer $\leq \frac{r}{2}$.

- (1) $(A_+^q, Q_-^q \oplus Q_Z^q)$: Q_-^q and Q_Z^q are the bilinear forms on A_-^q and G_Z^q defined below.
- (2) (A_-^q, Q_-^q) : Q_-^q is the restriction of the Hodge-Riemann form $Q_{\ell_+(0)}^q$ to A_-^q .
- (3) $(T_Z^q, Q_{\mathcal{F}}^q)$: $Q_{\mathcal{F}}^q$ is the Hodge-Riemann form associated to $\mathcal{F} := (1 \otimes \ell_Z - x_Z \otimes 1) \in T_Z^1$.
- (4) (G_Z^q, Q_Z^q) : Q_Z^q is the bilinear form defined by saying that $x_Z \cdot -$ gives an isometry

$$(T_Z^{q-1}, Q_{\mathcal{F}}^{q-1}) \simeq (G_Z^q, Q_Z^q).$$

We observe that $Q_-^q \oplus Q_Z^q$ satisfies the following version of Hodge-Riemann relations:

Proposition 8.5. The bilinear form $Q_-^q \oplus Q_Z^q$ is nondegenerate on A_+^q and has signature

$$\sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} A_+^p - \dim_{\mathbb{R}} A_+^{p-1} \right) \text{ for all nonnegative } q \leq \frac{r}{2}.$$

Proof. Theorem 6.18 shows that $\Phi_Z \otimes_{\mathbb{Z}} \mathbb{R}$ defines an isometry

$$\left(A^q(\mathbb{M}, \mathcal{P}_-)_{\mathbb{R}}, Q_{\ell_-}^q \right) \simeq \left(A_-^q, Q_-^q \right).$$

It follows from the assumption on $\Sigma_{\mathbb{M}, \mathcal{P}_-}$ that Q_-^q is nondegenerate on A_-^q and has signature

$$\sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} A_-^p - \dim_{\mathbb{R}} A_-^{p-1} \right).$$

It follows from the assumption on $\Sigma_{\mathbb{M}_Z}$ that Q_Z^q is nondegenerate on G_Z^q and has signature

$$\begin{aligned} \sum_{p=0}^{q-1} (-1)^{q-p-1} \left(\dim_{\mathbb{R}} T_Z^p - \dim_{\mathbb{R}} T_Z^{p-1} \right) &= \sum_{p=0}^{q-1} (-1)^{q-p-1} \left(\dim_{\mathbb{R}} G_Z^{p+1} - \dim_{\mathbb{R}} G_Z^p \right) \\ &= \sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} G_Z^p - \dim_{\mathbb{R}} G_Z^{p-1} \right). \end{aligned}$$

The assertion is deduced from the fact that the signature of the sum is the sum of the signatures. \square

We now construct a continuous family of symmetric bilinear forms Q_t^q on A_+^q parametrized by positive real numbers t . This family Q_t^q will be shown to have the following properties:

(1) For every positive real number t , there is an isometry

$$\left(A_+^q, Q_t^q \right) \simeq \left(A_+^q, Q_{\ell_+(t)}^q \right).$$

(2) The sequence Q_t^q as t goes to zero converges to the sum of Q_-^q and Q_Z^q :

$$\lim_{t \rightarrow 0} Q_t^q = Q_-^q \oplus Q_Z^q.$$

For positive real numbers t , we define a graded linear transformation

$$S_t : A_+^* \longrightarrow A_+^*$$

to be the sum of the identity on A_-^* and the linear transformations

$$\left(\text{im } \Psi_Z^{p,q} \right) \otimes_{\mathbb{Z}} \mathbb{R} \longrightarrow \left(\text{im } \Psi_Z^{p,q} \right) \otimes_{\mathbb{Z}} \mathbb{R}, \quad a \longmapsto t^{-\frac{\text{rk}(Z)}{2} + p} a.$$

The inverse transformation S_t^{-1} is the sum of the identity on A_-^* and the linear transformations

$$\left(\text{im } \Psi_Z^{p,q} \right) \otimes_{\mathbb{Z}} \mathbb{R} \longrightarrow \left(\text{im } \Psi_Z^{p,q} \right) \otimes_{\mathbb{Z}} \mathbb{R}, \quad a \longmapsto t^{\frac{\text{rk}(Z)}{2} - p} a.$$

Definition 8.6. The symmetric bilinear form Q_t^q is defined so that S_t defines an isometry

$$\left(A_+^q, Q_t^q\right) \simeq \left(A_+^q, Q_{\ell_+(t)}^q\right) \text{ for all nonnegative integers } q \leq \frac{r}{2}.$$

In other words, for any elements $a_1, a_2 \in A_+^q$, we set

$$Q_t^q(a_1, a_2) := (-1)^q \deg(S_t(a_1) \cdot \ell_+(t)^{r-2q} \cdot S_t(a_2)).$$

The first property of Q_t^q mentioned above is built into the definition. We verify the assertion on the limit of Q_t^q as t goes to zero.

Proposition 8.7. For all nonnegative integers $q \leq \frac{r}{2}$, we have

$$\lim_{t \rightarrow 0} Q_t^q = Q_-^q \oplus Q_Z^q.$$

Proof. We first construct a deformation of the Poincaré duality pairing $A_+^q \times A_+^{r-q} \rightarrow \mathbb{R}$:

$$\langle a_1, a_2 \rangle_t^q := \deg(S_t(a_1), S_t(a_2)), \quad t > 0.$$

We omit the upper index q when there is no danger of confusion.

Claim (1). For any $b_1, b_2 \in A_-^*$ and $c_1, c_2 \in G_Z^*$ and $a_1 = b_1 + c_1, a_2 = b_2 + c_2 \in A_+^*$,

$$\langle a_1, a_2 \rangle_0 := \lim_{t \rightarrow 0} \langle a_1, a_2 \rangle_t = \langle b_1, b_2 \rangle_{A_-^*} - \langle x_Z^{-1}c_1, x_Z^{-1}c_2 \rangle_{T_Z^*}.$$

We write $z := \text{rk}(Z)$ and choose bases of A_+^q and A_+^{r-q} that respect the decompositions

$$A_+^q = A_-^q \oplus \left(\text{im } \Psi_Z^{1,q} \oplus \text{im } \Psi_Z^{2,q} \oplus \cdots \oplus \text{im } \Psi_Z^{z-1,q} \right) \otimes_{\mathbb{Z}} \mathbb{R}, \text{ and}$$

$$A_+^{r-q} = A_-^{r-q} \oplus \left(\text{im } \Psi_Z^{z-1,r-q} \oplus \text{im } \Psi_Z^{z-2,r-q} \oplus \cdots \oplus \text{im } \Psi_Z^{1,r-q} \right) \otimes_{\mathbb{Z}} \mathbb{R}.$$

Let \mathcal{M}_- be the matrix of the Poincaré duality pairing between A_-^q and A_-^{r-q} , and let \mathcal{M}_{p_1, p_2} is the matrix of the Poincaré duality pairing between $\text{im } \Psi_Z^{p_1, q} \otimes_{\mathbb{Z}} \mathbb{R}$ and $\text{im } \Psi_Z^{p_2, r-q} \otimes_{\mathbb{Z}} \mathbb{R}$. Lemma 6.20 shows that the matrix of the deformed Poincaré pairing on A_+^* is

$$\begin{bmatrix} \mathcal{M}_- & 0 & 0 & 0 & \cdots & 0 \\ 0 & \mathcal{M}_{1, z-1} & t\mathcal{M}_{2, z-1} & t^2\mathcal{M}_{3, z-1} & \cdots & t^{z-2}\mathcal{M}_{z-1, z-1} \\ 0 & 0 & \mathcal{M}_{2, z-2} & t\mathcal{M}_{3, z-2} & \cdots & t^{z-3}\mathcal{M}_{z-1, z-2} \\ 0 & 0 & 0 & \mathcal{M}_{3, z-3} & \cdots & t^{z-4}\mathcal{M}_{z-1, z-3} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \mathcal{M}_{z-1, 1} \end{bmatrix}.$$

The claim on the limit of the deformed Poincaré duality pairing follows. The minus sign on the right-hand side of the claim comes from the following computation made in Proposition 6.17:

$$\deg(x_Z^{\text{rk}(Z)} x_{\mathcal{X}}) = (-1)^{\text{rk}(Z)-1}.$$

We use the deformed Poincaré duality pairing to understand the limit of the bilinear form Q_t^q . For an element a of A_+^1 , we write the multiplication with a by

$$M^a : A_+^* \longrightarrow A_+^{*+1}, \quad x \longmapsto a \cdot x,$$

and define its deformation $M_t^a := S_t^{-1} \circ M^a \circ S_t$. In terms of the operator $M_t^{\ell_+(t)}$, the bilinear form Q_t^q can be written

$$\begin{aligned} Q_t^q(a_1, a_2) &= (-1)^q \deg \left(S_t(a_1) \cdot M^{\ell_+(t)} \circ \dots \circ M^{\ell_+(t)} \circ S_t(a_2) \right) \\ &= (-1)^q \deg \left(S_t(a_1) \cdot S_t \circ M_t^{\ell_+(t)} \circ \dots \circ M_t^{\ell_+(t)}(a_2) \right) \\ &= (-1)^q \left\langle a_1, M_t^{\ell_+(t)} \circ \dots \circ M_t^{\ell_+(t)}(a_2) \right\rangle_t \end{aligned}$$

Define linear operators $M^{1 \otimes \ell_Z}$, $M^{x_Z \otimes 1}$, and $M^{\mathcal{F}}$ on G_Z^* by the isomorphisms

$$\begin{aligned} (G_Z^*, M^{1 \otimes \ell_Z}) &\simeq (T^{*-1}, 1 \otimes \ell_Z \cdot -), \\ (G_Z^*, M^{x_Z \otimes 1}) &\simeq (T^{*-1}, x_Z \otimes 1 \cdot -), \\ (G_Z^*, M^{\mathcal{F}}) &\simeq (T^{*-1}, \mathcal{F} \cdot -). \end{aligned}$$

Note that the linear operator $M^{\mathcal{F}}$ is the difference $M^{1 \otimes \ell_Z} - M^{x_Z \otimes 1}$.

Claim (2). The limit of the operator $M_t^{\ell_+(t)}$ as t goes to zero decomposes into the sum

$$\left(A_+^*, \lim_{t \rightarrow 0} M_t^{\ell_+(t)} \right) = \left(A_-^* \oplus G_Z^*, M^{\ell_+(0)} \oplus M^{\mathcal{F}} \right).$$

Assuming the second claim, we finish the proof as follows: We have

$$\lim_{t \rightarrow 0} Q_t^q(a_1, a_2) = (-1)^q \lim_{t \rightarrow 0} \left\langle a_1, M_t^{\ell_+(t)} \circ \dots \circ M_t^{\ell_+(t)}(a_2) \right\rangle_t$$

and from the first and the second claim, we see that the right-hand side is

$$(-1)^q \left\langle a_1, (M^{\ell_+(0)} \oplus M^{\mathcal{F}}) \circ \dots \circ (M^{\ell_+(0)} \oplus M^{\mathcal{F}})(a_2) \right\rangle_0 = Q_-^q(b_1, b_2) + Q_Z^q(c_1, c_2),$$

where $a_i = b_i + c_i$ for $b_i \in A_-^*$ and $c_i \in G_Z^*$. Notice that the minus sign in the first claim cancels with $(-1)^{q-1}$ in the Hodge-Riemann form

$$\left(T_Z^{q-1}, Q_{\mathcal{F}}^{q-1} \right) \simeq \left(G_Z^q, Q_Z^q \right).$$

We now prove the second claim made above. Write $M_t^{\ell_+(t)}$ as the difference

$$M_t^{\ell_+(t)} = S_t^{-1} \circ M^{\ell_+(t)} \circ S_t = S_t^{-1} \circ \left(M^{\ell_+(0)} - M^{tx_Z} \right) \circ S_t = M_t^{\ell_+(0)} - M_t^{tx_Z}.$$

By Lemma 6.20, the operators $M^{\ell_+(0)}$ and S_t commute, and hence

$$\left(A_+^*, M_t^{\ell_+(0)} \right) = \left(A_+^*, M^{\ell_+(0)} \right) = \left(A_-^* \oplus G_Z^*, M^{\ell_+(0)} \oplus M^{1 \otimes \ell_Z} \right).$$

Lemma 6.20 shows that the matrix of M^{xz} in the chosen bases of A_+^q and A_+^{q+1} is of the form

$$\begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & B_0 \\ A & 0 & 0 & \cdots & 0 & B_1 \\ 0 & \text{Id} & 0 & \cdots & 0 & B_2 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \text{Id} & 0 & B_{z-2} \\ 0 & 0 & \cdots & 0 & \text{Id} & B_{z-1} \end{bmatrix},$$

where “Id” are the identity matrices representing

$$A^{q-p}(\mathbb{M}_Z)_{\mathbb{R}} \simeq \text{im } \Psi_Z^{p,q} \longrightarrow \text{im } \Psi_Z^{p+1,q+1} \simeq A^{q-p}(\mathbb{M}_Z)_{\mathbb{R}}.$$

Note that the matrix of the deformed operator M_t^{xz} can be written

$$\begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & t^{\frac{\text{rk}(Z)}{2}} B_0 \\ t^{\frac{\text{rk}(Z)}{2}} A & 0 & 0 & \cdots & 0 & t^{\text{rk}(Z)-1} B_1 \\ 0 & \text{Id} & 0 & \cdots & 0 & t^{\text{rk}(Z)-1} B_2 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \text{Id} & 0 & t^2 B_{z-2} \\ 0 & 0 & \cdots & 0 & \text{Id} & t B_{z-1} \end{bmatrix}.$$

At the limit $t = 0$, the matrix represents the sum $0 \oplus M^{xz \otimes 1}$, and therefore

$$\begin{aligned} \left(A_+^*, \lim_{t \rightarrow 0} M_t^{\ell_+(t)} \right) &= \left(A_-^* \oplus G_Z^*, M^{\ell_+(0)} \oplus M^{1 \otimes \ell_Z} \right) - \left(A_-^* \oplus G_Z^*, 0 \oplus M^{xz \otimes 1} \right) \\ &= \left(A_-^* \oplus G_Z^*, M^{\ell_+(0)} \oplus M^{\mathcal{S}} \right). \end{aligned}$$

This completes the proof of the second claim. \square

Proof of Proposition 8.2. By Proposition 8.5 and Proposition 8.7, we know that $\lim_{t \rightarrow 0} Q_t^q$ is non-degenerate on A_+^q and has signature

$$\sum_{p=0}^q (-1)^{q-p} \left(\dim_{\mathbb{R}} A_+^p - \dim_{\mathbb{R}} A_+^{p-1} \right) \text{ for all nonnegative } q \leq \frac{r}{2}.$$

Therefore the same must be true for Q_t^q for all sufficiently small positive t . By construction, there is an isometry

$$\left(A_+^q, Q_t^q \right) \simeq \left(A_+^q, Q_{\ell_+(t)}^q \right),$$

and thus A_+^* satisfies $\text{HR}(\ell_+(t))$ for all sufficiently small positive t . \square

8.2. We are now ready to prove the main theorem. We write “deg” for the degree map of \mathbb{M} and, for an order filter \mathcal{P} of $\mathcal{P}_{\mathbb{M}}$, fix an isomorphism

$$A^r(\mathbb{M}, \mathcal{P}) \longrightarrow \mathbb{Z}, \quad a \longmapsto \text{deg}(\Phi_{\mathcal{P}^c}(a)).$$

Theorem 8.8 (Main Theorem). Let \mathbb{M} be a loopless matroid, and let \mathcal{P} be an order filter of $\mathcal{P}_{\mathbb{M}}$.

- (1) The Bergman fan $\Sigma_{M, \mathcal{P}}$ satisfies the hard Lefschetz property.
- (2) The Bergman fan $\Sigma_{M, \mathcal{P}}$ satisfies the Hodge-Riemann relations.

When $\mathcal{P} = \mathcal{P}_M$, the above implies Theorem 1.4 in the introduction because any strictly submodular function defines a strictly convex piecewise linear function on Σ_M .

We prove Theorem 8.8 by lexicographic induction on the rank of M and the cardinality of \mathcal{P} . Set $\mathcal{P} = \mathcal{P}_+$, and consider the matroidal flip from \mathcal{P}_- to \mathcal{P}_+ with center Z .

Proof. By Proposition 4.7 and Proposition 4.8, we may replace M by the associated combinatorial geometry \overline{M} . Thus we may assume that M has no rank 1 flat of cardinality greater than 1. In this case, Proposition 3.5 shows that the star of every ray in $\Sigma_{M, \mathcal{P}}$ is a product of at most two smaller Bergman fans to which the induction hypothesis applies.

By Proposition 7.7 and the induction hypothesis applied to the stars, $\Sigma_{M, \mathcal{P}}$ satisfies the local Hodge-Riemann relations. By Proposition 7.15, this implies that $\Sigma_{M, \mathcal{P}}$ satisfies the hard Lefschetz property.

Lastly, we show that $\Sigma_{M, \mathcal{P}}$ satisfies the Hodge-Riemann relations. Since $\Sigma_{M, \mathcal{P}}$ satisfies the hard Lefschetz property, Proposition 7.16 shows that it is enough to prove $\text{HR}(\ell)$ for some $\ell \in \mathcal{K}_{M, \mathcal{P}}$. This follows from Proposition 8.2 and the induction hypothesis applied to $\Sigma_{M, \mathcal{P}_-}$ and Σ_{M_Z} . \square

We remark that the same inductive approach can be used to prove the following stronger statement (see [Cat08] for an overview of the analogous facts in the context of convex polytopes and compact Kähler manifolds). We leave details to the interested reader.

Theorem 8.9. Let M be a loopless matroid on E , and let \mathcal{P} be an order filter of \mathcal{P}_M .

- (1) The Bergman fan $\Sigma_{M, \mathcal{P}}$ satisfies the *mixed hard Lefschetz theorem*: For any multiset

$$\mathcal{L} := \{\ell_1, \ell_2, \dots, \ell_{r-2q}\} \subseteq \mathcal{K}_{M, \mathcal{P}},$$

the linear map given by the multiplication

$$L_{\mathcal{L}}^q : A^q(M, \mathcal{P})_{\mathbb{R}} \longrightarrow A^{r-q}(M, \mathcal{P})_{\mathbb{R}}, \quad a \longmapsto (\ell_1 \ell_2 \cdots \ell_{r-2q}) \cdot a$$

is an isomorphism for all nonnegative integers $q \leq \frac{r}{2}$.

- (2) The Bergman fan $\Sigma_{M, \mathcal{P}}$ satisfies the *mixed Hodge-Riemann Relations*: For any multiset

$$\mathcal{L} := \{\ell_1, \ell_2, \dots, \ell_{r-2q}\} \subseteq \mathcal{K}_{M, \mathcal{P}} \text{ and any } \ell \in \mathcal{K}_{M, \mathcal{P}},$$

the symmetric bilinear form given by the multiplication

$$Q_{\mathcal{L}}^q : A^q(M, \mathcal{P})_{\mathbb{R}} \times A^q(M, \mathcal{P})_{\mathbb{R}} \longrightarrow \mathbb{R}, \quad (a_1, a_2) \longmapsto (-1)^q \deg(a_1 \cdot L_{\mathcal{L}}^q(a_2))$$

is positive definite on the kernel of $\ell \cdot L_{\mathcal{L}}^q$ for all nonnegative integers $q \leq \frac{r}{2}$.

9. LOG-CONCAVITY CONJECTURES

9.1. Let M be a loopless matroid of rank $r + 1$ on the ground set $E = \{0, 1, \dots, n\}$. The *characteristic polynomial* of M is defined to be

$$\chi_M(\lambda) = \sum_{I \subseteq E} (-1)^{|I|} \lambda^{\text{crk}(I)},$$

where the sum is over all subsets $I \subseteq E$ and $\text{crk}(I)$ is the corank of I in M . Equivalently,

$$\chi_M(\lambda) = \sum_{F \subseteq E} \mu_M(\emptyset, F) \lambda^{\text{crk}(F)},$$

where the sum is over all flats $F \subseteq E$ and μ_M is the Möbius function of the lattice of flats of M . Any one of the two descriptions clearly shows that

- (1) the degree of the characteristic polynomial is $r + 1$,
- (2) the leading coefficient of the characteristic polynomial is 1, and
- (3) the characteristic polynomial satisfies $\chi_M(1) = 0$.

See [Zas87, Aig87] for basic properties of the characteristic polynomial and its coefficients.

Definition 9.1. The *reduced characteristic polynomial* $\bar{\chi}_M(\lambda)$ is

$$\bar{\chi}_M(\lambda) := \chi_M(\lambda) / (\lambda - 1).$$

We define a sequence of integers $\mu^0(M), \mu^1(M), \dots, \mu^r(M)$ by the equality

$$\bar{\chi}_M(\lambda) = \sum_{k=0}^r (-1)^k \mu^k(M) \lambda^{r-k}.$$

The first number in the sequence is 1, and the last number in the sequence is the absolute value of the Möbius number $\mu_M(\emptyset, E)$. In general, $\mu^k(M)$ is the alternating sum of the absolute values of the coefficients of the characteristic polynomial

$$\mu^k(M) = w_k(M) - w_{k-1}(M) + \dots + (-1)^k w_0(M).$$

We will show that the Hodge-Riemann relations for $A^*(M)_{\mathbb{R}}$ imply the log-concavity

$$\mu^{k-1}(M) \mu^{k+1}(M) \leq \mu^k(M)^2 \quad \text{for } 0 < k < r.$$

Because the convolution of two log-concave sequences is log-concave, the above implies the log-concavity of the sequence $w_k(M)$.

Definition 9.2. Let $\mathcal{F} = \{F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_k\}$ be a k -step flag of nonempty proper flats of M .

- (1) The flag \mathcal{F} is said to be *initial* if $r(F_m) = m$ for all indices m .
- (2) The flag \mathcal{F} is said to be *descending* if $\min(F_1) > \min(F_2) > \dots > \min(F_k) > 0$.

We write $D_k(M)$ for the set of initial descending k -step flags of nonempty proper flats of M .

For inductive purposes it will be useful to consider the *truncation* of M , denoted $\text{tr}(M)$. This is the matroid on E whose rank function is defined by

$$\text{rk}_{\text{tr}(M)}(I) := \min(\text{rk}_M(I), r).$$

The lattice of flats of $\text{tr}(M)$ is obtained from the lattice of flats of M by removing all the flats of rank r . It follows that, for any nonnegative integer $k < r$, there is a bijection

$$D_k(M) \simeq D_k(\text{tr}(M)),$$

and an equality between the coefficients of the reduced characteristic polynomials

$$\mu^k(M) = \mu^k(\text{tr}(M)).$$

The second equality shows that all the integers $\mu^k(M)$ are positive, see [Zas87, Theorem 7.1.8].

Lemma 9.3. For every positive integer $k \leq r$, we have

$$\mu^k(M) = |D_k(M)|.$$

Proof. The assertion for $k = r$ is the known fact that $\mu^r(M)$ is the number of facets of Δ_M that are glued along their entire boundaries in its lexicographic shelling; see [Bjo92, Proposition 7.6.4]. The general case is obtained from the same equality applied to repeated truncations of M . See [HK12, Proposition 2.4] for an alternative approach using Weisner's theorem. \square

We now show that $\mu^k(M)$ is the degree of the product $\alpha_M^{r-k} \beta_M^k$. See Definition 5.7 for the elements $\alpha_M, \beta_M \in A^1(M)$, and Definition 5.9 for the degree map of M .

Lemma 9.4. For every positive integer $k \leq r$, we have

$$\beta_M^k = \sum_{\mathcal{F}} x_{\mathcal{F}} \in A^*(M),$$

where the sum is over all descending k -step flags of nonempty proper flats of M .

Proof. We prove by induction on the positive integer k . When $k = 1$, the assertion is precisely that $\beta_{M,0}$ represents β_M in the Chow ring of M :

$$\beta_M = \beta_{M,0} = \sum_{0 \notin F} x_F \in A^*(M).$$

In the general case, we use the induction hypothesis for k to write

$$\beta_M^{k+1} = \sum_{\mathcal{F}} \beta_M x_{\mathcal{F}},$$

where the sum is over all descending k -step flags of nonempty proper flats of M . For each of the summands $\beta_M x_{\mathcal{F}}$, we write

$$\mathcal{F} = \{F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_k\}, \text{ and set } i_{\mathcal{F}} := \min(F_1).$$

By considering the representative of β_M corresponding to the element $i_{\mathcal{F}}$, we see that

$$\beta_M x_{\mathcal{F}} = \left(\sum_{i_{\mathcal{F}} \notin F} x_F \right) x_{\mathcal{F}} = \sum_{\mathcal{G}} x_{\mathcal{G}},$$

where the second sum is over all descending flags of nonempty proper flats of M of the form

$$\mathcal{G} = \{F \subsetneq F_1 \subsetneq \cdots \subsetneq F_k\}.$$

This complete the induction. \square

Combining Lemma 9.3, Lemma 9.4, and Proposition 5.8, we see that the coefficients of the reduced characteristic polynomial of M are given by the degrees of the products $\alpha_M^{r-k} \beta_M^k$:

Proposition 9.5. For every nonnegative integer $k \leq r$, we have

$$\mu^k(M) = \deg(\alpha_M^{r-k} \beta_M^k).$$

9.2. Now we explain why the Hodge-Riemann relations imply the log-concavity of the reduced characteristic polynomial. We first state a lemma involving inequalities among degrees of products:

Lemma 9.6. Let ℓ_1 and ℓ_2 be elements of $A^1(M)_{\mathbb{R}}$. If ℓ_2 is nef, then

$$\deg(\ell_1 \ell_1 \ell_2^{r-2}) \deg(\ell_2 \ell_2 \ell_2^{r-2}) \leq \deg(\ell_1 \ell_2 \ell_2^{r-2})^2.$$

Proof. We first consider the case when ℓ_2 is ample. Let $Q_{\ell_2}^1$ be the Hodge-Riemann form

$$Q_{\ell_2}^1 : A^1(M)_{\mathbb{R}} \times A^1(M)_{\mathbb{R}} \longrightarrow \mathbb{R}, \quad (a_1, a_2) \longmapsto -\deg(a_1 \ell_2^{r-2} a_2).$$

Since the Chow ring $A^*(M)$ satisfies $\text{HL}(\ell_2)$, we have the Lefschetz decomposition

$$A^1(M)_{\mathbb{R}} = \langle \ell_2 \rangle \oplus P_{\ell_2}^1(M).$$

Note that the decomposition is orthogonal with respect to the Hodge-Riemann form $Q_{\ell_2}^1$. The property $\text{HR}(\ell_2)$ shows that $Q_{\ell_2}^1$ is negative definite on $\langle \ell_2 \rangle$ and positive definite on its orthogonal complement $P_{\ell_2}^1(M)$.

We consider $Q_{\ell_2}^1$ restricted to the subspace $\langle \ell_1, \ell_2 \rangle \subseteq A^1(M)_{\mathbb{R}}$. If ℓ_1 is not a multiple of ℓ_2 , then the restriction of $Q_{\ell_2}^1$ is neither positive definite nor negative definite, and hence

$$\deg(\ell_1 \ell_1 \ell_2^{r-2}) \deg(\ell_2 \ell_2 \ell_2^{r-2}) < \deg(\ell_1 \ell_2 \ell_2^{r-2})^2.$$

Next consider the case when ℓ_2 is nef. By Proposition 3.3, there is an element ℓ in the ample cone of M . Since ℓ_2 is nef, we have

$$\ell_2(t) := \ell_2 + t\ell \in \mathcal{X}_M \text{ for all positive real numbers } t.$$

Therefore for all positive real numbers t we have

$$\deg(\ell_1 \ell_1 \ell_2(t)^{r-2}) \deg(\ell_2(t) \ell_2(t) \ell_2(t)^{r-2}) \leq \deg(\ell_1 \ell_2(t) \ell_2(t)^{r-2})^2.$$

By taking the limit $t \rightarrow 0$, we obtain the desired inequality. \square

Lemma 9.7. Let M be a loopless matroid.

- (1) The element α_M is the class of a convex piecewise linear function on Σ_M .
- (2) The element β_M is the class of a convex piecewise linear function on Σ_M .

In other words, α_M and β_M are nef.

Proof. For the first assertion, it is enough to show that α_M is the class of a nonnegative piecewise linear function that is zero on a given cone $\sigma_{\emptyset < \mathcal{F}}$ in Σ_M . For this we choose an element i not in any of the flats in \mathcal{F} . The representative $\alpha_{M,i}$ of α_M has the desired property.

Similarly, for the second assertion, it is enough to show that β_M is the class of a nonnegative piecewise linear function that is zero on a given cone $\sigma_{\emptyset < \mathcal{F}}$ in Σ_M . For this we choose an element i in the flat $\min \mathcal{F}$. The representative $\beta_{M,i}$ of β_M has the desired property. \square

Proposition 9.8. For every positive integer $k < r$, we have

$$\mu^{k-1}(M)\mu^{k+1}(M) \leq \mu^k(M)^2.$$

Proof. We prove by induction on the rank of M . When k is less than $r - 1$, the induction hypothesis applies to the truncation of M . When k is $r - 1$, Proposition 9.5 shows that the assertion is equivalent to the inequality

$$\deg(\alpha_M^2 \beta_M^{r-2}) \deg(\beta_M^2 \beta_M^{r-2}) \leq \deg(\alpha_M^1 \beta_M^{r-1})^2.$$

This follows from Lemma 9.6 applied to the nef classes α_M and β_M . \square

As an implication of Proposition 9.8, we conclude with the proof of the announced log-concavity results.

Theorem 9.9. Let M be a matroid, and let G be a graph.

- (1) The coefficients of the reduced characteristic polynomial of M form a log-concave sequence.
- (2) The coefficients of the characteristic polynomial of M form a log-concave sequence.
- (3) The number of independent subsets of size i of M form a log-concave sequence in i .
- (4) The coefficients of the chromatic polynomial of G form a log-concave sequence.

The second item proves the aforementioned conjecture of Heron [Her72], Rota [Rot71], and Welsh [Wel76]. The third item proves the conjecture of Mason [Mas72] and Welsh [Wel71]. The last item proves the conjecture of Read [Rea68] and Hoggar [Hog74].

Proof. It follows from Proposition 9.8 that the coefficients of the reduced characteristic polynomial of M form a log-concave sequence. Since the convolution of two log-concave sequences is a log-concave sequence, the coefficients of the characteristic polynomial of M also form a log-concave sequence.

To justify the third assertion, we consider the free dual extension of M . It is defined by taking the dual of M , placing a new element p in general position (taking the free extension), and again taking the dual. In symbols,

$$M \times p := (M^* + p)^*.$$

The free dual extension $M \times p$ has the following property: The number of independent subsets of size k of M is the absolute value of the coefficient of λ^{r-k} of the reduced characteristic polynomial of M . We refer to [Len12] and also to [Bry77, Bry86] for these facts. It follows that the number of independent subsets of size k of M form a log-concave sequence in k .

For the last assertion, we recall that the chromatic polynomial of a graph is given by the characteristic polynomial of the associated graphic matroid [Wel76]. More precisely, we have

$$\chi_G(\lambda) = \lambda^{n_G} \cdot \chi_{M_G}(\lambda),$$

where n_G is the number of connected components of G . It follows that the coefficients of the chromatic polynomial of G form a log-concave sequence. \square

REFERENCES

- [AS14] Karim Adiprasito and Raman Sanyal, *Log-concavity of Whitney numbers via measure concentration*, preprint.
- [AB14] Karim Adiprasito and Anders Björner, *Filtered geometric lattices and Lefschetz Section Theorems over the tropical semiring*, arXiv:1401.7301.
- [AHK15] Karim Adiprasito, June Huh, Eric Katz, *Hodge theory for simplicial fans*, in preparation.
- [Aig87] Martin Aigner, *Whitney numbers*, Combinatorial Geometries, 139–160, Encyclopedia of Mathematics and its Applications **29**, Cambridge University Press, Cambridge, 1987.
- [Aig07] Martin Aigner, *A course in enumeration*, Graduate Texts in Mathematics **23**, Springer-Verlag, Berlin, 2007.
- [AK06] Federico Ardila and Caroline Klivans, *The Bergman complex of a matroid and phylogenetic trees*, Journal of Combinatorial Theory Series B **96** (2006), no. 1, 38–49.
- [BH15] Farhad Babaee and June Huh, *A tropical approach to the strongly positive Hodge conjecture*, arXiv:1502.00299.
- [BB11] Victor Batyrev and Mark Blume, *The functor of toric varieties associated with Weyl chambers and Losev-Manin moduli spaces*, Tohoku Mathematical Journal **63** (2011), no. 4, 581–604.
- [BDP90] Emili Bifet, Corrado De Concini, and Claudio Procesi, *Cohomology of regular embeddings*, Advances in Mathematics **82** (1990), no. 1, 1–34.
- [Bil89] Louis Billera, *The algebra of continuous piecewise polynomials*, Advances in Mathematics **76** (1989), no. 2, 170–183.
- [Bir12] George Birkhoff, *A determinant formula for the number of ways of coloring a map*, Annals of Mathematics **14** (1912), no. 1, 42–46.
- [Bjo92] Anders Björner, *The homology and shellability of matroids and geometric lattices*, Matroid Applications, 226–283, Encyclopedia of Mathematics and its Applications **40**, Cambridge University Press, Cambridge, 1992.
- [Bri96] Michel Brion, *Piecewise polynomial functions, convex polytopes and enumerative geometry*, Parameter spaces (Warsaw, 1994), 25–44, Banach Center Publications **36**, Polish Academy of Sciences, Warsaw, 1996.
- [Bry77] Thomas Brylawski, *The broken-circuit complex*, Transactions of the American Mathematical Society **234** (1977), no. 2, 417–433.
- [Bry86] Thomas Brylawski, *Constructions*, Theory of Matroids, 127–223, Encyclopedia of Mathematics and its Applications **26**, Cambridge University Press, Cambridge, 1986.
- [Cat08] Eduardo Cattani, *Mixed Lefschetz theorems and Hodge-Riemann bilinear relations*, International Mathematics Research Notices, Vol. 2008, no. 10, Article ID rnn025, 20 pp.

- [CM02] Mark de Cataldo and Luca Migliorini, *The hard Lefschetz theorem and the topology of semismall maps*, Annales Scientifiques de l'école Normale Supérieure **35** (2002), no. 5, 759–772.
- [Dan78] Vladimir Danilov, *The geometry of toric varieties*, Russian Mathematical Surveys **33** (1978), 97–154.
- [DP95] Corrado De Concini and Claudio Procesi, *Wonderful models of subspace arrangements*, Selecta Mathematica. New Series **1** (1995), no. 3, 459–494.
- [Koz08] Dimitry Kozlov, *Combinatorial algebraic topology*, Algorithms and Computation in Mathematics **21**, Springer-Verlag Berlin, 2008.
- [FY04] Eva Maria Feichtner and Sergey Yuzvinsky, *Chow rings of toric varieties defined by atomic lattices*, Inventiones Mathematicae **155** (2004), no. 3, 515–536.
- [Ful93] William Fulton, *Introduction to Toric Varieties*, Annals of Mathematics Studies, **131**, Princeton University Press, Princeton, NJ, 1993.
- [FS97] William Fulton and Bernd Sturmfels, *Intersection theory on toric varieties*, Topology **36** (1997), no. 2, 335–353.
- [GM92] Israel Gel'fand and Robert MacPherson, *A combinatorial formula for the Pontrjagin classes*, Bulletin of the American Mathematical Society (New Series) **26** (1992), no. 2, 304–309.
- [GM12] Angela Gibney and Diane Maclagan, *Lower and upper bounds on nef cones*, International Mathematics Research Notices (2012), no. 14, 3224–3255.
- [Her72] Andrew Heron, *Matroid polynomials*, Combinatorics (Proc. Conf. Combinatorial Math., Math. Inst., Oxford, 1972), Inst. of Math. and its Appl., Southend-on-Sea, 1972, pp. 164–202.
- [Hog74] Stuart Hoggar, *Chromatic polynomials and logarithmic concavity*, Journal of Combinatorial Theory Series B **16** (1974), 248–254.
- [Huh12] June Huh, *Milnor numbers of projective hypersurfaces and the chromatic polynomial of graphs*, Journal of the American Mathematical Society **25** (2012), 907–927.
- [HK12] June Huh and Eric Katz, *Log-concavity of characteristic polynomials and the Bergman fan of matroids*, Mathematische Annalen **354** (2012), no. 3, 1103–1116.
- [Hru92] Ehud Hrushovski, *Unimodular minimal structures*, Journal of the London Mathematical Society, II. **46** (1992), no.3, 385–396.
- [IKMZ] Ilya Itenberg, Ludmil Katzarkov, Grigory Mikhalkin and Ilya Zharkov, *Tropical homology*, in preparation.
- [KP11] Eric Katz and Sam Payne, *Realization spaces for tropical fans*, Combinatorial Aspects of Commutative Algebra and Algebraic Geometry, 73–88, Abel Symposium **6**, Springer, Berlin, 2011.
- [Kun95] Joseph Kung, *The geometric approach to matroid theory*, Gian-Carlo Rota on Combinatorics, 604–622, Contemporary Mathematicians, Birkhäuser Boston, Boston, MA, 1995.
- [Laz04] Robert Lazarsfeld, *Positivity in Algebraic Geometry I*, Ergebnisse der Mathematik und ihrer Grenzgebiete **48**, Springer-Verlag, Berlin, 2004.
- [Lee96] Carl Lee, *P.L.-spheres, convex polytopes, and stress*, Discrete and Computational Geometry **15** (1996), no. 4, 389–421.
- [Len12] Matthias Lenz, *The f-vector of a representable-matroid complex is log-concave*, Advances in Applied Mathematics **51** (2013), no. 5, 543–545.
- [Mas72] John Mason, *Matroids: unimodal conjectures and Motzkin's theorem*, Combinatorics (Proc. Conf. Combinatorial Math., Math. Inst., Oxford, 1972), pp. 207–220. Inst. Math. Appl., Southend-on-Sea, 1972.
- [McM93] Peter McMullen, *On simple polytopes*, Inventiones mathematicae **113** (1993), no. 2, 419–444.
- [MS05] Dagmar Meyer and Larry Smith, *Poincaré duality algebras, Macaulay's dual systems, and Steenrod operations*, Cambridge Tracts in Mathematics **167**, Cambridge University Press, Cambridge, 2005.
- [MS05] Ezra Miller and Bernd Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics **227**, Springer-Verlag, New York, 2005.
- [vN60] John von Neumann, *Continuous geometry*, Princeton Landmarks in Mathematics Princeton University Press, Princeton, NJ (1960).
- [Oxl11] James Oxley, *Matroid Theory*, Oxford Science Publications, Oxford University Press, New York, 2011.

- [Pil96] Anand Pillay, *Geometric stability theory*, Oxford Logic Guides **32**, The Clarendon Press, Oxford University Press, New York, 1996.
- [Rea68] Ronald Read, *An introduction to chromatic polynomials*, Journal of Combinatorial Theory **4** (1968), 52–71.
- [Rot64] Gian-Carlo Rota, *On the foundations of combinatorial theory. I. Theory of Möbius functions*. Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete **2** (1964), 340–368.
- [Rot71] Gian-Carlo Rota, *Combinatorial theory, old and new*, Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 3, Gauthier-Villars, Paris, 1971, pp. 229–233.
- [Sta07] Richard Stanley, *An introduction to hyperplane arrangements*, Geometric combinatorics, 389–496, IAS/Park City Mathematics Series, **13**, American Mathematical Society, Providence, RI, 2007.
- [Wel71] Dominic Welsh, *Combinatorial problems in matroid theory*, Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969) pp. 291–306, Academic Press, London, 1971.
- [Wel76] Dominic Welsh, *Matroid Theory*, London Mathematical Society Monographs, **8**, Academic Press, London-New York, 1976.
- [Whi87] Neil White, *Combinatorial geometries*, Encyclopedia of Mathematics and its Applications **29**, Cambridge University Press, Cambridge, 1987.
- [Whi32] Hassler Whitney, *A logical expansion in mathematics*, Bulletin of the American Mathematical Society **38**, no. 8 (1932), 572–579.
- [Whi35] Hassler Whitney, *On the abstract properties of linear dependence*. American Journal of Mathematics **57** (1935), no. 3, 509–533.
- [Zas87] Thomas Zaslavsky, *The Möbius function and the characteristic polynomial*, Combinatorial geometries, 114–138, Encyclopedia of Mathematics and its Applications **29**, Cambridge University Press, Cambridge, 1987.

EINSTEIN INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, ISRAEL

E-mail address: `adiprasito@math.huji.ac.il`

INSTITUTE FOR ADVANCED STUDY, FULD HALL, 1 EINSTEIN DRIVE, PRINCETON, NEW JERSEY, USA.

E-mail address: `huh@princeton.edu`, `juneuh@ias.edu`

DEPARTMENT OF COMBINATORICS & OPTIMIZATION, UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVENUE WEST, WATERLOO, ONTARIO, CANADA.

E-mail address: `eekatz@uwaterloo.ca`