

The Strichartz inequality for orthonormal functions

Rupert L. Frank
Caltech

Joint work with **Mathieu Lewin, Elliott Lieb** *and* **Robert Seiringer**

Strichartz inequality for orthonormal functions

Preprint: arXiv:1306.1309

Quantum Spectra And Transport, Jerusalem, July 1, 2013

INTRODUCTION – THE SCHRÖDINGER EQUATION

By spectral theory the solution $e^{-itH}\psi$ of the **time-dependent Schrödinger equation**

$$i\frac{\partial}{\partial t}\Psi = H\Psi, \quad \Psi|_{t=0} = \psi$$

with H self-adjoint satisfies $\|e^{-itH}\psi\| = \|\psi\|$ for all $t \in \mathbb{R}$.

Here we are interested in the phenomenon of **dispersion**.

Example: $H = -\Delta$ in $L^2(\mathbb{R}^d)$ and $\psi(x) = (\pi\sigma^2)^{-d/4} e^{ip \cdot x} e^{-x^2/2\sigma^2}$. Then

$$|(e^{it\Delta}\psi)(x)|^2 = \left(\frac{\sigma^2}{\pi(\sigma^4 + 4t^2)}\right)^{d/2} e^{-\sigma^2(x-2tp)^2/(\sigma^4 + 4t^2)}.$$

Dispersion is quantified by **Strichartz inequalities**. Simplest form:

$$\int_{\mathbb{R}} \int_{\mathbb{R}^d} |(e^{it\Delta}\psi)(x)|^{2(d+2)/d} dx dt \leq C_d \left(\int_{\mathbb{R}^d} |\psi(x)|^2 dx \right)^{(d+2)/d}.$$

Due to **Strichartz** (1977); see also **Lindblad–Sogge**, **Ginibre–Velo**, **Keel–Tao**, **Foschi**, ...

GOAL – A STRICHARTZ INEQUALITY FOR ORTHONORMAL FUNCTIONS

Is there an inequality for

$$\int_{\mathbb{R}} \int_{\mathbb{R}^d} \left(\sum_j |(e^{it\Delta} \psi_j)(x)|^2 \right)^{(d+2)/d} dx dt$$

with ψ_j orthonormal in $L^2(\mathbb{R}^d)$?

Obvious answer: By triangle inequality (without using orthogonality!)

$$\int_{\mathbb{R}} \int_{\mathbb{R}^d} \left(\sum_{j=1}^N |(e^{it\Delta} \psi_j)(x)|^2 \right)^{(d+2)/d} dx dt \leq C_d N^{(d+2)/d}$$

Can we do better than that?

Main result: Yes, we can!

$$\int_{\mathbb{R}} \int_{\mathbb{R}^d} \left(\sum_{j=1}^N |(e^{it\Delta} \psi_j)(x)|^2 \right)^{(d+2)/d} dx dt \leq C'_d N^{(d+1)/d}$$

And this is best possible!

COMPARE WITH LIEB–THIRRING INEQUALITIES

The **Sobolev interpolation inequality** says that for $\gamma \geq 1$ (and more)

$$\int_{\mathbb{R}^d} |\nabla \psi|^2 dx \geq S_{d,\gamma} \left(\int_{\mathbb{R}^d} |\psi|^2 dx \right)^{-\frac{\gamma-1}{d/2}} \left(\int_{\mathbb{R}^d} |\psi|^{\frac{2(\gamma+d/2)}{\gamma+d/2-1}} dx \right)^{\frac{\gamma+d/2-1}{d/2}} .$$

This was generalized by **Lieb–Thirring** (1976) to **orthonormal functions** ψ_j

$$\sum_{j=1}^N \int_{\mathbb{R}^d} |\nabla \psi_j|^2 dx \geq K_{d,\gamma} N^{-\frac{\gamma-1}{d/2}} \left(\int_{\mathbb{R}^d} \left(\sum_{j=1}^N |\psi_j|^2 \right)^{\frac{\gamma+d/2}{\gamma+d/2-1}} dx \right)^{\frac{\gamma+d/2-1}{d/2}} .$$

This is better than $N^{-\frac{\gamma}{d/2}}$ (from triangle inequality) and **optimal** in the semi-classical limit. Case $\gamma = 1$ is used in the Lieb–Thirring proof of **stability of matter**.

Slightly more precise version: for any operator $\Gamma \geq 0$ on $L^2(\mathbb{R}^d)$,

$$\mathrm{Tr}(-\Delta)\Gamma \geq K_{d,\gamma} \left(\mathrm{Tr} \Gamma^{\frac{\gamma}{\gamma-1}} \right)^{-\frac{\gamma-1}{d/2}} \left(\int_{\mathbb{R}^d} \Gamma(x, x)^{\frac{\gamma+d/2}{\gamma+d/2-1}} dx \right)^{\frac{\gamma+d/2-1}{d/2}} .$$

‘SEMI-CLASSICAL’ INTUITION BEHIND STRICHARTZ

Why is $\iint \left(\sum_{j=1}^N |(e^{it\Delta}\psi_j)(x)|^2 \right)^{(2+d)/d} dx dt \leq C'_d N^{(d+1)/d}$ **best possible?**

Heuristics: At $t = 0$ consider N electrons in a box of size L with const. density $\rho = L^{-d}N$. For $|t| \geq T$ the electrons have (approximately) disjoint supports and therefore

$$\iint_{|t| \geq T} \left(\sum_{j=1}^N |(e^{it\Delta}\psi_j)(x)|^2 \right)^{(2+d)/d} dx dt \approx N \ll N^{(d+1)/d}.$$

We think of T as the typical time it takes an electron to move a distance comparable with the size of the system. By **Thomas–Fermi theory** the expected momentum per particle is $\approx \rho^{1/d}$ and therefore, if the electrons move **ballistically** $T \approx L\rho^{-1/d}$. Thus,

$$\iint_{|t| \leq T} \left(\sum_{j=1}^N |(e^{it\Delta}\psi_j)(x)|^2 \right)^{(2+d)/d} dx dt \approx TL^d \rho^{(2+d)/d} \approx N^{(d+1)/d}.$$

THE MAIN RESULT

Theorem 1. *Let $d \geq 1$ and assume that $1 < p, q < \infty$ satisfy*

$$1 < q \leq 1 + \frac{2}{d} \quad \text{and} \quad \frac{2}{p} + \frac{d}{q} = d.$$

*Then, for any **orthonormal** ψ_j and any $n_j \in \mathbb{C}$*

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}^d} \left| \sum_j n_j |(e^{it\Delta}\psi_j)(x)|^2 \right|^q dx \right)^{\frac{p}{q}} dt \leq C_{d,q}^p \left(\sum_j |n_j|^{\frac{2q}{q+1}} \right)^{\frac{p(q+1)}{2q}}. \quad (1)$$

that is, with the notations $\gamma(t) = e^{it\Delta}\gamma e^{-it\Delta}$ and $\rho_\gamma(x) = \gamma(x, x)$,

$$\|\rho_{\gamma(t)}\|_{L_t^p(\mathbb{R}, L_x^q(\mathbb{R}^d))} \leq C_{d,q} \|\gamma\|_{\mathfrak{S}^{\frac{2q}{q+1}}}.$$

This is best possible in the sense that

$$\sup_{\gamma} \frac{\|\rho_{\gamma(t)}\|_{L_t^p(\mathbb{R}, L_x^q(\mathbb{R}^d))}}{\|\gamma\|_{\mathfrak{S}^r}} = \infty \quad \text{if } r > \frac{2q}{q+1}.$$

REMARKS

Recall

$$\|\rho_\gamma(t)\|_{L_t^p(\mathbb{R}, L_x^q(\mathbb{R}^d))} \leq C_{d,q} \|\gamma\|_{\mathfrak{S}^{\frac{2q}{q+1}}} \quad \text{if } 1 < q \leq 1 + \frac{2}{d}.$$

Remarks. (1) The inequality with the **trace norm** $\|\gamma\|_{\mathfrak{S}^1}$ on the right side is known, even for the full range $1 \leq p, q \leq \infty$ with $(p, q, d) \neq (1, \infty, 2)$ (plus scaling condition).

(2) This implies an **inhomogeneous Strichartz inequality**: if

$$i\dot{\gamma}(t) = [-\Delta, \gamma(t)] + iR(t), \quad \gamma(t_0) = 0,$$

with $R(t)$ self-adjoint, then for q as in our theorem

$$\|\rho_\gamma(t)\|_{L_t^p(\mathbb{R}, L_x^q(\mathbb{R}^d))} \leq C \left\| \int_{\mathbb{R}} e^{-is\Delta} |R(s)| e^{is\Delta} ds \right\|_{\mathfrak{S}^{\frac{2q}{q+1}}}.$$

(3) We prove that the inequality **fails** for $q \geq (d+1)/(d-1)$ and **conjecture** that it holds even for $1 + 2/d < q < (d+1)/(d-1)$. At the **endpoint** $q = (d+1)/(d-1)$ we conjecture that

$$\|\rho_\gamma(t)\|_{L_t^p(\mathbb{R}, L_x^q(\mathbb{R}^d))} \leq C'_d \|\gamma\|_{\mathfrak{S}^{\frac{2q}{q+1}, \mathbf{1}}}.$$

THE DUAL FORMULATION

Using **Hölder's inequality** (for operators and for functions) and the fact that

$$\iint_{\mathbb{R} \times \mathbb{R}^d} V(t, x) \rho_{\gamma(t)}(x) dx dt = \text{Tr } \gamma \int_{\mathbb{R}} e^{-it\Delta} V(t, \cdot) e^{it\Delta} dt$$

we see that Theorem 1 is **equivalent** to

Theorem 2. *Let $d \geq 1$ and assume that $1 < p', q' < \infty$ satisfy*

$$1 + \frac{d}{2} \leq q' < \infty \quad \text{and} \quad \frac{2}{p'} + \frac{d}{q'} = 2.$$

Then, with the same constant as in Theorem 1,

$$\left\| \int_{\mathbb{R}} e^{-it\Delta} V(t, \cdot) e^{it\Delta} dt \right\|_{\mathfrak{S}^{2q'}} \leq C_{d,q} \|V\|_{L_t^{p'}(\mathbb{R}, L_x^{q'}(\mathbb{R}^d))}.$$

By **interpolation** it suffices to prove this for $q' = p' = 1 + d/2$.

PROOF OF THEOREM 2

For $V \geq 0$,

$$\begin{aligned} \left\| \int_{\mathbb{R}} e^{-it\Delta} V(t, \cdot) e^{it\Delta} dt \right\|_{\mathfrak{S}^{d+2}}^{d+2} &= \text{Tr} \left(\int_{\mathbb{R}} e^{-it\Delta} V(t, \cdot) e^{it\Delta} dt \right)^{d+2} \\ &= \int_{\mathbb{R}} \cdots \int_{\mathbb{R}} \text{Tr} V(t_1, x + 2t_1 p) \cdots V(t_{d+2}, x + 2t_{d+2} p) dt_{d+2} \cdots dt_1 \end{aligned}$$

Here we use the **notation**

$$f(x + 2tp) = e^{-it\Delta} f(x) e^{it\Delta}.$$

Lemma 3 (Generalized Kato–Simon–Seiler ineq.). For $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ and $r \geq 2$,

$$\|f(\alpha x + \beta p) g(\gamma x + \delta p)\|_{\mathfrak{S}^r} \leq \frac{\|f\|_{L^r(\mathbb{R}^d)} \|g\|_{L^r(\mathbb{R}^d)}}{(2\pi)^{\frac{d}{r}} |\alpha\delta - \beta\gamma|^{\frac{d}{r}}}.$$

Thus,

$$\left| \text{Tr} \left(V(t_1, x + 2t_1 p) \cdots V(t_{d+2}, x + 2t_{d+2} p) \right) \right| \leq \frac{\|V(t_1, \cdot)\|_{L_x^{1+d/2}} \cdots \|V(t_{d+2}, \cdot)\|_{L_x^{1+d/2}}}{(4\pi)^d |t_1 - t_2|^{\frac{d}{d+2}} \cdots |t_{d+2} - t_1|^{\frac{d}{d+2}}}$$

PROOF OF THEOREM 2, CONT'D

We have shown that

$$\left\| \int_{\mathbb{R}} e^{-it\Delta} V(t, \cdot) e^{it\Delta} dt \right\|_{\mathfrak{S}^{d+2}}^{d+2} \leq \int_{\mathbb{R}} \cdots \int_{\mathbb{R}} \frac{\|V(t_1, \cdot)\|_{L_x^{1+d/2}} \cdots \|V(t_{d+2}, \cdot)\|_{L_x^{1+d/2}}}{(4\pi)^d |t_1 - t_2|^{\frac{d}{d+2}} \cdots |t_{d+2} - t_1|^{\frac{d}{d+2}}} dt_{d+2} \cdots dt_1$$

Lemma 4 (Multi-linear HLS inequality; Christ, Beckner). *Assume that $(\beta_{ij})_{1 \leq i, j \leq N}$ and $(r_k)_{1 \leq k \leq N}$ are real-numbers such that*

$$\beta_{ii} = 0, \quad 0 \leq \beta_{ij} = \beta_{ji} < 1, \quad r_k > 1, \quad \sum_{k=1}^N \frac{1}{r_k} > 1, \quad \sum_{i=1}^N \beta_{ik} = \frac{2(r_k - 1)}{r_k}.$$

Then

$$\left| \int_{\mathbb{R}} \cdots \int_{\mathbb{R}} \frac{f_1(t_1) \cdots f_N(t_N)}{\prod_{i < j} |t_i - t_j|^{\beta_{ij}}} dt_N \cdots dt_1 \right| \leq C \prod_{k=1}^N \|f_k\|_{L^{r_k}(\mathbb{R})}.$$

For us, $N = d + 2$, $\beta_{ij} = \delta_{j, i+1} d / (d + 2)$ and $r_k = 1 + d/2$ and thus

$$\left\| \int_{\mathbb{R}} e^{-it\Delta} V(t, \cdot) e^{it\Delta} dt \right\|_{\mathfrak{S}^{d+2}}^{d+2} \leq C \|V\|_{L_{t,x}^{1+d/2}}^{d+2}. \quad \square$$

AN APPLICATION

Consider the **unitary propagator** $U_V(t, t_0)$ satisfying

$$i \frac{\partial}{\partial t} U_V(t, t_0) = (-\Delta + V(t, x)) U_V(t, t_0), \quad U_V(t_0, t_0) = 1,$$

and the **wave operator**

$$\mathcal{W}_V(t, t_0) := U_0(t_0, t) U_V(t, t_0) = e^{i(t_0 - t)\Delta} U_V(t, t_0). \quad (2)$$

The wave operator can be formally expanded in a **Dyson series**.

Theorem 5. *Let $d \geq 1$ and assume that $1 < p', q' < \infty$ satisfy*

$$1 + \frac{d}{2} \leq q' < \infty \quad \text{and} \quad \frac{2}{p'} + \frac{d}{q'} = 2.$$

If $V \in L_t^{p'}(\mathbb{R}, L_x^{q'}(\mathbb{R}^d))$, then $\lim_{t \rightarrow \pm\infty} \mathcal{W}_V(t, t_0) - 1 \in \mathfrak{S}^{2q'}$ and the Dyson series converges in $\mathfrak{S}^{2q'}$.

Improves parts of results of **Howland, Yajima, Jensen, ...**

HAPPY BIRTHDAY, YOSI!

AND

THANK YOU FOR YOUR ATTENTION!