


Spectral Transition for Random Quantum Walks on Trees*

Alain JOYE



* Joint work with Eman HAMZA, Cairo University

Warm Up: Quantum Walk on $\mathbb{Z} = \mathcal{T}_2$

Quantum particle with spin 1/2 on 1-dim lattice, i.e. $\mathcal{K}_2 = l^2(\mathbb{Z}) \otimes \mathbb{C}^2$

Spin evol.: C unitary op. on \mathbb{C}^2 , “coin” space

Spin dependent shift:

Let S_{\pm} shift to the right/left on $l^2(\mathbb{Z})$, $|\pm\rangle\langle\pm|$ proj. on $|\pm\rangle \in \mathbb{C}^2$

$S := S_+ \otimes |+\rangle\langle+| + S_- \otimes |-\rangle\langle-|$ on $l^2(\mathbb{Z}) \otimes \mathbb{C}^2$

One step dynamics: $U := S(\mathbb{I} \otimes C)$ s.t.

$$U|x \otimes \pm\rangle = \langle -|C \pm\rangle|(x-1) \otimes -\rangle + \langle +|C \pm\rangle|(x+1) \otimes +\rangle$$

Warm Up: Quantum Walk on $\mathbb{Z} = \mathcal{T}_2$

Quantum particle with spin 1/2 on 1-dim lattice, i.e. $\mathcal{K}_2 = l^2(\mathbb{Z}) \otimes \mathbb{C}^2$

Spin evol.: C unitary op. on \mathbb{C}^2 , “coin” space

Spin dependent shift:

Let S_{\pm} shift to the right/left on $l^2(\mathbb{Z})$, $|\pm\rangle\langle\pm|$ proj. on $|\pm\rangle \in \mathbb{C}^2$

$S := S_+ \otimes |+\rangle\langle+| + S_- \otimes |-\rangle\langle-|$ on $l^2(\mathbb{Z}) \otimes \mathbb{C}^2$

One step dynamics: $U := S(\mathbb{I} \otimes C)$ s.t.

$$U|x \otimes \pm\rangle = \langle -|C \pm\rangle|(x-1) \otimes -\rangle + \langle +|C \pm\rangle|(x+1) \otimes +\rangle$$

Analogy:

If $H = -\Delta + V$ then $U = e^{-iH}$ suggests $S \simeq e^{-i\Delta}$ and $C \simeq e^{iV}$

Warm Up: Quantum Walk on $\mathbb{Z} = \mathcal{T}_2$

Quantum particle with spin 1/2 on 1-dim lattice, i.e. $\mathcal{K}_2 = l^2(\mathbb{Z}) \otimes \mathbb{C}^2$

Spin evol.: C unitary op. on \mathbb{C}^2 , “coin” space

Spin dependent shift:

Let S_{\pm} shift to the right/left on $l^2(\mathbb{Z})$, $|\pm\rangle\langle\pm|$ proj. on $|\pm\rangle \in \mathbb{C}^2$

$$S := S_+ \otimes |+\rangle\langle+| + S_- \otimes |-\rangle\langle-| \text{ on } l^2(\mathbb{Z}) \otimes \mathbb{C}^2$$

One step dynamics: $U := S(\mathbb{I} \otimes C)$ s.t.

$$U|x \otimes \pm\rangle = \langle -|C \pm\rangle|(x-1) \otimes -\rangle + \langle +|C \pm\rangle|(x+1) \otimes +\rangle$$

Analogy:

If $H = -\Delta + V$ then $U = e^{-iH}$ suggests $S \simeq e^{-i\Delta}$ and $C \simeq e^{iV}$

Random Quantum Walk:

$$C \rightsquigarrow \{C_{\omega}(x)\}_{x \in \mathbb{Z}}, \text{ i.i.d. set of } C_{\omega}(x) \in U(2)$$

$$U \rightsquigarrow U_{\omega}|x \otimes \pm\rangle = \langle -|C_{\omega}(x)|\pm\rangle|(x-1) \otimes -\rangle + \langle +|C_{\omega}(x)|\pm\rangle|(x+1) \otimes +\rangle$$

Homogeneous tree \mathcal{T}_3 with coord. number $q = 3$

$A_3 = \{a, b, c\}$ generators of a group with unit e s.t. $a^2 = b^2 = c^2 = e$.

Root: Origin e

Edges: Labels $\{a, b, c\}$ in trig. order


Any $x \in \mathcal{T}_3$: Finite reduced word

$$x = x_1 x_2 \cdots x_n, x_j \in A_3$$

Length: $|x| = n$

Distance: For any $x, y \in \mathcal{T}_3$,

$$d(x, y) = |x^{-1}y|$$



Homogeneous tree \mathcal{T}_3 with coord. number $q = 3$

$A_3 = \{a, b, c\}$ generators of a group with unit e s.t. $a^2 = b^2 = c^2 = e$.

Root: Origin e

Edges: Labels $\{a, b, c\}$ in trig. order

Any $x \in \mathcal{T}_3$: Finite reduced word

$$x = x_1 x_2 \cdots x_n, x_j \in A_3$$


Length: $|x| = n$

Distance: For any $x, y \in \mathcal{T}_3$,

$$d(x, y) = |x^{-1}y|$$

$l^2(\mathcal{T}_3)$:

$x \mapsto |x\rangle$, s.t. $\{|x\rangle\}_{x \in \mathcal{T}_3}$ ONB of $l^2(\mathcal{T}_3)$



$$\psi = \sum_{x \in \mathcal{T}_3} \psi(x) |x\rangle, \quad \sum_{x \in \mathcal{T}_3} |\psi(x)|^2 < \infty.$$

Shifts on \mathcal{T}_3

Even / Odd: Sites x_e / x_o s.t. $|x_e|$ even / $|x_o|$ odd

Let $S_{ab} : l^2(\mathcal{T}_3) \rightarrow l^2(\mathcal{T}_3)$

$$S_{a b}|x_e\rangle = |x_e a\rangle$$

$$S_{a b}|x_o\rangle = |x_o b\rangle$$

Shifts on \mathcal{T}_3

Even / Odd: Sites x_e / x_o s.t. $|x_e|$ even / $|x_o|$ odd

Let $S_{ab} : l^2(\mathcal{T}_3) \rightarrow l^2(\mathcal{T}_3)$


$$S_{a\bar{b}}|x_e\rangle = |x_e a\rangle$$

$$S_{a\bar{b}}|x_o\rangle = |x_o b\rangle$$

Cyclic subspaces:

$$\mathcal{H}_x^{ab} = \text{span} \{ S_{ab}^n |x\rangle, n \in \mathbb{Z} \}$$

$$S_{ab}|_{\mathcal{H}_x^{ab}} \simeq \text{shift on } l^2(\mathbb{Z}).$$



Shifts on \mathcal{T}_3

Even / Odd: Sites x_e / x_o s.t. $|x_e|$ even / $|x_o|$ odd

Let $S_{ab} : l^2(\mathcal{T}_3) \rightarrow l^2(\mathcal{T}_3)$


$$S_{ab}|x_e\rangle = |x_e a\rangle$$

$$S_{ab}|x_o\rangle = |x_o b\rangle$$

Cyclic subspaces:

$$\mathcal{H}_x^{ab} = \text{span} \{ S_{ab}^n |x\rangle, n \in \mathbb{Z} \}$$

$$S_{ab}|_{\mathcal{H}_x^{ab}} \simeq \text{shift on } l^2(\mathbb{Z}).$$



Spectrum:

$$\sigma(S_{ab}) = \sigma_{ac}(S_{ab}) = \{z \text{ s.t. } |z| = 1\}$$

Similarly for S_{bc} , S_{ca}

Quantum Walk on \mathcal{T}_3

- Unitary evolution:

Particle with spin 1 on \mathcal{T}_3 jumping on nearest neighbors

Hilbert space: $\mathcal{K}_3 = l^2(\mathcal{T}_3) \otimes \mathbb{C}^3$

ONB of \mathbb{C}^3 : $\{|a\rangle, |b\rangle, |c\rangle\}$,

ONB of \mathcal{K}_3 : $\{\underline{x} \otimes \tau = |x\rangle \otimes |\tau\rangle\}_{x \in \mathcal{T}_3, \tau \in A_3}$

- Spin dep. shift on \mathcal{K}_3 :

$$\underline{S} = S_{bc} \otimes |a\rangle\langle a| + S_{ca} \otimes |b\rangle\langle b| + S_{ab} \otimes |c\rangle\langle c|$$

- Spin update: For $\underline{C} \in U(3)$ a unitary op. on \mathbb{C}^3

$$\mathbb{I} \otimes C : \mathcal{K}_3 \rightarrow \mathcal{K}_3$$

Quantum Walk on \mathcal{T}_3

- Unitary evolution:

Particle with spin 1 on \mathcal{T}_3 jumping on nearest neighbors

Hilbert space: $\mathcal{K}_3 = l^2(\mathcal{T}_3) \otimes \mathbb{C}^3$

ONB of \mathbb{C}^3 : $\{|a\rangle, |b\rangle, |c\rangle\}$,

ONB of \mathcal{K}_3 : $\{\mathbf{x} \otimes \tau = |x\rangle \otimes |\tau\rangle\}_{x \in \mathcal{T}_3, \tau \in A_3}$

- Spin dep. shift on \mathcal{K}_3 :

$$S = S_{bc} \otimes |a\rangle\langle a| + S_{ca} \otimes |b\rangle\langle b| + S_{ab} \otimes |c\rangle\langle c|$$

- Spin update: For $C \in U(3)$ a unitary op. on \mathbb{C}^3

$$\mathbb{I} \otimes C : \mathcal{K}_3 \rightarrow \mathcal{K}_3$$

- Time one dynamics of the QW:

$$U(C) := S(\mathbb{I} \otimes C)$$

s.t.

$$U(C)\mathbf{x}_e \otimes a = C_{aa}\mathbf{x}_e b \otimes a + C_{ba}\mathbf{x}_e c \otimes b + C_{ca}\mathbf{x}_e a \otimes c$$

$$U(C)\mathbf{x}_o \otimes a = C_{aa}\mathbf{x}_o c \otimes a + C_{ba}\mathbf{x}_o a \otimes b + C_{ca}\mathbf{x}_o b \otimes c \quad \text{etc...}$$

Random Environment \Rightarrow RQW

Spatial disorder

Different coin op. C_ω at each site:

$$C \mapsto \{C_\omega(x)\}_{x \in \mathcal{T}_3}$$

with

$$C_\omega(x_e)_{\tau, \sigma} = \exp(i\omega_{x_e \tau}^\tau) C_{\tau, \sigma},$$

$$C_\omega(x_o)_{\tau, \sigma} = \exp(i\omega_{x_o \sigma}^\tau) C_{\tau, \sigma}, \quad \tau, \sigma \in \{a, b, c\}.$$

- Random Quantum Walk:

$$U(C) \mapsto U_\omega(C)$$

Property: Set $\mathbb{D}(\omega) = \text{diag}(\exp(i\omega_x^\tau))$, then random time one dynamics

$$U_\omega(C) = \mathbb{D}(\omega)U(C)$$

Random Environment \Rightarrow RQW

Spatial disorder

Different coin op. C_ω at each site:

$$C \mapsto \{C_\omega(x)\}_{x \in \mathcal{T}_3} \quad \text{with} \quad \begin{aligned} C_\omega(x_e)_{\tau, \sigma} &= \exp(i\omega_{x_e \tau}^\tau) C_{\tau, \sigma}, \\ C_\omega(x_o)_{\tau, \sigma} &= \exp(i\omega_{x_o \sigma}^\tau) C_{\tau, \sigma}, \quad \tau, \sigma \in \{a, b, c\}. \end{aligned}$$

- Random Quantum Walk:

$$U(C) \mapsto U_\omega(C)$$

Property: Set $\mathbb{D}(\omega) = \text{diag}(\exp(i\omega_x^\tau))$, then random time one dynamics

$$U_\omega(C) = \mathbb{D}(\omega)U(C)$$

Assumptions:

- $\{\omega_x^\tau\}_{x \in \mathcal{T}_3}^{\tau \in A_3}$ are i.i.d. \mathbb{T} -valued random variables
 $\omega_x^\tau(\omega) \simeq d\mu(\theta) = l(\theta)d\theta$ with $l \in L^\infty$

Random Environment \Rightarrow RQW

Spatial disorder Different coin op. C_ω at each site:

$$C \mapsto \{C_\omega(x)\}_{x \in \mathcal{T}_3} \quad \text{with} \quad \begin{aligned} C_\omega(x_e)_{\tau, \sigma} &= \exp(i\omega_{x_e \tau}^\tau) C_{\tau, \sigma}, \\ C_\omega(x_o)_{\tau, \sigma} &= \exp(i\omega_{x_o \sigma}^\tau) C_{\tau, \sigma}, \quad \tau, \sigma \in \{a, b, c\}. \end{aligned}$$

- Random Quantum Walk:

$$U(C) \mapsto U_\omega(C)$$

Property: Set $\mathbb{D}(\omega) = \text{diag}(\exp(i\omega_x^\tau))$, then random time one dynamics

$$U_\omega(C) = \mathbb{D}(\omega)U(C)$$

Assumptions:

- $\{\omega_x^\tau\}_{x \in \mathcal{T}_3}^{\tau \in A_3}$ are i.i.d. \mathbb{T} -valued random variables
 $\omega_x^\tau(\omega) \simeq d\mu(\theta) = l(\theta)d\theta$ with $l \in L^\infty$

Remarks:

- The transition prob. $|\langle x \otimes \tau | U_\omega(C) | y \otimes \sigma \rangle|^2$ are deterministic.
- Spectral transition expected between "large" and "small" disorder regimes,
Abou-Chakra, Anderson, Thouless '73, Kunz Souillard '83, Klein '94
- Used in Quantum Dynamics, Quantum Computing, Quantum Probability, ...

Landmarks in $U(3)$

Permutation mat. For $\pi \in \mathfrak{S}_3$ on $A_3 = \{a, b, c\}$ set $C_\pi = \sum_{\tau \in A_3} |\pi(\tau)\rangle\langle\tau|$

Consider

$$U_\omega(C_\pi) = \mathbb{D}(\omega)U(C_\pi)$$

Landmarks in $U(3)$

Permutation mat. For $\pi \in \mathfrak{S}_3$ on $A_3 = \{a, b, c\}$ set $C_\pi = \sum_{\tau \in A_3} |\pi(\tau)\rangle\langle\tau|$

Consider

$$U_\omega(C_\pi) = \mathbb{D}(\omega)U(C_\pi)$$

- $C_{(a)(b)(c)} = \mathbb{I}$ $\Rightarrow U_\omega(\mathbb{I}) = \mathbb{D}(\omega)S \simeq S$ a.c. \leftrightarrow deloc.

Landmarks in $U(3)$

Permutation mat. For $\pi \in \mathfrak{S}_3$ on $A_3 = \{a, b, c\}$ set $C_\pi = \sum_{\tau \in A_3} |\pi(\tau)\rangle\langle\tau|$

Consider

$$U_\omega(C_\pi) = \mathbb{D}(\omega)U(C_\pi)$$

- $C_{(a)(b)(c)} = \mathbb{I}$ $\Rightarrow U_\omega(\mathbb{I}) = \mathbb{D}(\omega)S \simeq S$ a.c. \leftrightarrow deloc.

- $C_{(abc)} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$ $\Rightarrow U_\omega(C_{(abc)})$ p.p. \leftrightarrow loc. (as is $U_\omega(C_{(acb)})$)

$$\mathcal{H}_{x_0} = \text{span}\{x_0 \otimes a, x_0a \otimes b, x_0 \otimes c, x_0c \otimes a, x_0 \otimes b, x_0b \otimes c\}$$

invariant under $U_\omega(C_{(abc)})$, $\forall x_o \in \mathcal{T}_3$

Landmarks in $U(3)$

Permutation mat. For $\pi \in \mathfrak{S}_3$ on $A_3 = \{a, b, c\}$ set $C_\pi = \sum_{\tau \in A_3} |\pi(\tau)\rangle\langle\tau|$

Consider

$$U_\omega(C_\pi) = \mathbb{D}(\omega)U(C_\pi)$$

- $C_{(a)(b)(c)} = \mathbb{I}$ $\Rightarrow U_\omega(\mathbb{I}) = \mathbb{D}(\omega)S \simeq S$ a.c. \leftrightarrow deloc.

- $C_{(abc)} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$ $\Rightarrow U_\omega(C_{(abc)})$ p.p. \leftrightarrow loc. (as is $U_\omega(C_{(acb)})$)

$$\mathcal{H}_{x_0} = \text{span}\{x_0 \otimes a, x_0a \otimes b, x_0 \otimes c, x_0c \otimes a, x_0 \otimes b, x_0b \otimes c\}$$

invariant under $U_\omega(C_{(abc)})$, $\forall x_o \in \mathcal{T}_3$


- $C_{(c)(ab)} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ $\Rightarrow U_\omega(C_{(c)(ab)}) \simeq$ p.p. \oplus a.c. \leftrightarrow mixed
(as are $U_\omega(C_{(a)(bc)})$, $U_\omega(C_{(b)(ac)})$)

$$\mathcal{H}_{x_0} = \text{span}\{\dots, x_0ba \otimes a, x_0b \otimes b, x_0 \otimes a, x_0a \otimes b, x_0ab \otimes a, \dots\}$$

$$\mathcal{H}_{x_e} = \text{span}\{x_e \otimes a, x_ec \otimes b\} \text{ invar. under } U_\omega(C_{(ab)(c)}), \forall x_e, x_o \in \mathcal{T}_3$$

Spectral Phase Diagram

Theorem:



Interpolating Matrices I

Localizing Matrices: For $0 \leq r \leq 1$ and $t = \sqrt{1 - r^2}$,

$$C_1^l(r) = \begin{pmatrix} 0 & r & t \\ 1 & 0 & 0 \\ 0 & t & -r \end{pmatrix} \text{ s.t. } C_1^l(\mathbf{1}) \simeq C_{(\mathbf{c})(ab)} \text{ and } C_1^l(\mathbf{0}) = C_{(abc)}$$


and similar $C_2^l(r), C_3^l(r), \dots, C_6^l(r)$.

Interpolating Matrices I

Localizing Matrices: For $0 \leq r \leq 1$ and $t = \sqrt{1 - r^2}$,

$$C_1^l(r) = \begin{pmatrix} 0 & r & t \\ 1 & 0 & 0 \\ 0 & t & -r \end{pmatrix} \text{ s.t. } C_1^l(1) \simeq C_{(c)(ab)} \text{ and } C_1^l(0) = C_{(abc)}$$

and similar $C_2^l(r), C_3^l(r), \dots, C_6^l(r)$.



For all $x_e \in \mathcal{T}_3$, let $\mathcal{H}_{x_e \otimes a}$ be the $U_\omega(C_1^l(r))$ -cyclic subsp. for $x_e \otimes a$,

Reduction to 1D model

Property

For each $x_e \in \mathcal{T}_3$, $U_{\omega}(C_1^l(r))|_{\mathcal{H}_{x_e} \otimes a} \simeq V_{\omega}(r)$

where

$$V_{\omega}(r) = \mathbb{D}(\omega) \left(\begin{array}{ccccccc} \ddots & & 0 & & & & \\ 0 & 0 & & & 1 & & \\ 0 & 0 & 0 & t & 0 & r & \\ r & -t & 0 & 0 & 0 & 0 & \\ & & 0 & 0 & 0 & 0 & t & r \\ & & 1 & 0 & 0 & 0 & 0 & 0 \\ & & r & & -t & 0 & 0 & \\ & & & & & 0 & 0 & \\ & & & & & 0 & 0 & t \\ & & & & & r & -t & 0 & \ddots \end{array} \right)$$

Allows for Transfer Matrix based analysis that yields localization $\forall r \in [0, 1]$.

Interpolating Matrices II

Delocalizing Matrices: For $0 \leq r \leq 1$ and $t = \sqrt{1 - r^2}$,

$$C_1^d(r) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & r & t \\ 0 & -t & r \end{pmatrix} \text{ s.t. } C_1^d(\mathbf{0}) \simeq C_{(a)(bc)} \text{ and } C_1^d(\mathbf{1}) = C_{(a)(b)(c)} = \mathbb{I}$$

and similar $C_2^d(r), C_3^d(r)$.

Interpolating Matrices II

Delocalizing Matrices: For $0 \leq r \leq 1$ and $t = \sqrt{1 - r^2}$,

$$C_1^d(r) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & r & t \\ 0 & -t & r \end{pmatrix} \text{ s.t. } C_1^d(0) \simeq C_{(a)(bc)} \text{ and } C_1^d(1) = C_{(a)(b)(c)} = \mathbb{I}$$

and similar $C_2^d(r), C_3^d(r)$.

Prop: $U_\omega(C_1^d(r))$ is purely ac. $\forall r \in (0, 1]$, and $\forall \omega$

Interpolating Matrices II

Delocalizing Matrices: For $0 \leq r \leq 1$ and $t = \sqrt{1 - r^2}$,

$$C_1^d(r) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & r & t \\ 0 & -t & r \end{pmatrix} \text{ s.t. } C_1^d(0) \simeq C_{(a)(bc)} \text{ and } C_1^d(1) = C_{(a)(b)(c)} = \mathbb{I}$$

and similar $C_2^d(r), C_3^d(r)$.

Prop: $U_\omega(C_1^d(r))$ is purely ac. $\forall r \in (0, 1]$, and $\forall \omega$

Path counting argument:

$\sum_{n \in \mathbb{Z}} |\langle x \otimes \tau | U^n x \otimes \tau \rangle|^2 < \infty \Rightarrow x \otimes \tau \in \text{a.c. spectral subspace of } U.$

$$\langle x \otimes \tau | U^n x \otimes \tau \rangle = \sum_{\substack{y_j \in \mathcal{T}_3 \\ \sigma_j \in A_3}} \langle x \otimes \tau | U y_1 \otimes \sigma_1 \rangle \langle y_1 \otimes \sigma_1 | U y_2 \otimes \sigma_2 \rangle \cdots \langle y_{n-1} \otimes \sigma_{n-1} | U x \otimes \tau \rangle$$

Spin $|b\rangle$ or $|c\rangle$:

s.t. $x_e \mapsto x_e c$ or $x_e a$ whereas $x_o \mapsto x_o a$ or $x_o b$.

Only possibility for return in $2n$ steps $x \mapsto xaaa\dots a \rightsquigarrow \sum_{n \in \mathbb{N}} t^{2n} < \infty$.

Neighborhood of $C_{(a)(b)(c)} = \mathbb{I}$

Perturbation: $C = \mathbb{I} + E$, with $\|E\| \leq \epsilon$

$$C_{\tau\tau} = O(1), \quad C_{\tau\sigma} = O(\epsilon), \quad \tau \neq \sigma$$

Path counting argument to show $\langle x \otimes \tau | U_{\omega}^n(C) | x \otimes \tau \rangle \in l^2(\mathbb{Z})$ if ϵ small

Expansion \rightsquigarrow

Path: $x = x \ y_1 \ y_2 \ y_3 \ \cdots \ y_{2n} \leftrightarrow$ Weight: $|C_{\tau\sigma_1} C_{\sigma_1\sigma_2} \cdots C_{\sigma_{2n}\tau}|$

Neighborhood of $C_{(a)(b)(c)} = \mathbb{I}$

Perturbation: $C = \mathbb{I} + E$, with $\|E\| \leq \epsilon$

$$C_{\tau\tau} = O(1), \quad C_{\tau\sigma} = O(\epsilon), \quad \tau \neq \sigma$$

Path counting argument to show $\langle x \otimes \tau | U_{\omega}^n(C) | x \otimes \tau \rangle \in l^2(\mathbb{Z})$ if ϵ small

Expansion \rightsquigarrow

$$\text{Path: } x = x \ y_1 \ y_2 \ y_3 \ \cdots \ y_{2n} \leftrightarrow \text{Weight: } |C_{\tau\sigma_1} C_{\sigma_1\sigma_2} \cdots C_{\sigma_{2n}\tau}|$$

Lemma:

Weight of length $2n$ path from x to x contains at least n off diag. terms

Argument:

Strings with of m consecutive diag. elemnts cannot reduce one another
 \Rightarrow the need for enough off-diag. elemnts to ensure reduction.

Then: Weight $\leq \epsilon^n$, $\#\{\text{contrib. paths}\} \leq k^n$, $k \simeq 72 \Rightarrow$

$$\sum_{n \in \mathbb{N}} |\langle x \otimes \tau | U_{\omega}^n(C) | x \otimes \tau \rangle|^2 \leq \sum_{n \in \mathbb{N}} (\epsilon k)^n < \infty, \text{ if } \epsilon > 0 \text{ small enough.}$$

Neighborhoods of $C_{(abc)}$ and $C_{(acb)}$

Theorem

Let $\pi \in \{(abc), (acb)\}$, C_π and $U_\omega(C)$ be as above and $\mathbb{U} := \{|z| = 1\}$.

For all $\gamma > 0$, there exists $\delta > 0$, $K < \infty$, s.t. $\forall C \in U(3)$, $\|C - C_\pi\| < \delta \implies \forall x, y \in \mathcal{T}_3$ and $\forall \sigma, \tau \in A_3$

$$\mathbb{E} \left[\sup_{f \in C(\mathbb{U}), \|f\|_\infty \leq 1} |\langle x \otimes \tau | f(U_\omega(C)) y \otimes \sigma \rangle| \right] \leq K e^{-\gamma d(x, y)}$$

à la "Aizenman-Molchanov" '09 Hamza-J.-Stolz
Similar approach Asch, Bourget, J. 11', J. '12

Neighborhoods of $C_{(abc)}$ and $C_{(acb)}$

Theorem Let $\pi \in \{(abc), (acb)\}$, C_π and $U_\omega(C)$ be as above and $\mathbb{U} := \{|z| = 1\}$.

For all $\gamma > 0$, there exists $\delta > 0$, $K < \infty$, s.t. $\forall C \in U(3)$,
 $\|C - C_\pi\| < \delta \implies \forall x, y \in \mathcal{T}_3$ and $\forall \sigma, \tau \in A_3$

$$\mathbb{E} \left[\sup_{f \in C(\mathbb{U}), \|f\|_\infty \leq 1} |\langle x \otimes \tau | f(U_\omega(C)) y \otimes \sigma \rangle| \right] \leq K e^{-\gamma d(x, y)}$$

à la "Aizenman-Molchanov" '09 Hamza-J.-Stolz
Similar approach Asch, Bourget, J. 11', J. '12

⇒ "Exponential" Dynamical Localization:

with X $x \otimes \tau = x x \otimes \tau$, ψ_0 cpct. support, and $p > 0$,

$$\sup_{n \in \mathbb{Z}} \|e^{p|X|} U_\omega^n(C) \psi_0\| < K_\omega, \text{ a.s.}$$

Remarks

- Generalizations to homog. trees of coord. number $q \geq 3$
- RQW "analog" of weak disorder Anderson deloc. on trees '94 Klein
- First RQW: '09 Konno

$$C_{\omega}(\mathbf{x}) = \begin{pmatrix} te^{i\omega_x} & r \\ r & -te^{-i\omega_x} \end{pmatrix} \Rightarrow \text{ballistic behaviour!}$$

- $d = 1$: $\forall 0 < r, t < 1$ s.t. $r^2 + t^2 = 1$ '10 J.-Merkli, '11 Ahlbrecht et al

$$C_{\omega}(\mathbf{x}) = \begin{pmatrix} te^{i\omega_x^{-1}} & re^{i\omega_x^{-1}} \\ re^{i\omega_x+1} & -te^{i\omega_x+1} \end{pmatrix} \Rightarrow \text{dyn. localized}$$

- Related random CMV matrices: 06' Simon
- Large disorder localization: RQW on \mathbb{Z}^d : 12' J.

Thank you and...

Happy Birthday Yosi!