

RELATIVE RIEMANN-ZARISKI SPACES

MICHAEL TEMKIN

ABSTRACT. In this paper we study relative RiemannZariski spaces associated to a morphism of schemes and generalizing the classical RiemannZariski space of a field. We prove that similarly to the classical RZ spaces, the relative ones can be described either as projective limits of schemes in the category of locally ringed spaces or as certain spaces of valuations. We apply these spaces to prove the following two new results: a strong version of stable modification theorem for relative curves; a decomposition theorem which asserts that any separated morphism between quasi-compact and quasi-separated schemes factors as a composition of an affine morphism and a proper morphism. In particular, we obtain a new proof of Nagatas compactification theorem.

1. INTRODUCTION

Let K/k be a finitely generated field extension. In the first half of the 20-th century, Zariski defined a Riemann variety $\mathrm{RZ}_K(k)$ as the projective limit of all projective k -models of K . Zariski showed that this topological space, which is now called a Riemann-Zariski (or Zariski-Riemann) space, possesses the following set-theoretic description: to give a point $\mathbf{x} \in \mathrm{RZ}_K$ is equivalent to give a valuation ring $\mathcal{O}_{\mathbf{x}}$ with fraction field K and such that $k \subset \mathcal{O}_{\mathbf{x}}$. The Riemann-Zariski space possesses a sheaf of rings \mathcal{O} whose stalks are valuation rings of K as above. Zariski made extensive use of these spaces in his desingularization works.

Let S be a scheme and U be a subset closed under generalizations, for example $U = S_{\mathrm{reg}}$ is the regular locus of S , or $U = \eta$ is a maximal point of S . In many birational problems one wants to consider only U -modifications $S' \rightarrow S$, i.e. modifications which do not modify U . Then it is natural to consider the projective limit $\mathfrak{S} = \mathrm{RZ}_U(S)$ of all U -modifications of S . It was remarked in [Tem2, §3.3] that working with such relative Riemann-Zariski spaces one can extend the P -modification results of [Tem2] to the case of general U and S , and this plan is realized in §2. In §2.2 we give a preliminary description of the space \mathfrak{S} , which is used in §2.3 to prove the first main result of the paper, the stable modification theorem 2.3.3 generalizing its analog from [Tem2]. Our improvement to the stable modification theorem [Tem2, 1.5] is in the control on the base change one has to perform in order to construct a stable modification of a relative curve $C \rightarrow S$. Namely, we prove that in order to find a stable modification of a relative curve with semi-stable U -fibers it suffices to replace the base S with a U -étale covering.

I want to express my deep gratitude to B. Conrad for pointing out various gaps and mistakes in an earlier version of the article and to thank R. Huber for a useful discussion. Also I thank D. Rydh and the referee for pointing out some mistakes in §2.3. A first version of the article was written during my stay at the Max Planck Institute for Mathematics at Bonn. The final revision was made when the author was staying at IAS and supported by NFS grant DMS-0635607.

Although a very rough study of relative RZ spaces suffices for the proof of Theorem 2.3.3, it seems natural to investigate these spaces deeper. Furthermore, the definition of relative Riemann-Zariski spaces can be naturally generalized to the case of an arbitrary morphism $f : Y \rightarrow X$, and the case when f is a dominant point was already applied in [Tem1]. So, it is natural to investigate the relative RZ spaces associated to a morphism $f : Y \rightarrow X$. We will see that under a very mild assumption that f is a separated morphism between quasi-compact quasi-separated schemes, one obtains a very specific description of the space $\mathrm{RZ}_Y(X)$ which is similar to the classical case of $\mathrm{RZ}_K(k)$. Let us say that f is *decomposable* if it factors into a composition of an affine morphism $Y \rightarrow Z$ and a proper morphism $Z \rightarrow X$. Actually, in §2.2 we study $\mathrm{RZ}_Y(X)$ in the case of a general decomposable morphism because this case is not essentially easier than the case of an open immersion $Y \hookrightarrow X$. We define a set $\mathrm{Val}_Y(X)$ whose points are certain X -valuations of Y , and construct a surjection $\psi : \mathrm{Val}_Y(X) \rightarrow \mathrm{RZ}_Y(X)$. It will require some additional work to prove in Corollary 3.4.7 that ψ is actually a bijection (and even a homeomorphism with respect to natural topologies defined in the paper). Now, a natural question to ask is if the decomposition assumption is essential. Slightly surprisingly, the answer is negative because the assumption is actually empty. A second main result of this paper is decomposition theorem 1.1.3 which states that a morphism of quasi-compact quasi-separated schemes is decomposable if and only if it is separated. Thus, the description of relative RZ spaces obtained in the decomposable case is actually the general one.

We give two proofs of the decomposition theorem in this paper. The first proof is based on Nagata compactification and Thomason approximation theorems. Actually, we prove in §1.1 that the decomposition theorem is essentially equivalent to the union of these two theorems. This accomplishes the first proof. On the other hand, it turns out that a deeper study of relative RZ spaces leads to an independent proof of the decomposition theorem as explained in §3.5. In particular, we obtain new proofs of Nagata's and Thomason's theorems. Though there are few known proofs of Nagata's theorem, see [Con] and [Lüt], the author expects that the new proof might be better suited for applying to algebraic spaces and (perhaps) certain classes of stacks (joint project with I. Tyomkin).

Let us describe briefly the structure of the paper. In §1.1 we prove a slight generalization of Thomason's theorem and show that the decomposition theorem is essentially equivalent to the union of Nagata's and Thomason's theorems. In §2 we start our study of relative RZ spaces and apply them to the strong stable modification theorem. Then, §3 is devoted to further study of the relative RZ spaces. In §3.1 we establish an interesting connection between Riemann-Zariski spaces and adic spaces of R. Huber; in particular, we obtain an intrinsic topology on $\mathrm{Val}_Y(X)$. However, it turns out that the notion of an open subdomain in the spaces $\mathrm{Val}_Y(X)$ is much finer than its analog in the adic spaces. It requires some work to prove in Theorem 3.3.4 that open subdomains of the form $\mathrm{Val}_{\mathrm{Spec}(B)}(\mathrm{Spec}(A))$ form a basis for the topology of $\mathrm{Val}_Y(X)$. In §3.4 we study Y -blow ups of X , which are analogs of U -admissible or formal blow ups from Raynaud's theory, see [BL]. As a corollary, we prove that $\psi : \mathrm{Val}_Y(X) \rightarrow \mathrm{RZ}_Y(X)$ is a homeomorphism in the decomposable case. Finally, we prove in Theorem 3.5.1 that any open quasi-compact subset of $\mathrm{Val}_Y(X)$ admits a scheme model of the form $\mathrm{Val}_{\bar{Y}}(\bar{X})$ with \bar{Y}

being \overline{X} -affine. This result implies the decomposition theorem, and, therefore, leads to a new proof of Nagata's theorem.

I want to mention that I was motivated by Raynaud's theory in my study of Riemann-Zariski spaces in the decomposable case, and some basic ideas are taken from [BL]. I give a simple illustration of those ideas in the proof of the generalized Thomason's theorem.

When this paper was almost finished I was informed about a recent paper [FK] by Fujiwara and Kato, which contains a survey on a theory of generalized Riemann-Zariski spaces they are developing. The survey announces many interesting results, including Nagata compactification for algebraic spaces. It is clear that there is a certain overlap between that theory and the present paper which can be rather large, though it is difficult to make any conclusion on this subject until the actual proofs are published. The generalized RZ spaces mentioned in [FK] are exactly the relative RZ spaces of open immersions $Y \hookrightarrow X$ (the same case which is used in the proof of the stable modification theorem).

Finally, let us discuss the most recent progress that was made during the last year. Nagata compactification for algebraic spaces was proved independently by Conrad-Lieblich-Olsson in [CLO] (implementing Gabber's approach) and D. Rydh in [Rydh]. In both cases one reduces this to the scheme case rather than proving it from scratch. It should also be noted in this context that important particular cases of the latter theorem (when the algebraic spaces are normal or when the target is a field) were proved much earlier by Raoult, see [R1] and [R2].

1.1. On noetherian approximation and Nagata compactification. For shortness, a filtered projective family of schemes with affine transition morphisms will be called *affine filtered family*. Also, we abbreviate the words "quasi-compact and quasi-separated" by the single "word" qcqs. In [TT, C.9], Thomason proved a very useful approximation theorem, which states that any qcqs scheme Y over a ring Λ is isomorphic to a scheme $\text{proj lim } Y_\alpha$, where $\{Y_\alpha\}_\alpha$ is an affine filtered family of Λ -schemes of finite presentation. Due to the following lemma, this theorem may be reformulated in a more laconic way as follows: Y is affine over a Λ -scheme Y_0 of finite presentation.

Lemma 1.1.1. *A morphism of qcqs schemes $f : Y \rightarrow X$ is affine if and only if $Y \xrightarrow{\sim} \text{proj lim } Y_\alpha$, where $\{Y_\alpha\}_\alpha$ is a filtered family of X -affine finitely presented X -schemes.*

Proof. If $Y \xrightarrow{\sim} \text{proj lim } Y_\alpha$ is as in the lemma then $Y_\alpha = \mathbf{Spec}(\mathcal{E}_\alpha)$ for an \mathcal{O}_X -algebra \mathcal{E}_α , hence $Y = \mathbf{Spec}(\mathcal{E})$ where $\mathcal{E} = \text{inj lim } \mathcal{E}_\alpha$. Conversely, suppose that f is affine. By [EGA I, 6.9.16(iii)], $f_*(\mathcal{O}_Y) \xrightarrow{\sim} \text{inj lim } \mathcal{E}_\alpha$, where $\{\mathcal{E}_\alpha\}$ is a filtered family of finitely presented \mathcal{O}_X -algebras. Hence $Y = \text{proj lim } \mathbf{Spec}(\mathcal{E}_\alpha)$. \square

We generalize Thomason's theorem below. As a by-product, we obtain a simplified proof of the original theorem.

Theorem 1.1.2. *Let $f : Y \rightarrow X$ be a (separated) morphism of qcqs schemes. Then f can be factored into a composition of an affine morphism $Y \rightarrow Z$ and a (separated) morphism $Z \rightarrow X$ of finite presentation.*

Proof. Step 1. *Preliminary work.* First we observe that if f is separated and $Y \rightarrow Z \rightarrow X$ is a factorization as in the theorem, then Y is the projective limit of schemes Y_α which are affine over Z and of finite presentation. By [TT, C.7], already

some Y_α is separated over X , hence replacing Z with Y_α , we achieve a factorization with X -separated Z . This allows us to deal only with the general (not necessarily separated) case in the sequel.

If Y is affine and $f(Y)$ is contained in an open affine subscheme $X' \subset X$ then the claim is obvious. So, Y admits a finite covering by open qcqs subschemes Y_1, \dots, Y_n such that the induced morphisms $Y_i \rightarrow X$ satisfy the conclusion of the theorem. It suffices to prove that one can decrease the natural number n until it becomes 1, and, obviously, it suffices to deal only with the case of $n = 2$. Then the schemes $U := Y_1$ and $V := Y_2$ can be represented as $U = \text{proj lim } U_\beta$ and $V = \text{proj lim } V_\gamma$, where the limits are taken over X -affine filtered families of X -schemes of finite presentation.

Step 2. Affine domination. By [EGA, IV₃, 8.2.11], for $\beta \geq \beta_0$ and $\gamma \geq \gamma_0$, the schemes U_β and V_γ contain open subschemes U'_β and V'_γ , whose preimages in U and V coincide with $W := U \cap V$. By [EGA, IV₃, 8.13.1], the morphism $W \rightarrow U'_{\beta_0}$ factors through V'_γ for sufficiently large γ . Replace γ_0 by γ . By the same reason, the morphism $W \rightarrow V'_{\gamma_0}$ factors through some U'_β and the morphism $W \rightarrow U'_\beta$ factors through some V'_γ . Let us denote the corresponding morphisms as $f_{\gamma,\beta} : V'_\gamma \rightarrow U'_\beta$, f_{β,γ_0} and f_{γ_0,β_0} . Now comes an obvious but critical argument: f_{β,γ_0} is separated because the composition $f_{\gamma_0,\beta_0} \circ f_{\beta,\gamma_0} : U'_\beta \rightarrow U'_{\beta_0}$ is separated (and even affine); $f_{\gamma,\beta}$ is affine because its composition with the separated morphism f_{β,γ_0} is affine. We gather the already defined objects in the left diagram below. Note that everything is defined over X , the horizontal arrows are open immersions, the vertical arrows are affine morphisms and the indexed schemes are of finite X -presentation.

$$\begin{array}{ccc}
 V & \longleftarrow W \hookrightarrow & U \\
 \downarrow & & \downarrow \phi' \\
 V_\gamma & \longleftarrow V'_\gamma \hookrightarrow & U \\
 & & \downarrow h \\
 & & U'_\beta \hookrightarrow U_\beta
 \end{array}
 \qquad
 \begin{array}{ccc}
 V & \longleftarrow W \hookrightarrow & U \\
 \downarrow & & \downarrow \phi' \\
 V_\gamma & \longleftarrow V'_\gamma \hookrightarrow & U_\gamma \\
 & & \downarrow f_{\gamma,\beta} \\
 & & U'_\beta \hookrightarrow U_\beta
 \end{array}$$

Step 3. Affine extension. The main task of this step is to produce the right diagram from the left one. It follows from the previous stage that $V'_\gamma = \mathbf{Spec}(\mathcal{E}')$, where \mathcal{E}' is a finitely presented $\mathcal{O}_{U'_\beta}$ -algebra. The morphism $\phi' : W \rightarrow V'_\gamma$ to a U'_β -affine scheme corresponds to a homomorphism $\phi' : \mathcal{E}' \rightarrow h'_*(\mathcal{O}_W)$, where $h' : W \rightarrow U'_\beta$ is the projection. Obviously $h_*(\mathcal{O}_U)|_{U'_\beta} \xrightarrow{\sim} h'_*(\mathcal{O}_W)$, where $h : U \rightarrow U_\beta$ is the projection. Hence we can apply [EGA I, 6.9.10.1], to find a finitely presented \mathcal{O}_{U_β} -algebra \mathcal{E} and a homomorphism $\varphi : \mathcal{E} \rightarrow h_*(\mathcal{O}_U)$ such that $\mathcal{E}|_{U'_\beta} \xrightarrow{\sim} \mathcal{E}'$ and the restriction of φ to U'_β is ϕ' . Set $U_\gamma = \mathbf{Spec}(\mathcal{E})$, then $U_\gamma \rightarrow U_\beta$ is an affine morphism whose restriction over U'_β is $f_{\gamma,\beta}$, and φ induces a morphism $\phi : U \rightarrow U_\gamma$. Finally, we glue U_γ and V_γ along V'_γ obtaining a finitely presented X -scheme Z , and notice that the affine morphisms $U \rightarrow U_\gamma$ and $V \rightarrow V_\gamma$ glue to an affine morphism $Y \rightarrow Z$ over X . \square

Our proof is a simple analog of Raynaud's theory. Thomason used the first two steps (induction argument in the proof of Theorem C.9 and Lemma C.6). Our simplification of his proof is due to the third step. The same arguments are used in Raynaud's theory, see the end of the proof of [BL, 4.1(d)] and [BL, 2.6(a)]. In our

paper, they also appear in the proofs of Lemmas 3.4.2(i) and 3.4.4, and Theorem 3.5.1.

Next, we recall Nagata compactification theorem, see [Nag]. A scheme theoretic proof of the theorem can be found in [Con] or [Lüt]. Recall that a morphism $f : Y \rightarrow X$ is called *compactifiable* if it can be factored as a composition of an open immersion $g : Y \rightarrow Z$ and a proper morphism $h : Z \rightarrow X$. Nagata proved that a finite type morphism $f : Y \rightarrow X$ of qcqs schemes is compactifiable if and only if it is separated. Actually, Nagata considered noetherian schemes, and the general case was proved by B. Conrad in [Con].

Assume that f is factored as above. Let $\mathcal{I} \subset \mathcal{O}_Z$ be an ideal with support $Z \setminus Y$ and let Z' be the blow up of Z along \mathcal{I} . We can choose a finitely generated \mathcal{I} because the morphism $Y \hookrightarrow Z$ is quasi-compact. The open immersion $g' : Y \rightarrow Z'$ is affine because $Z' \setminus Y$ is a locally principal divisor. It follows that g is a composition of an affine morphism g' of finite type and a proper morphism $Z' \rightarrow X$. Conversely, assume that $g : Y \rightarrow Z$ is affine of finite type and $Z \rightarrow X$ is proper. Then Y is quasi-projective over Z , hence there exists an open immersion of finite type $Y \hookrightarrow \bar{Y}$ with Z -projective and, therefore, X -proper \bar{Y} . Thus, Nagata's theorem can be reformulated as follows: a finite type morphism is separated if and only if it can be represented as a composition of an affine morphism of finite type and a proper morphism. Now, one sees that a weak form of Theorem 1.1.2 (f is separated and $Z \rightarrow X$ is of finite type) and Nagata's theorem are together equivalent to the following decomposition theorem, which will be also proved in §3.5 by a different method.

Theorem 1.1.3. *A morphism $f : Y \rightarrow X$ of quasi-compact quasi-separated schemes is separated if and only if it can be factored as a composition of an affine morphism $Y \rightarrow Z$ and a proper morphism $Z \rightarrow X$.*

2. PRELIMINARY DESCRIPTION OF RELATIVE RZ SPACES AND APPLICATIONS

Throughout §2, $f : Y \rightarrow X$ denotes a separated morphism between qcqs schemes.

2.1. Valuations and projective limits. We are going to recall some notions introduced in [Tem2, §3.3]. Consider a factorization of f into a composition of a schematically dominant morphism $f_i : Y \rightarrow X_i$ and a proper morphism $g_i : X_i \rightarrow X$. We call the pair (f_i, g_i) a *Y-modification* of X , and usually it will be denoted simply as X_i . Given two Y -modifications of X , we say that X_j *dominates* or *refines* X_i , if there exists an X -morphism $g_{ji} : X_j \rightarrow X_i$ compatible with f_i, f_j, g_i and g_j . A standard graph argument shows that if g_{ji} exists then it is unique (one uses only that f_j is schematically dominant and X_i is X -separated). The family $\{X_i\}_{i \in I}$ of all Y -modifications of X is filtered because any two Y -modifications X_i, X_j are dominated by the scheme-theoretic image of Y in $X_i \times_X X_j$, and it has an initial object corresponding to the schematic image of Y in X .

A relative Riemann-Zariski space $\mathfrak{X} = \text{RZ}_Y(X)$ is defined as the projective limit of the underlying topological spaces of Y -modifications of X . Note that if X is integral and Y is its generic point then one recovers the classical Riemann-Zariski spaces. A slightly more general case, when Y is a dominant point, was considered in [Tem1, §1]. Let $\pi_i : \mathfrak{X} \rightarrow X_i$ be the projections and $\eta : Y \rightarrow \mathfrak{X}$ be the map induced by f_i 's. We provide \mathfrak{X} with the sheaf $\mathcal{M}_{\mathfrak{X}} = \eta_*(\mathcal{O}_Y)$, which will be called the sheaf of *meromorphic functions*, and with the sheaf $\mathcal{O}_{\mathfrak{X}} = \text{inj lim } \pi_i^{-1}(\mathcal{O}_{X_i})$,

which will be called the sheaf of *regular functions*. The natural homomorphisms $\alpha_i : \pi_i^{-1}(\mathcal{O}_{X_i}) \rightarrow \mathcal{M}_{\mathfrak{X}}$ induce a homomorphism $\alpha : \mathcal{O}_{\mathfrak{X}} \rightarrow \mathcal{M}_{\mathfrak{X}}$, and we will prove later that η is injective and α is a monomorphism. Actually, we will give in Corollary 3.5.2 a rather precise meaning to a claim that $\mathcal{M}_{\mathfrak{X}}$ is a sheaf of semi-fractions of the sheaf $\mathcal{O}_{\mathfrak{X}}$.

Remark 2.1.1. For any filtered projective family of locally ringed spaces $\{Y_j\}_{j \in J}$ the projective limit $\mathfrak{Y} = \text{proj lim}_{j \in J} Y_j$ always exists and satisfies $|\mathfrak{Y}| := \text{proj lim } |Y_j|$ and $\mathcal{O}_{\mathfrak{Y}} = \text{inj lim } \pi_j^{-1} \mathcal{O}_{Y_j}$ where $\pi_j : \mathfrak{Y} \rightarrow Y_j$'s are the projections. Assume now that Y_j 's are schemes. Then \mathfrak{Y} is known to be a scheme when the transition morphisms are affine: this situation is studied very extensively in [EGA, IV₃, §8] and the obtained results have a plenty of various very important applications. Although \mathfrak{Y} does not have to be a scheme in general, it is a locally ringed space of a rather special form which deserves a study. Our relative RZ spaces $(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}})$ provide a nice example of such pro-schemes (while $\mathcal{M}_{\mathfrak{X}}$ corresponds to an extra-structure related to Y), and we will later obtain a very detailed description of these spaces (e.g. we will describe the stalks of $\mathcal{O}_{\mathfrak{X}}$). Another interesting example of a pro-scheme which is not a scheme but has a very nice realization is as follows: let X be a scheme with a subset U closed with respect to generalization, then $(U, \mathcal{O}_X|_U)$ is the projective limit of all open neighborhoods of U . Note that this locally ringed space does not have to be a scheme: for example, take U to be the set of all non-closed points on an algebraic surface X .

The classical absolute RZ spaces viewed either as topological spaces or, more generally, as locally ringed spaces admit two alternative descriptions: (a) a projective limit of schemes, (b) a space whose points are valuations. We defined the relative spaces $\text{RZ}_Y(X)$ using projective limits, but they also admit a "valuative" description as spaces $\text{Val}_Y(X)$. In §2 we only introduce the sets $\text{Val}_Y(X)$ and establish a certain connection between $\text{RZ}_Y(X)$ and $\text{Val}_Y(X)$ which suffices for application to the stable modification theorem 2.3.3. Throughout this paper by a *valuation* on a ring B we mean a commutative ordered group Γ with a multiplicative map $|\cdot| : B \rightarrow \Gamma \cup \{0\}$ which satisfies the strong triangle inequality and sends 1 to 1. Recall that if B is a field then $R = \{x \in B \mid |x| \leq 1\}$ is a valuation ring of B (i.e. $\text{Frac}(R) = B$) which defines $|\cdot|$ up to an equivalence. In general, a valuation is defined up to an equivalence by its kernel p , which is a prime ideal, and by the induced valuation on the residue field $\text{Frac}(B/p)$. By slight abuse of language, the point of $\text{Spec}(B)$ given by p will be also called the *kernel* of $|\cdot|$. Also, we will often identify equivalent valuations.

Remark 2.1.2. We follow R. Huber by using the notion of a valuation. Since these valuations may have a non-empty kernel, a reasonable alternative, however, would be the notion of a semivaluation. Note also that in the literature on abstract algebra this object is often called Manis valuation.

Now, let $\text{Val}_Y(X)$ be the set of triples $\mathbf{y} = (y, R, \phi)$, where $y \in Y$ is a point, R is a valuation ring of $k(y)$ (in particular $\text{Frac}(R) = k(y)$) and $\phi : S = \text{Spec}(R) \rightarrow X$ is a morphism compatible with $y = \text{Spec}(k(y)) \rightarrow Y$ and such that the induced morphism $y \rightarrow S \times_X Y$ is a closed immersion. Let $\mathcal{O}_{\mathbf{y}}$ denote the preimage of R in $\mathcal{O}_{Y,y}$ (currently, it is just a ring attached to \mathbf{y}). We would like to axiomatize the properties of $\mathcal{O}_{\mathbf{y}}$ as follows. By a *semi-valuation ring* we mean a ring \mathcal{O} with a valuation $|\cdot|$ such that any zero divisor of \mathcal{O} lies in the kernel $m = \text{Ker}(|\cdot|)$ and

for any pair $g, h \in \mathcal{O}$ with $|g| \leq |h| \neq 0$ one has that $h|g$. Two structures of a semi-valuation ring on \mathcal{O} are *equivalent* if their valuations are equivalent.

Note that \mathcal{O} embeds into $A = \mathcal{O}_m$ by our assumption on zero divisors, $mA = m$ because the prime ideal m is $(\mathcal{O} \setminus m)$ -divisible, and $R = \mathcal{O}/m$ is the valuation ring of A/m corresponding to the valuation induced by $|\cdot|$. Therefore, \mathcal{O} is *composed* from the local ring A and the valuation ring $R \subset A/m$ in the sense that \mathcal{O} is the preimage of R in A . We say that A is a *semi-fraction ring* of \mathcal{O} . Conversely, any ring composed from a local ring and a valuation ring is easily seen to be a semi-valuation ring. Semi-valuation rings play the same role in the theory of relative RZ spaces as valuation rings do in the theory of usual RZ spaces.

Remark 2.1.3. (i) The structure of a semi-valuation ring on an abstract local ring \mathcal{O} is uniquely defined (up to an equivalence) by its kernel m because \mathcal{O}/m is a valuation ring and hence defines the valuation. Since $A = \mathcal{O}_m$ we obtain that the semi-valuation ring structure on \mathcal{O} is uniquely defined by its embedding into the semi-fraction ring A .

(ii) An abstract ring \mathcal{O} can admit many semi-valuation ring structures. For example, if \mathcal{O} is a valuation ring then any its localization (i.e. a larger valuation ring in its field of fractions) can serve as its semi-fraction ring.

Here is a generalization of the classical criterion that an integral domain \mathcal{O} is a valuation ring if and only if for any pair of elements $f, g \in \mathcal{O}$ either $f|g$ or $g|f$.

Lemma 2.1.4. *Let $\mathcal{O} \subset A$ be two rings. Then the following conditions are equivalent:*

(i) \mathcal{O} admits a structure of a semi-valuation ring such that A is \mathcal{O} -isomorphic to the semi-fraction ring of \mathcal{O} ,

(ii) if $f, g \in A$ are co-prime (i.e. $fA + gA = A$) then either $f \in g\mathcal{O}$ or $g \in f\mathcal{O}$.

Proof. We should only prove that (ii) implies (i), since the opposite implication is obvious. We claim that A is a local ring. Indeed, if it is not local then $A \setminus A^\times$ is not an ideal, hence there exist non-invertible f, g with invertible $f + g$. But by our assumption either $f \in gA$ or $g \in fA$, hence $f + g$ is contained in a proper ideal equal to either fA or gA , that is an absurd. Let $m \subset A$ be the maximal ideal, then taking $f \in m$ and $g = 1$ and observing that f does not divide 1 in \mathcal{O} (and even in A), we deduce that $f \in \mathcal{O}$. Thus, we proved that $m \subset \mathcal{O}$, in particular, \mathcal{O} is the preimage of the ring $\mathcal{O}/m \subset A/m$ under the surjection $A \rightarrow A/m$. It remains to show that \mathcal{O}/m is a valuation ring of A/m . For a pair of elements $\tilde{f}, \tilde{g} \in \mathcal{O}/m$ choose liftings $f, g \in \mathcal{O}$. Since either $f|g$ or $g|f$ in \mathcal{O} , it follows that either $\tilde{f}|\tilde{g}$ or $\tilde{g}|\tilde{f}$. Hence \mathcal{O}/m is a valuation ring, and we are done. \square

2.2. RZ space of a decomposable morphism. Let $\mathbf{y} = (y, R, \phi)$ be a point of $\text{Val}_Y(X)$ and let $S = \text{Spec}(R)$. By the valuative criterion of properness, ϕ factors uniquely through a morphism $\phi_i : Y \rightarrow X_i$ for any Y -modification $X_i \rightarrow X$. Since $S \times_{X_i} Y$ is a closed subscheme of $S \times_X Y$ by X -separatedness of X_i , we obtain that ϕ_i induces a closed immersion $y \rightarrow S \times_{X_i} Y$, and, in particular, (y, R, ϕ_i) is an element of $\text{Val}_Y(X_i)$. It follows that the natural map $\text{Val}_Y(X_i) \rightarrow \text{Val}_Y(X)$ is a bijection. So, $\text{RZ}_Y(X)$ and $\text{Val}_Y(X)$ depend on X and Y only up to replacing X with its Y -modification.

Now we will construct a map of sets $\psi : \text{Val}_Y(X) \rightarrow \text{RZ}_Y(X)$. For any $i \in I$, let $x_i \in X_i$ be the *center* of R on X_i , i.e. the image of the closed point of S under

ϕ_i . Then the family of points (x_i) defines a point $\mathbf{x} \in \mathfrak{X}$ and we obtain a map ψ as above. For any i , x_i is a specialization of $f_i(y)$, hence we obtain a homomorphism $\mathcal{O}_{X_i, x_i} \rightarrow \mathcal{O}_{X_i, f_i(y)} \rightarrow \mathcal{O}_{Y, y} \rightarrow k(y)$ whose image lies in R because x_i is the center of R on X_i . Therefore, the image of \mathcal{O}_{X_i, x_i} in $\mathcal{O}_{Y, y}$ lies in $\mathcal{O}_{\mathbf{y}}$, and we obtain a natural homomorphism $\mathcal{O}_{\mathfrak{X}, \mathbf{x}} = \text{inj lim } \mathcal{O}_{X_i, x_i} \rightarrow \mathcal{O}_{\mathbf{y}}$.

Proposition 2.2.1. *Suppose that f is decomposable. Then any point $\mathbf{x} \in \mathfrak{X}$ possesses a preimage $\mathbf{y} = \lambda(\mathbf{x})$ in $\text{Val}_Y(X)$ such that the homomorphism $\mathcal{O}_{\mathfrak{X}, \mathbf{x}} \rightarrow \mathcal{O}_{\mathbf{y}}$ is an isomorphism. In particular, λ is a section of ψ .*

Actually, we will prove in §3 that ψ is a bijection (so λ is its inverse), but the proposition as it is already covers our applications in §2.

Proof. Factor f into a composition of an affine morphism $Y \rightarrow Z$ and a proper morphism $Z \rightarrow X$. After replacing X with the scheme-theoretic image of Y in Z , we can assume that f is affine. Note that then for any Y -modification $X_i \rightarrow X$, the morphism $f_i : Y \rightarrow X_i$ is affine. Let x_i be the image of \mathbf{x} in X_i . Obviously, the schemes $U_i = \text{Spec}(\mathcal{O}_{X_i, x_i}) \times_{X_i} Y$ are affine. In addition, on the level of sets each U_i consists of points $y \in Y$ such that x_i is a specialization of $f_i(y)$, the morphisms $U_i \rightarrow Y$ are topological embeddings and $\mathcal{O}_Y|_{U_i} \xrightarrow{\sim} \mathcal{O}_{U_i}$. Notice that the schemes $U_i = \text{Spec}(B_i)$ form a filtered family, hence $U_\infty := \text{proj lim } U_i = \text{Spec}(B_\infty)$, where $B_\infty = \text{inj lim } B_i$. By [EGA, IV₃, §8], $U_\infty = \bigcap U_i$ set-theoretically. Since $f_i : Y \rightarrow X_i$ is schematically dominant and the latter property is preserved under (possibly infinite) localizations on the base, the morphism $U_i \rightarrow \text{Spec}(\mathcal{O}_{X_i, x_i})$ is schematically dominant too. So, for each $i \in I$ we have that $\mathcal{O}_{X_i, x_i} \hookrightarrow B_i$, and then an embedding of the direct limits $\mathcal{O}_{\mathfrak{X}, \mathbf{x}} \hookrightarrow B_\infty$ arises.

Lemma 2.2.2. *Suppose that elements $g, h \in B_\infty$ do not have common zeros on U_∞ . Then either $g \in h\mathcal{O}_{\mathfrak{X}, \mathbf{x}}$ or $h \in g\mathcal{O}_{\mathfrak{X}, \mathbf{x}}$.*

Proof. Find i such that g and h are defined and do not have common zeros on U_i . Note that $U_i = \bigcap f^{-1}(V_j)$, where V_j runs over affine neighborhoods of x_i . Hence we can choose a neighborhood $X'_i = \text{Spec}(A)$ of x_i such that $g, h \in B$ and $gB + hB = 1$, where $Y' = \text{Spec}(B)$ is the preimage of X'_i in Y . To ease the notation we will write X and x instead of X_i and x_i (we can freely replace X with X_i because $\text{RZ}_Y(X)$ remains unchanged). Now, the pair (g, h) induces a morphism $\alpha' : Y' \rightarrow P' := \text{Proj}(A[T_g, T_h])$, whose scheme-theoretic image \overline{X}' is a Y' -modification of X' . It would suffice to extend the Y' -modification $\alpha' : \overline{X}' \rightarrow X'$ to a Y -modification $\alpha : \overline{X} \rightarrow X$. Indeed, either $T_g \in T_h\mathcal{O}_{\overline{X}', x'}$ or $T_h \in T_g\mathcal{O}_{\overline{X}', x'}$, where $x' \in \overline{X}'$ is the image of \mathbf{x} in \overline{X}' . So, existence of α would imply that $g|h$ or $h|g$ already in the image of $\mathcal{O}_{\overline{X}, x'}$ in B_∞ , which is by definition contained in $\mathcal{O}_{\mathfrak{X}, \mathbf{x}}$.

It can be difficult to extend α' (without applying Nagata compactification), but fortunately we can replace \overline{X}' with any its Y' -modification \overline{X}'' and it suffices to extend $\overline{X}'' \rightarrow X'$ to a Y -modification of X . Choose a, b such that $ag + bh = 1$. Then there exists a natural morphism $\beta' : Y' \rightarrow P'' := \text{Proj}(A[T_{ag}, T_{ah}, T_{bg}, T_{bh}])$ which takes Y' to the affine chart on which $T_{ag} + T_{bh}$ is invertible. We define \overline{X}'' to be the scheme-theoretic image of β' . Since β' factors through Segre embedding $\text{Proj}(A[T_g, T_h]) \times \text{Proj}(A[T_a, T_b]) \hookrightarrow P''$, we obtain that \overline{X}'' is a closed subscheme of the source which is mapped to \overline{X}' by the projection onto the first factor. In

particular, \overline{X}'' is a Y' -modification of \overline{X}' . We will show that $\overline{X}'' \rightarrow X'$ extends to a Y -modification $\overline{X} \rightarrow X$.

Let $E \subset B$ be the A -submodule generated by ag, ah, bg, bh and consider the graded algebra $A_E := \bigoplus_{n=0}^{\infty} E^n$, where E^n is the n -th power of E in B and E^0 is the image of A . Note that $1 \in E$, and we will denote by 1_E the associated 1-graded element of A_E . Set $P := \text{Proj}(A_E)$ and observe that the affine chart corresponding to 1_E is $P_1 = \text{Spec}(\bigcup_{n=0}^{\infty} E^n)$ (where the union is taken inside of B). Clearly, P is a closed subscheme in P'' and the morphism $Y \rightarrow P''$ factors through P_1 . In particular, \overline{X}'' is the schematical image of $Y \rightarrow P$, and the latter coincides with the schematical closure of P_1 because the morphism $Y \rightarrow P_1$ is schematically dominant by injectivity of the homomorphism $\bigcup_{n=0}^{\infty} E^n \rightarrow B$. By [EGA I, 6.9.7], E can be extended to a finitely generated \mathcal{O}_X -submodule $\mathcal{E} \subset f_*(\mathcal{O}_Z)$, and replacing \mathcal{E} by $\mathcal{E} + \mathcal{O}_X$ we achieve in addition that \mathcal{E} contains the image of \mathcal{O}_X in $f_*(\mathcal{O}_Y)$. Let \mathcal{E}^n be the n -th power of \mathcal{E} in the sheaf of \mathcal{O}_X -algebras $f_*(\mathcal{O}_Y)$ (so, \mathcal{E}^0 is the image of \mathcal{O}_X) and form the graded \mathcal{O}_X -algebra $\mathcal{A}_{\mathcal{E}} := \bigoplus_{n=0}^{\infty} \mathcal{E}^n$. Then exactly the same computation as was used above shows that the schematical closure of $\text{Spec}(\bigcup_{n=0}^{\infty} \mathcal{E}^n)$ in $\text{Proj}(\mathcal{A}_{\mathcal{E}})$ is a Y -modification of X , which we denote \overline{X} . Since $\overline{X} \rightarrow X$ obviously extends $\overline{X}'' \rightarrow X'$, we are done. \square

The above lemma combined with Lemma 2.1.4 provides $\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}$ with a semi-valuation ring structure such that B_{∞} is its semi-fraction ring. In particular, B_{∞} is a local ring and so U_{∞} possesses a unique closed point y . Thus, $B_{\infty} = \mathcal{O}_{Y, y}$, its subring $\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}$ contains m_y and $R := \mathcal{O}_{\mathfrak{x}, \mathfrak{x}}/m_y$ is a valuation ring of $k(y)$. Define $\phi : S = \text{Spec}(R) \rightarrow X$ as the composition of the closed immersion $S \rightarrow \text{Spec}(\mathcal{O}_{\mathfrak{x}, \mathfrak{x}})$ with the natural morphism $\text{Spec}(\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}) \rightarrow X$. Since $\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}$ is composed from $\mathcal{O}_{Y, y}$ and R , the triple $\mathbf{y} := (y, R, \phi)$ is a candidate for being $\lambda(\mathbf{x})$ and it only remains to check that $y \rightarrow S \times_X Y$ is a closed immersion (and so \mathbf{y} is indeed an element of $\text{Val}_Y(X)$).

For any i , $U_i = \text{Spec}(\mathcal{O}_{X_i, x_i}) \times_{X_i} Y$ is a closed subscheme of $\text{Spec}(\mathcal{O}_{X_i, x_i}) \times_X Y$, hence $U_{\infty} \xrightarrow{\sim} \text{proj lim}_{i \in I} U_i$ is a closed subscheme of

$$\text{Spec}(\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}) \times_X Y \xrightarrow{\sim} \text{proj lim}_{i \in I} \text{Spec}(\mathcal{O}_{X_i, x_i}) \times_X Y$$

Since y is closed in U_{∞} , we obtain that the morphism $y \rightarrow \text{Spec}(\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}) \times_X Y$ is a closed immersion. Hence the morphism from y to a closed subscheme $S \times_X Y$ of $\text{Spec}(\mathcal{O}_{\mathfrak{x}, \mathfrak{x}}) \times_X Y$ is a closed immersion too and we are done. \square

2.3. Applications. A preliminary description of relative Riemann-Zariski spaces obtained in the previous section, suffices for some applications. Assume we are given a qcqs scheme S with a schematically dense quasi-compact subset U (i.e. any neighborhood of U is schematically dense) which is closed under generalizations. An S -scheme X is called *U -admissible* if the preimage of U in X is schematically dense. By a *U -étale covering* we mean a separated finite type morphism $\phi : S' \rightarrow S$ such that ϕ is étale over U , S' is U -admissible, and for any valuation ring R any morphism $\text{Spec}(R) \rightarrow S$ taking the generic point to U lifts to a morphism $\text{Spec}(R') \rightarrow S'$ where R' is a valuation ring dominating R and such that $\text{Frac}(R')/\text{Frac}(R)$ is finite. (Actually those are finite type h -covers of S which are étale over U .) Note that in [BLR] one considers a more restrictive class of coverings, namely U -étale maps $S' \rightarrow S$, which split to a composition of a surjective flat U -étale morphism and

a U -modification. However, it follows from the flattening theorem [RG, 5.2.2] of Raynaud-Gruson that the latter class of coverings is cofinal in ours.

In order to make use of Riemann-Zariski spaces we have first to establish some properties of schemes over semi-valuation rings. So, let \mathcal{O} be a semi-valuation ring with semi-fraction ring A and let m be the maximal ideal of A . Recall that $A = \mathcal{O}_m$, $R := \mathcal{O}/m$ is the valuation ring in $K := A/m$, the scheme $S = \text{Spec}(\mathcal{O})$ is covered by pro-open subscheme $U = \text{Spec}(A)$ (i.e. U is the intersection of open subschemes) and closed subscheme $T = \text{Spec}(R)$, and the intersection $U \cap T$ is a single point $\eta = \text{Spec}(K)$, which is the generic point of T and the closed point of U . Note that in some sense S is glued from U and T along η , for example, there is a bi-Cartesian square

$$\begin{array}{ccc} \eta & \longrightarrow & U \\ \downarrow & & \downarrow \\ T & \longrightarrow & S \end{array}$$

Next we will study how U -admissible S -schemes (resp. quasi-coherent \mathcal{O}_S -modules) can be glued from T -schemes and U -schemes (resp. modules), and we will call such gluing (U, T) -descent. Given a quasi-coherent \mathcal{O}_S -module M , which we identify with an \mathcal{O} -module, set $M_U = M \otimes_{\mathcal{O}} A$, $M_T = M \otimes_{\mathcal{O}} R = M/mM$ and $M_\eta = M \otimes_{\mathcal{O}} K$. We say that M is U -admissible if the localization homomorphism $M \rightarrow M_U$ is injective. Note that any \mathcal{O} -module M defines a descent datum consisting of M_U, M_T and an isomorphism $\phi_M : M_U \otimes_A K \xrightarrow{\sim} M_T \otimes_R K$, and a similar claim holds for S -schemes. The corresponding categories of descent data are defined in an obvious way, and, naturally, we have a (U, T) -descent lemma below. Slightly more generally, we fix a qcqs U -admissible S -scheme \bar{S} with $\bar{U} = U \times_S \bar{S}$, $\bar{T} = T \times_S \bar{S}$ and $\bar{\eta} = \eta \times_S \bar{S}$ and we will glue objects defined over \bar{U} and \bar{T} along their restrictions over $\bar{\eta}$. For example, an $\mathcal{O}_{\bar{S}}$ -module \mathcal{M} induces a descent data $\phi_{\mathcal{M}} : \mathcal{M}_{\bar{U}}|_{\bar{\eta}} \xrightarrow{\sim} \mathcal{M}_{\bar{T}}|_{\bar{\eta}}$. If we want to stress the choice of \bar{S} we will call such gluing (\bar{U}, \bar{T}) -descent. For an \bar{S} -scheme X we will use the notation $X_U = X \times_{\bar{S}} \bar{U}$ (and so $X_U \xrightarrow{\sim} X \times_S U$), $X_T = X \times_{\bar{S}} \bar{T}$ and $X_\eta = X \times_{\bar{S}} \bar{\eta}$.

Lemma 2.3.1. *Keep the above notation.*

(i) *The natural functor from the category of U -admissible quasi-coherent $\mathcal{O}_{\bar{S}}$ -modules (resp. $\mathcal{O}_{\bar{S}}$ -algebras) \mathcal{M} to the category of descent data $(\mathcal{M}_{\bar{U}}, \mathcal{M}_{\bar{T}}, \phi_{\mathcal{M}})$ with quasi-coherent $\mathcal{O}_{\bar{U}}$ -module (resp. $\mathcal{O}_{\bar{U}}$ -algebra) $\mathcal{M}_{\bar{U}}$ and quasi-coherent $\bar{\eta}$ -admissible $\mathcal{O}_{\bar{T}}$ -module (resp. $\mathcal{O}_{\bar{T}}$ -algebra) $\mathcal{M}_{\bar{T}}$ is an equivalence of categories.*

(ii) *The (\bar{U}, \bar{T}) -descent is effective on \bar{U} -flat \bar{S} -projective schemes with fixed relatively ample sheaves. More concretely, assume that we are given a descent datum $((X_U, \mathcal{L}_U), (X_T, \mathcal{L}_T), (\phi_X, \phi_{\mathcal{L}}))$, where $f_U : X_U \rightarrow \bar{U}$ and $f_T : X_T \rightarrow \bar{T}$ are projective morphisms with relatively ample invertible modules \mathcal{L}_U and \mathcal{L}_T , respectively, f_U is flat, X_T is η -admissible, $\phi_X : X_U \times_U \eta \xrightarrow{\sim} X_T \times_T \eta$ and $\phi_{\mathcal{L}}$ is an isomorphism between the restrictions of \mathcal{L}_U and \mathcal{L}_T on the η -fibers which agrees with ϕ_X . Then there exists a projective morphism $f : X \rightarrow \bar{S}$ with a relatively ample \mathcal{O}_X -module \mathcal{L} whose restriction over \bar{U} and \bar{T} give rise to the above descent datum.*

(iii) *A qcqs U -admissible S -scheme X is of finite type if and only if X_U and X_T are so. If in addition $X \times_S U \rightarrow U$ is flat and finitely presented then $X \rightarrow S$ is flat and finitely presented.*

Proof. The claim of (i) is local on \bar{S} , so we can assume that \bar{S} is affine. Then $\mathcal{O}_{\bar{S}}$, $\mathcal{O}_{\bar{U}}$ or $\mathcal{O}_{\bar{T}}$ -modules can be viewed simply as \mathcal{O} , A or R -modules, and this reduces our problem to the case when $\bar{S} = S$. In particular, we will now denote the modules as M , M_T , etc. Next we note that $mM_U = mM$ because $mA = m$, and hence $M_T = M/mM$ embeds into $M_\eta = M_U/mM_U$. So, M_T is η -admissible and the embedding $M \hookrightarrow M_U$ identifies M with the preimage of M_T under the projection $M_U \rightarrow M_\eta$. In particular, an exact sequence $0 \rightarrow M \rightarrow M_U \oplus M_T \rightarrow M_\eta \rightarrow 0$ arises. Conversely, given a descent datum as in (i), we can define an \mathcal{O} -module $M = \text{Ker}(M_U \oplus M_T \rightarrow M_\eta)$, and one easily sees that M is actually the preimage of $M_T \subset M_\eta$ under the projection $M_U \rightarrow M_U/mM_U \xrightarrow{\sim} M_\eta$ and hence M_U and M_T are the base changes of this M . We constructed maps from \mathcal{O}_S -modules to descent data and vice versa, and one immediately sees that these maps extend to functors. Then it is obvious from the above that these functors are equivalences of categories which are inverse one to another.

To prove (ii) we find sufficiently large n so that the n -th tensor powers of the initial sheaves induce closed immersions $X_U \rightarrow \mathbf{P}((f_U)_*(\mathcal{L}_U^{\otimes n}))$ and $X_T \rightarrow \mathbf{P}((f_T)_*(\mathcal{L}_T^{\otimes n}))$ into the associated projective fibers. Moreover, the higher direct images of $\mathcal{L}_U^{\otimes n}$ vanish for large n and then $(f_\eta)_*(\mathcal{L}_\eta^{\otimes n}) \xrightarrow{\sim} ((f_U)_*(\mathcal{L}_U^{\otimes n}))_\eta$ by the theorem on base changes and direct images, see [Har, III.12.9]. By part (i) the sheaves $(f_U)_*(\mathcal{L}_U^{\otimes n})$ and $(f_T)_*(\mathcal{L}_T^{\otimes n})$ glue along $(f_\eta)_*(\mathcal{L}_\eta^{\otimes n})$ to an $\mathcal{O}_{\bar{S}}$ -module \mathcal{M} and so $\mathbf{P} := \mathbf{P}(\mathcal{M})$ is glued from $\mathbf{P}_U := \mathbf{P}((f_U)_*(\mathcal{L}_U^{\otimes n}))$ and $\mathbf{P}_T := \mathbf{P}((f_T)_*(\mathcal{L}_T^{\otimes n}))$ along $\mathbf{P}((f_\eta)_*(\mathcal{L}_\eta^{\otimes n}))$. In particular, the closed subschemes $X_U \hookrightarrow \mathbf{P}_U$ and $X_T \hookrightarrow \mathbf{P}_T$ glue to a closed subscheme $i : X \hookrightarrow \mathbf{P}$ with a relatively very ample sheaf $\mathcal{K} := i^*(\mathcal{O}_{\mathbf{P}}(1))$. Note that \mathcal{K} is glued from $\mathcal{L}_U^{\otimes n}$ and $\mathcal{L}_T^{\otimes n}$. Finally, the modules \mathcal{L}_U and \mathcal{L}_T glue to an invertible \mathcal{O}_X -sheaf \mathcal{L} with $\mathcal{L}^{\otimes n} \xrightarrow{\sim} \mathcal{K}$ by (X_U, X_T) -descent of modules, which was established in (i). In particular, \mathcal{L} is relatively ample.

The first assertion of (iii) is exactly Step 2 from the proof of [Tem2, 2.5.3]. So, let us assume that $X \times_{\bar{S}} \bar{U} \rightarrow \bar{U}$ is flat and finitely presented (in addition to the assumption that X is of finite type over \bar{S} and \bar{U} -admissible). The claim is local on X so we can assume that $X = \text{Spec}(C)$ is affine. Note also that X_T is η -admissible, so C/mC embeds into $(C/mC) \otimes_R K$ and then C/mC is flat and finitely presented over R by [Tem2, 3.5.1]. We first deal with finite presentation, so fix an epimorphism $\phi : \mathcal{O}[T] \rightarrow C$ with $T = (T_1, \dots, T_k)$ and let us prove that its kernel I is finitely generated. Localizing at m we obtain an epimorphism $\phi \otimes_{\mathcal{O}} A : A[T] \rightarrow B = C_m$ with kernel $J = I_m$. Then B is A -flat by our assumption and we claim that this implies that $J \cap m[T] = mJ$. Indeed, if x is contained in $J \cap m[T]$ but not in mJ then it reduces to a non-zero element \tilde{x} in the kernel of $J/mJ \rightarrow A[T]/m[T]$. However, this kernel is an epimorphic image of $\text{Tor}_1^A(B, m) = 0$ and hence $\tilde{x} = 0$. By finite presentation of $A \rightarrow B$ we have that $J = \sum_{i=1}^n f_i B$ and multiplying f_i 's by elements of $\mathcal{O} \setminus m$ we can achieve that $f_i \in I$ and so they generate an ideal $I' := \sum_{i=1}^n f_i C \subset I$. Since m is $(\mathcal{O} \setminus m)$ -divisible, $mJ = mI' \subset I'$ and hence $\bar{I} := I \cap m[T] \subset mJ \subset I'$. Note that I/\bar{I} is the kernel of $\phi \otimes_{\mathcal{O}} R : R[T] \rightarrow C/mC$, and so is finitely generated over $R[T]$ (and hence over $\mathcal{O}[T]$). Choose any finite set of generators $\tilde{g}_1, \dots, \tilde{g}_l \in I/\bar{I}$, lift each \tilde{g}_j to $g_j \in I$ and consider the ideal $I'' = (I', g_1, \dots, g_l)$ in $C[T]$. Then I'' contains \bar{I} and I''/\bar{I} contains I/\bar{I} , and so $I = I''$ is finitely generated.

Finally, let us show that X is S -flat. We already know that X is of finite presentation over S , therefore the flattening theorem of Raynaud-Gruson [RG, 5.2.2]

asserts that X can be flattened by performing a U -modification (and even a U -admissible blow up) on S and replacing X with its strict transform. However, S has no non-trivial U -modifications because T (being the spectrum of a valuation ring) is the only modification of itself. Thus, X has to be S -flat and we conclude the proof. \square

In the first version of the paper, the lemma was formulated in a larger (and incorrect) generality, as was pointed out by D. Rydh. So, let us discuss briefly true and false generalizations.

Remark 2.3.2. (i) Lemma 2.3.1(i) implies that descent data of the form $X_U \times_U \eta \xrightarrow{\sim} X_T \times_T \eta$ is always effective for U -admissible \bar{S} -affine schemes. Some examples show that for general U -admissible schemes the descent of this type is not effective, though it exists as an algebraic space. More generally, Rydh recently showed in [Rydh, §6] that general descent of this type can be made in the category of stacks with quasi-finite diagonal.

(ii) Also, Rydh observed that the flatness assumption in Lemma 2.3.1(iii) is essential for finite presentation. Without flatness finite presentation can be lost after gluing even in the case when \mathcal{O} is a height two valuation ring composed from DVR's A and R and X is a (non-reduced) closed subscheme in S .

We assume again that S is a qcqs scheme with a schematically dense quasi-compact subset U which is closed under generalizations. We will prove a stable modification theorem which strengthens its analog from [Tem2], and we refer to the introduction of loc.cit. for terminology. Our strengthening is in imposing natural restrictions on the base change required in order to construct a stable modification. It is reasonable to expect that in some sense one can preserve the locus U of S over which the given curve is already semi-stable. Since already when U is the generic point of an integral base scheme S one has to allow its finite étale coverings (i.e. one has to allow separable alterations rather than modifications), it seems that one cannot hope for something more restrictive than admitting general U -étale coverings of the base.

Theorem 2.3.3. *Let (C, D) be an S -multipointed curve with semi-stable U -fibers. Then there exists a U -étale covering $S' \rightarrow S$ such that the curve $(C, D) \times_S S'$ admits a stable U -modification.*

Proof. Step 1. *The theorem holds over a semi-valuation ring \mathcal{O} .* More concretely, throughout Step 1 we assume that \mathcal{O} is composed from a local ring (A, m) and a valuation ring R of $K = A/m$, $S = \text{Spec}(\mathcal{O})$ and $U = \text{Spec}(A)$. Set also $T = \text{Spec}(R)$ and $\eta = \text{Spec}(K)$. By [Tem2, 1.5], the theorem is known in the case of a valuation ring, i.e. the case when $m = 0$. Thus, there exists a finite separable extension K'/K with a valuation ring R' lying over R and such that $(C, D) \times_S T'$ admits a stable modification, where $T' = \text{Spec}(R')$. Lift the extension K'/K to a finite étale extension of local rings A'/A , and let \mathcal{O}' be the semi-valuation ring composed from A' and R' . We will show that the stable modification exists over \mathcal{O}' , but let us explain first how this concludes the Step. Clearly, $R' = \cup R_i$ where R_i 's are finitely generated R -subalgebras of R' such that $\text{Frac}(R_i) = K'$. Therefore $\mathcal{O}' = \cup \mathcal{O}_i$ where \mathcal{O}_i is the preimage of R_i in A' . It remains to note that for any i we have that $\text{Spec}(\mathcal{O}_i) \times_S U \xrightarrow{\sim} \text{Spec}(A')$ is étale over U , and by approximation the stable modification exists already over some \mathcal{O}_i .

Now we can work over \mathcal{O}' and to simplify the notation we replace \mathcal{O} by \mathcal{O}' achieving that already $(C_T, D_T) := (C, D) \times_S T$ admits a stable modification $(\overline{C}_T, \overline{D}_T)$. By [Tem2, 1.1] there is a canonical C_T -ample sheaf on \overline{C}_T , namely the sheaf $\mathcal{L}_T := \omega_{(\overline{C}_T, \overline{D}_T)/T}$. Set also $\mathcal{L}_U := \omega_{(C_U, D_U)/U}$ and note that these sheaves agree over η because the formation of ω 's commutes with base changes (see [Tem2, §1]). By Lemma 2.3.1(ii) applied with $\overline{S} = C$, we can glue (C_U, \mathcal{L}_U) and $(\overline{C}_T, \mathcal{L}_T)$ to a U -modification $\overline{C} \rightarrow C$. In addition, \overline{C} is flat and finitely presented over S by Lemma 2.3.1(iii). Clearly, the closed subschemes $D_U \hookrightarrow C_U$ and $\overline{D}_T \hookrightarrow \overline{C}_T$ glue to a closed subscheme $\overline{D} \hookrightarrow \overline{C}$, and checking the S -fibers we obtain that $(\overline{C}, \overline{D})$ is a stable U -modification of (C, D) .

Step 2. The general case. Since (C, D) is semi-stable over an open subscheme of S , we can enlarge U to an open schematically dense qcqs subscheme. Note that by noetherian approximation there exists a scheme S' of finite type over \mathbf{Z} with a morphism $S \rightarrow S'$ such that U and (C, D) are induced from a schematically dense open subscheme $U' \hookrightarrow S'$ and a multipointed curve $(C', D') \rightarrow S'$. Then it suffices to solve our problem for S', U' and (C', D') , so we can assume that S is of finite type over \mathbf{Z} . By [Tem2, 3.3.1], $\mathfrak{S} = \text{RZ}_U(S)$ is a qcqs topological space. For any point $\mathbf{x} = (y, R, \phi) \in \mathfrak{S}$, set $S_{\mathbf{x}} = \text{Spec}(\mathcal{O}_{\mathfrak{S}, \mathbf{x}})$, $U_{\mathbf{x}} = \text{Spec}(\mathcal{O}_{Y, y})$ and $(C_{\mathbf{x}}, D_{\mathbf{x}}) = (C, D) \times_S S_{\mathbf{x}}$. Since the embedding $U \hookrightarrow S$ is obviously decomposable, Proposition 2.2.1 implies that $\mathcal{O}_{\mathfrak{S}, \mathbf{x}}$ is a semi-valuation ring with the semi-fraction ring $\mathcal{O}_{Y, y}$. By Step 1, there exists a $U_{\mathbf{x}}$ -étale covering $S'_{\mathbf{x}} \rightarrow S_{\mathbf{x}}$ such that the $S'_{\mathbf{x}}$ -multipointed curve $(C_{\mathbf{x}}, D_{\mathbf{x}}) \times_{S_{\mathbf{x}}} S'_{\mathbf{x}} \xrightarrow{\sim} (C, D) \times_S S'_{\mathbf{x}}$ admits a stable $U'_{\mathbf{x}}$ -modification for $U'_{\mathbf{x}} = U_{\mathbf{x}} \times_{S_{\mathbf{x}}} S'_{\mathbf{x}}$. Note also that the morphism $S'_{\mathbf{x}} \rightarrow S_{\mathbf{x}}$ is flat and finitely presented by Lemma 2.3.1(iii).

Consider the family $\{S_i\}_{i \in I}$ of all U -modifications of S , and let x_i be the center of \mathbf{x} on S_i . Recall that $\mathcal{O}_{\mathfrak{S}, \mathbf{x}} = \text{injlim } \mathcal{O}_{S_i, x_i}$. By approximation, there exists $i = i(\mathbf{x})$ and a flat finitely presented U -étale morphism $h_{\mathbf{x}} : S' \rightarrow S_i$ such that x_i lies in its image and $(C, D) \times_S S'_i$ admits a stable U -modification. By flatness of $h_{\mathbf{x}}$, $h_{\mathbf{x}}(S')$ is open in S_i , and hence its preimage in \mathfrak{S} is an open neighborhood of \mathbf{x} . Note that in the sequel we can replace i by any larger index k simply by replacing $h_{\mathbf{x}}$ by its base change with respect to the U -modification $S_k \rightarrow S_i$. Since \mathfrak{S} is quasi-compact, there exist finitely many points \mathbf{x}_j , $1 \leq j \leq n$ with associated flat morphisms $h_j : S'_j \rightarrow S_{i_j}$ so that \mathfrak{S} is covered by the preimages of the sets $h_j(S'_j)$. By the above argument we can enlarge all indexes so that $i := i_1 = \dots = i_n$. The open subschemes $h_j(S'_j) \hookrightarrow S_i$ with $1 \leq j \leq n$ cover S_i because their preimages cover \mathfrak{S} , and so $S' := \sqcup_{j=1}^n S'_j$ is a flat cover of S_i . In particular, S' is a U -étale covering of S over which (C, D) possesses a stable U -modification. \square

A scheme version of the reduced fiber theorem of Bosch-Lütkebohmert-Raynaud [BLR, 2.1'], can be proved absolutely similarly.

Theorem 2.3.4. *Let $X \rightarrow S$ be a schematically dominant finitely presented morphism whose U -fibers are geometrically reduced. Then there exists a U -étale covering $S' \rightarrow S$ and a finite U -modification $X' \rightarrow X \times_S S'$ such that X' is flat, finitely presented and has reduced geometric fibers over S' .*

Proof. If S is the spectrum of a valuation ring and U is its generic point then the theorem follows from [Tem2, 3.5.5] (actually it was the content of Steps 2–4 of the loc.cit.). Acting as in Step 1 of the previous proof, we deduce the case when S

is the spectrum of a semi-valuation ring and U is the corresponding local scheme. Then it remains to repeat the argument of Step 2. \square

3. RELATIVE RZ SPACES AND THE DECOMPOSITION THEOREM

Throughout §3, $f : Y \rightarrow X$ is a morphism of schemes and $\mathfrak{X} = \text{Val}_Y(X)$. Later we will also introduce a topological space $\text{Spa}(Y, X)$ and then we will use the notation $\overline{\mathfrak{X}} = \text{Spa}(Y, X)$. Sometimes we will consider another morphism of schemes $f' : Y' \rightarrow X'$ and then $\mathfrak{X}' = \text{Val}_{Y'}(X')$, $\overline{\mathfrak{X}'} = \text{Spa}(Y', X')$.

3.1. Connection to adic spaces. Let A be a ring and B be an A -algebra. R. Huber considers in [Hub1] the set $\text{Spv}(B)$ of all equivalence classes of valuations on B and provides it with the weakest topology in which the sets of the form $\{ | \in \text{Spv}(B) \mid |a| \leq |b| \neq 0 \}$ are open for any $a, b \in B$. Huber proves in [Hub1, 2.2] that the resulting topological space is quasi-compact. Furthermore, he considers the quasi-compact subspace $\text{Spa}(B, A) \subset \text{Spv}(B)$ consisting of the valuations of B with $|A| \leq 1$: see the definition on p. 467 in loc.cit., where one treats A and B as topological rings with discrete topology (note also that Huber actually considers the case when A is an integrally closed subring of B , but this does not really restrict the generality because replacing A by the integral closure of its image in B has no impact on the topological space $\text{Spa}(B, A)$). Actually, the topological space $\text{Spa}(B, A)$ has a much finer structure of an adic space but we will not use it.

Let us generalize the above paragraph to schemes. Note that a valuation on a ring A is defined by its kernel $x \in \text{Spec}(A)$ and the induced valuation on $k(x)$. So, by a *valuation on a scheme* Y we mean a pair $\mathbf{y} = (y, R)$, where $y \in Y$ is a point called the *kernel* of \mathbf{y} and R is a valuation ring of $k(y)$. One can define \mathbf{y} by giving a valuation $|_{\mathbf{y}} : \mathcal{O}_{Y,y} \rightarrow \Gamma_{\mathbf{y}}$ whose kernel is $m_{\mathbf{y}}$. By $\mathcal{O}_{\mathbf{y}}$ we denote the subring of $\mathcal{O}_{Y,y}$ given by the condition $|_{\mathbf{y}} \leq 1$; it is the preimage of R in $\mathcal{O}_{Y,y}$. Remark that $\mathcal{O}_{\mathbf{y}}$ is a semi-valuation ring with the semi-fraction ring $\mathcal{O}_{Y,y}$. Often it is convenient to describe a valuation locally by choosing an affine neighborhood $\text{Spec}(A)$ of y and giving a valuation $A \rightarrow \mathcal{O}_{Y,y} \rightarrow \Gamma_{\mathbf{y}}$ on A .

Furthermore, if $f : Y \rightarrow X$ is a morphism of schemes then by an *X -valuation on Y* we mean a valuation $\mathbf{y} = (y, R)$ provided with a morphism $\phi : S = \text{Spec}(R) \rightarrow X$ which is compatible with the natural morphism $\eta = \text{Spec}(k(y)) \rightarrow X$. Recall that in the valuative criteria of properness/separatedness one considers commutative diagrams of the form

$$\begin{array}{ccc} \eta & \xrightarrow{i} & Y \\ \downarrow & & \downarrow f \\ S & \xrightarrow{\phi} & X \end{array} \quad (1)$$

where $S = \text{Spec}(R)$ is the spectrum of a valuation ring and $\eta = \text{Spec}(K)$ is its generic point, and studies liftings of S to Y . It is easy to see (and will be proved in Lemma 3.2.1) that it suffices to consider only the case when $k(y) \overset{\sim}{\rightarrow} K$ for $y = i(\eta)$ in the valuative criteria. In the latter particular case, diagrams of type (1) are exactly the diagrams which correspond to X -valuations of Y . Note also that an

X -valuation $\mathbf{y} = (y, R, \phi)$ gives rise to the following finer diagram

$$\begin{array}{ccccc} \eta & \longrightarrow & \mathrm{Spec}(\mathcal{O}_{Y,y}) & \longrightarrow & Y \\ \downarrow & & \downarrow & & \downarrow \\ S & \longrightarrow & \mathrm{Spec}(\mathcal{O}_{\mathbf{y}}) & \longrightarrow & X \end{array} \quad (2)$$

Indeed, the center $x \in X$ of R is a specialization the image of y and the induced homomorphism $\mathcal{O}_{X,x} \rightarrow \mathcal{O}_{Y,y} \rightarrow k(y)$ coincides with $\mathcal{O}_{X,x} \rightarrow R \rightarrow \mathrm{Frac}(R) \xrightarrow{\sim} k(y)$. Hence these homomorphisms factor through $\mathcal{O}_{\mathbf{y}}$ (actually, we have just shown that the left square is co-Cartesian).

Let $\mathrm{Spa}(Y, X)$ denote the set of all isomorphism classes of X -valuations on Y . We claim that $\mathrm{Spa}(Y, X)$ depends functorially on f . Indeed, given a morphism $f' : Y' \rightarrow X'$ and a morphism $g : f' \rightarrow f$ consisting of a compatible pair of morphisms $g_Y : Y' \rightarrow Y$ and $g_X : X' \rightarrow X$, there is a natural map $\mathrm{Spa}(g) : \mathrm{Spa}(Y', X') \rightarrow \mathrm{Spa}(Y, X)$ which to a point (y', R', ϕ') associates a point (y, R, ϕ) , where $y = g_Y(y')$, $R = R' \cap k(y)$ and ϕ is defined as follows. The morphism $g_X \circ \phi' : \mathrm{Spec}(R') \rightarrow X$ factors through $\mathrm{Spec}(\mathcal{O}_{X,x})$, where x is the image of the closed point of the source, hence we obtain a homomorphism $\alpha : \mathcal{O}_{X,x} \rightarrow R'$. Since the morphism $\mathrm{Spec}(k(y')) \rightarrow X$ factors uniquely through $\mathrm{Spec}(k(y))$, the image of α is contained in R . So, $g_X \circ \phi'$ factors uniquely through a morphism $\phi : \mathrm{Spec}(R) \rightarrow X$ and the map $\mathrm{Spa}(g)$ is constructed.

If g_Y is an immersion and g_X is separated then $\mathrm{Spa}(g)$ is injective. Indeed, if a point $\mathbf{y} = (y, R, \phi) \in \overline{\mathfrak{X}} := \mathrm{Spa}(Y, X)$ has a non-empty preimage in $\overline{\mathfrak{X}}' := \mathrm{Spa}(Y', X')$, then $y \in Y'$ and any preimage of \mathbf{y} is given by a lifting of $\phi : \mathrm{Spec}(R) \rightarrow X$ to X' , which is unique by the valuative criterion of separatedness. Furthermore, we say that $\overline{\mathfrak{X}}'$ is an *affine subset* of $\overline{\mathfrak{X}}$ if Y' and X' are affine, g_Y is an open immersion and g_X is of finite type. We provide $\overline{\mathfrak{X}}$ with the weakest topology in which all affine subsets are open. Note that if we are given another morphism between morphisms $h : (Y_1 \rightarrow X_1) \rightarrow (Y \rightarrow X)$ with the corresponding map $\mathrm{Spa}(h) : \overline{\mathfrak{X}}_1 \rightarrow \overline{\mathfrak{X}}$, then $Y_1' := Y' \times_Y Y_1$ is a subscheme in Y_1 and $X_1' := X' \times_X X_1$ is separated over X_1 , hence $\overline{\mathfrak{X}}_1' := \mathrm{Spa}(Y_1', X_1')$ embeds into $\overline{\mathfrak{X}}_1$.

Lemma 3.1.1. *Let $\mathrm{Spa}(g) : \overline{\mathfrak{X}}' \rightarrow \overline{\mathfrak{X}}$ and $\mathrm{Spa}(h) : \overline{\mathfrak{X}}_1 \rightarrow \overline{\mathfrak{X}}$ be as above and assume that g_Y is an immersion and g_X is separated.*

- (i) $\overline{\mathfrak{X}}_1'$ is the preimage of $\overline{\mathfrak{X}}'$ under $\mathrm{Spa}(h)$.
- (ii) If X and Y are separated, $\overline{\mathfrak{X}}'$ is an affine subset of $\overline{\mathfrak{X}}$ and both X_1 and Y_1 are affine, then $\overline{\mathfrak{X}}_1'$ is an affine subset of $\overline{\mathfrak{X}}_1$. In particular, if X and Y are separated then the intersection of affine subsets in $\overline{\mathfrak{X}}$ is an affine subset.
- (iii) Affine subsets form a basis of the topology on $\overline{\mathfrak{X}}$, and if X and Y are qcqs then any intersection of two affine subsets is a finite union of affine subsets.
- (iv) If g_Y is an open immersion and g_X is of finite type then $\overline{\mathfrak{X}}'$ is open in $\overline{\mathfrak{X}}$;
- (v) The maps $\mathrm{Spa}(h)$ are continuous.

Proof. The first claim is proved by a straightforward check. If Y and X are separated then $Y' \times_Y Y_1$ and $X' \times_X X_1$ are affine, hence (i) implies (ii). Furthermore, in general (i) implies that the intersection of affine subsets in $\overline{\mathfrak{X}}$ is of the form $\mathrm{Spa}(\overline{Y}, \overline{X})$. Since an affine subset in $\mathrm{Spa}(\overline{Y}, \overline{X})$ is also an affine subset in $\overline{\mathfrak{X}}$, to prove (iii) it suffices to show that any space $\mathrm{Spa}(\overline{Y}, \overline{X})$ (resp. with qcqs \overline{X} and \overline{Y})

is covered by (resp. finitely many) affine subsets. Find open affine (resp. finite) coverings $\bar{X} = \cup \bar{X}_i$ and $\bar{Y} = \cup \bar{Y}_j$ such that each \bar{Y}_j is mapped to some $\bar{X}_{i(j)}$, and note that $\text{Spec}(\bar{Y}, \bar{X})$ is the union of affine subsets $\text{Spa}(\bar{Y}_j, \bar{X}_{i(j)})$. This proves (iii), and the same argument proves (iv). Finally, (v) follows from the fact the preimage of each affine subset is open due to (i) and (iv). \square

We claim that in the affine case the above topology agrees with the topology defined by Huber.

Lemma 3.1.2. *If $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$ are affine then the canonical bijection $\phi : \text{Spa}(Y, X) \rightarrow \text{Spa}(B, A)$ is a homeomorphism.*

Proof. It follows from the definitions of the topologies that the map is continuous, so we have only to establish openness. Let $\bar{\mathfrak{X}}' = \text{Spa}(\text{Spec}(C), \text{Spec}(A'))$ be an affine subset in $\bar{\mathfrak{X}}$, where $\text{Spec}(C)$ is an open subscheme of $\text{Spec}(B)$ and A' is a finitely generated A -algebra. It suffices to prove that $\phi(\bar{\mathfrak{X}}')$ is a neighborhood of each point \mathbf{z} it contains. Replacing A' with its image in C we can assume that it is an A -subalgebra of C generated by $h_1, \dots, h_n \in C$. Note that if $\{U_i\}$ is an open covering of $\text{Spec}(C)$ then the sets $\text{Spa}(U_i, \text{Spec}(A'))$ cover $\bar{\mathfrak{X}}'$. Therefore, shrinking $\text{Spec}(C)$ we can assume that $C = B_b$ for an element $b \in B$. Then $h_i = b_i/b^m$ with $b_i \in B$ and $m \in \mathbf{N}$, and $\phi(\bar{\mathfrak{X}}')$ consists of all valuations of B with $|b_i| \leq |b^m| \neq 0$ for any i . Thus, $\phi(\bar{\mathfrak{X}}')$ is open in $\text{Spa}(B, A)$, and we are done. \square

Since Huber's spaces $\text{Spa}(B, A)$ are qcqs, we obtain the following corollary.

Corollary 3.1.3. *If X and Y are qcqs schemes then the space $\text{Spa}(Y, X)$ is qcqs.*

Let B be a ring provided with a valuation $|\cdot| : B \rightarrow \Gamma \cup \{0\}$, and let $y \in \text{Spec}(B)$ be its kernel. We say that a convex subgroup $\Gamma' \subseteq \Gamma$ *bounds* B , if for any element $b \in B$, there exists an element $h \in \Gamma'$ with $|b| \leq h$. For any such subgroup we can define a valuation $|\cdot|' : B \rightarrow \Gamma'$ by the rule $|x|' = |x|$ if $|x| \in \Gamma'$ and $|x|' = 0$ otherwise. Obviously, the kernel y' of $|\cdot|'$ is a specialization of y . Recall that $|\cdot|'$ is called a *primary specialization* of $|\cdot|$, see [Hub1, 2.3]. Here are simple properties of primary specializations.

Remark 3.1.4. (i) Primary specialization is a transitive operation and the set P of primary specializations of $|\cdot|$ is ordered.

(ii) The set P possesses a minimal element corresponding to the intersection of all subgroups bounding B ; it is called the *minimal primary specialization*.

(iii) A valuation on B is called *minimal* if it has no non-trivial primary specializations. For a valuation given by a point $y \in \text{Spec}(B)$ and a valuation ring $R \subset k(y)$ the following conditions are equivalent: (a) (y, R) is minimal; (b) $k(y)$ is generated by R and the image of B ; (c) the morphism $\text{Spec}(k(y)) \rightarrow \text{Spec}(R) \times \text{Spec}(B)$ is a closed immersion.

(iv) Let $|\cdot| : B \rightarrow \Gamma \cup \{0\}$ be a valuation with kernel y , $\Gamma' \subseteq \Gamma$ be a convex subgroup, and $R \subseteq R'$ be the valuation ring of $k(y)$ corresponding to the induced valuations $k(y) \rightarrow \Gamma$ and $k(y) \rightarrow \Gamma \rightarrow \Gamma/\Gamma'$. Then the following conditions are equivalent: (a) there exists a primary specialization $|\cdot|'$ corresponding to Γ' ; (b) the image of B in $k(y)$ is contained in R' ; (c) the morphism $y \rightarrow \text{Spec}(B)$ extends to a morphism $\text{Spec}(R') \rightarrow \text{Spec}(B)$. Moreover, if the conditions are satisfied then the kernel y' of $|\cdot|'$ is the center of R' on $\text{Spec}(B)$. The equivalences (a) \Leftrightarrow (b) and

(b) \Leftrightarrow (c) are obvious. As for the additional claim, we note that the center of R' corresponds to the kernel of the homomorphism $B \rightarrow R' \rightarrow R'/m_{R'}$, and the latter consists of the elements $b \in B$ with $|b| \notin \Gamma'$, i.e. coincides with the kernel of $|\cdot|'$.

Let, more generally, $\mathbf{y} = (y, R)$ be a valuation on a scheme Y . By a *primary specialization* of \mathbf{y} we mean a valuation $\bar{\mathbf{y}} = (\bar{y}, \bar{R})$ such that \bar{y} is a specialization of y and the valuation $|\cdot|_{\bar{\mathbf{y}}}$ on $\mathcal{O}_{Y, \bar{y}}$ is a primary specialization of the valuation induced from \mathbf{y} via the homomorphism $\mathcal{O}_{Y, \bar{y}} \rightarrow \mathcal{O}_{Y, y}$. Equivalently, if $\text{Spec}(A)$ is an affine neighborhood of \bar{y} (and hence of y) then the valuation induced by (\bar{y}, \bar{R}) on A is a primary specialization of the valuation induced by (y, R) .

Lemma 3.1.5. *Let (y, R) be a valuation on a separated scheme Y .*

- (i) *The set of primary valuations of (y, R) is totally ordered by specialization;*
- (ii) *If Y is also quasi-compact then (y, R) admits a minimal primary specialization.*

Proof. We claim that (i) follows from Remark 3.1.4(iv). Indeed, for any R' with $R \subseteq R' \subseteq k(y)$ there exists at most one possibility to extend y to a morphism $\text{Spec}(R') \rightarrow Y$. So, if we have two primary specializations (y_1, R_1) and (y_2, R_2) corresponding to valuation rings $R \subseteq R', R'' \subseteq k(y)$, then without loss of generality we have that $R' \subseteq R''$ and the unique morphism $\text{Spec}(R'') \rightarrow Y$ is obtained by localizing the morphism $\text{Spec}(R') \rightarrow Y$. Thus, y_1 is a specialization of y_2 , and everything reduces to affine theory of primary specializations on \mathcal{O}_{Y, y_1} , see Remark 3.1.4(i). To prove (ii) we note that (y, R) admits a minimal primary specialization because if $\{(y_i, R_i)\}_{i \in I}$ denotes the set of all primary specializations then the set of kernels $\{y_i\}_{i \in I}$ is totally ordered with respect to specialization. By quasi-compactness there exists a point $\bar{y} \in Y$ which is a specialization all y_i 's. So, the claim reduces to the affine theory on $\mathcal{O}_{Y, \bar{y}}$, see Remark 3.1.4(ii). \square

Finally, taking a morphism $f : Y \rightarrow X$ into account, by a *primary specialization* of an X -valuation $\mathbf{y} = (y, R, \phi)$ we mean an X -valuation $\bar{\mathbf{y}} = (\bar{y}, \bar{R}, \bar{\phi})$ such that (\bar{y}, \bar{R}) is a primary specialization of (y, R) and the image of $\bar{\phi}$ in X is contained in the image of ϕ in X . Primary specialization is a particular case of a specialization relation in $\text{Spa}(Y, X)$. An X -valuation (y, R, ϕ) (resp. a valuation (y, R)) on Y is called *minimal* if it has no non-trivial primary specializations.

Lemma 3.1.6. *Let (y, R, ϕ) be an X -valuation on Y . Then any primary specialization (\bar{y}, \bar{R}) of the valuation (y, R) admits at most one extension to a primary specialization $(\bar{y}, \bar{R}, \bar{\phi})$ of (y, R, ϕ) , and the extension exists if and only if $f(\bar{y})$ belongs to the image of ϕ . The latter is automatically the case when X is separated.*

Proof. Obviously, the assertion on $f(\bar{y})$ is necessary for an extension to exist. Furthermore, by Remark 3.1.4(iv) there exists a valuation ring R' with $R \subseteq R' \subseteq k(y)$ such that y extends to a morphism $\text{Spec}(R') \rightarrow Y$ with \bar{y} being the image of the closed point. If X is separated then the induced map $\text{Spec}(R') \rightarrow X$ must coincide with the corresponding localization of $\phi : \text{Spec}(R) \rightarrow X$, hence we obtain the last assertion of the lemma. The remaining claims are local at the center $x \in X$ of R (i.e. the image of the closed point of ϕ). So, we can replace X and Y with a neighborhood of x and its preimage achieving that the schemes become separated. The uniqueness is now clear. To establish existence we should check that the image of the homomorphism $\mathcal{O}_{X, x} \rightarrow \mathcal{O}_{Y, \bar{y}} \rightarrow k(\bar{y})$ is in \bar{R} . The latter follows from the following two facts: (a) by existence of ϕ the image of $\mathcal{O}_{X, x}$ in $k(y)$ is in R , (b) \bar{R}

is induced from R in the sense that an element $\bar{f} \in \mathcal{O}_{Y, \bar{y}}$ satisfies $\bar{f}(\bar{y}) \in \bar{R}$ if and only if $f(y) \in R$. \square

The lemma shows that we can actually ignore ϕ when X is separated. In particular, minimality of (y, R, ϕ) is then equivalent to that of (y, R) .

Corollary 3.1.7. *Let $f : Y \rightarrow X$ be a separated morphism of qcqs schemes and $\mathbf{y} = (y, R, \phi)$ be an X -valuation on Y . Then*

i) the set of primary specializations of \mathbf{y} is totally ordered and contains a minimal element,

ii) \mathbf{y} is minimal if and only if the morphism $h : \text{Spec}(k(y)) \rightarrow Y \times_X \text{Spec}(R)$ is a closed immersion.

Proof. The claim is local at the center of $x \in X$ of R (with respect to ϕ), hence we can assume that X and, hence, Y are separated. Then primary specializations of \mathbf{y} can be identified with primary specializations of the valuation (y, R) , hence (i) follows from Lemma 3.1.5. To prove (ii) we note that as soon as X is separated, h is a closed immersion if and only if the morphism $\text{Spec}(k(y)) \rightarrow Y \times \text{Spec}(R)$ is a closed immersion. Hence the claim follows from Remark 3.1.4(iii). \square

Until the end of §3, we assume that $f : Y \rightarrow X$ is a separated morphism of qcqs schemes, unless the contrary is said explicitly. We define the subset $\mathfrak{X} = \text{Val}_Y(X) \subset \bar{\mathfrak{X}}$ as the set of all minimal valuations and note that in view of Lemma 3.1.7 this agrees with the case studied in 2.2. We do not introduce $\text{Val}_Y(X)$ when f is not separated: although the formal definition makes sense, it is not clear if the obtained object is interesting. Note also that in affine situation such subsets were considered by Huber, see [Hub1, 2.6 and 2.7]. We provide \mathfrak{X} with the induced topology. The following lemma follows easily from the valuative criterion of properness and Lemma 3.1.1.

Lemma 3.1.8. *(i) If X' is a Y -modification of X then there are natural homeomorphisms $\text{Spa}(Y, X') \xrightarrow{\sim} \text{Spa}(Y, X)$ and $\text{Val}_Y(X') \xrightarrow{\sim} \text{Val}_Y(X)$.*

(ii) If X' is an open subscheme of X then its preimage in $\text{Val}_Y(X)$ (resp. $\text{Spa}(Y, X)$) is canonically homeomorphic to $\text{Val}_{Y'}(X')$ (resp. $\text{Spa}(Y', X')$), where $Y' = X' \times_X Y$.

Remark 3.1.9. (i) If $f' : Y' \rightarrow X'$ and $f : Y \rightarrow X$ are separated morphisms of qcqs schemes, and $g : f' \rightarrow f$ is a morphism such that g_Y is an open immersion and g_X is separated and of finite type, then $\text{Spa}(Y', X')$ maps homeomorphically onto an open subspace of $\bar{\mathfrak{X}}$. However, it may (and usually does) happen that the image of $\text{Val}_{Y'}(X')$ in $\bar{\mathfrak{X}}$ is not contained in \mathfrak{X} . The problem originates from the fact that a minimal valuation on Y' may admit non-trivial primary specializations on Y .

(ii) There exists a natural contraction $\pi_{\mathfrak{X}} : \bar{\mathfrak{X}} \rightarrow \mathfrak{X}$ which maps any valuation to its minimal primary specialization, but it is a difficult fact that $\pi_{\mathfrak{X}}$ is continuous.

(iii) Using $\pi_{\mathfrak{X}}$ we can extend Val to a functor by composing $\text{Spa}(g)$ with the contraction $\pi_{\mathfrak{X}}$ as $\text{Val}(g) : \mathfrak{X}' \hookrightarrow \bar{\mathfrak{X}}' \rightarrow \bar{\mathfrak{X}} \rightarrow \mathfrak{X}$. However, we do not know that it is continuous until continuity of $\pi_{\mathfrak{X}}$ is established.

Actually, the above problems are closely related, and we will solve them only in the end of §3.3. Recall that if X and Y are qcqs then so are $\text{Spa}(Y, X)$ and $\text{RZ}_Y(X)$ (by Corollary 3.1.3 and [Tem2, 3.3.1]). Here is a partial (so far) result for $\text{Val}_Y(X)$.

Proposition 3.1.10. *Assume that $f : Y \rightarrow X$ is a separated morphism of qcqs schemes. Then the spaces $\text{Val}_Y(X)$ is quasi-compact and the map $\psi : \text{Val}_Y(X) \rightarrow \text{RZ}_Y(X)$ is continuous.*

Proof. Let $\{\mathfrak{X}_i\}_{i \in I}$ be an open covering of \mathfrak{X} . Find open sets $\overline{\mathfrak{X}}_i \subset \overline{\mathfrak{X}}$ such that $\mathfrak{X}_i = \overline{\mathfrak{X}}_i \cap \mathfrak{X}$. Since any point of $\overline{\mathfrak{X}}$ has a specialization in \mathfrak{X} by Corollary 3.1.7, $\{\overline{\mathfrak{X}}_i\}_{i \in I}$ is a covering of $\overline{\mathfrak{X}}$. By quasi-compactness of $\overline{\mathfrak{X}}$, we can find a subcovering $\{\overline{\mathfrak{X}}_i\}_{i \in J}$ with a finite J , and then $\{\mathfrak{X}_i\}_{i \in J}$ is a finite covering of \mathfrak{X} . Thus, \mathfrak{X} is quasi-compact.

We claim that for any Y -modification $X' \rightarrow X$, the map $\phi : \mathfrak{X} \rightarrow X'$ is continuous. Indeed, if $U \subset X'$ is open then its preimage in $\overline{\mathfrak{X}}$ is the open subspace $\overline{\mathfrak{X}}' \xrightarrow{\sim} \text{Spa}(Y \times_{X'} U, U)$. Therefore, the preimage of U in \mathfrak{X} is the open set $\overline{\mathfrak{X}}' \cap \mathfrak{X}$, as required. Continuity of the maps ϕ (for each X') implies that the map $\psi : \mathfrak{X} \rightarrow \text{RZ}_Y(X)$ is continuous. \square

3.2. Valuative criteria. In the sequel, we will need to strengthen the classical valuative criteria of separatedness and properness, [EGA, II, 7.2.3 and 7.3.8]. Our aim is to show that it suffices to consider valuative diagrams of specific types. We say that a morphism is compatible with a commutative diagram, if the diagram remains commutative after adjoining this morphism. Throughout §3.2 f is not assumed to be separated.

Lemma 3.2.1. *Keep the notation of diagram (1) and set $K' = k(i(\eta))$ and $R' = R \cap K$. Then diagram (1) completes uniquely to a commutative diagram*

$$\begin{array}{ccccc} \text{Spec}(K) & \longrightarrow & \text{Spec}(K') & \longrightarrow & Y \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(R) & \longrightarrow & \text{Spec}(R') & \longrightarrow & X \end{array}$$

and any morphism $h : \text{Spec}(R) \rightarrow Y$ compatible with the above diagram is induced from a morphism $h' : \text{Spec}(R') \rightarrow Y$ compatible with the diagram. In addition, h determines h' uniquely.

Proof. The morphism $\text{Spec}(K) \rightarrow Y$ obviously factors through $\text{Spec}(K')$. The morphism $\text{Spec}(R) \rightarrow X$ factors through $\text{Spec}(\mathcal{O}_{X,x})$, where x is the image of the closed point of $\text{Spec}(R)$. The image of $\mathcal{O}_{X,x}$ in $R \subset K$ is contained in K' , hence the morphism $\text{Spec}(R) \rightarrow X$ factors through $\text{Spec}(R')$. By the same reasoning, a morphism $h : \text{Spec}(R) \rightarrow Y$ compatible with the diagram factors through $h' : \text{Spec}(R') \rightarrow Y$, and they both are determined uniquely by the image of the closed point in Y . \square

Lemma 3.2.2. *Keep the notation of diagram (1) and assume that $R \subseteq R' \subseteq K$ is such that the morphism $\text{Spec}(R') \rightarrow X$ admits a lifting $g : \text{Spec}(R') \rightarrow Y$ compatible*

with the diagram. Let \tilde{K} be the residue field of R' and \tilde{R} be the image of R in \tilde{K} .

$$\begin{array}{ccccc}
 & & \text{Spec}(K) & \longrightarrow & Y \\
 & & \downarrow & \nearrow g & \downarrow f \\
 \text{Spec}(\tilde{K}) & \longrightarrow & \text{Spec}(R') & & \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Spec}(\tilde{R}) & \longrightarrow & \text{Spec}(R) & \longrightarrow & X
 \end{array}$$

Then any morphism $\tilde{h} : \text{Spec}(\tilde{R}) \rightarrow Y$ compatible with the above diagram is induced from a morphism $h : \text{Spec}(R) \rightarrow Y$ compatible with the diagram and \tilde{h} determines h uniquely.

Proof. Consider a morphism $\tilde{h} : \text{Spec}(\tilde{R}) \rightarrow Y$ compatible with the diagram. It suffices to show that it factors through $\text{Spec}(R)$, since the uniqueness is again trivial. Let y be the image of the closed point of $\text{Spec}(\tilde{R})$, so \tilde{h} induces a homomorphism $\mathcal{O}_{Y,y} \rightarrow \tilde{R}$. Since y is a specialization of the image y' of the closed point of $\text{Spec}(R')$, we have also a homomorphism $\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{Y,y'} \rightarrow R'$. Then the compatibility implies that the image of $\mathcal{O}_{Y,y}$ in $\tilde{K} = R'/m_{R'}$ lies in \tilde{R} . Therefore, the image of $\mathcal{O}_{Y,y}$ in R' lies in R which is the preimage of \tilde{R} under $R' \rightarrow \tilde{K}$, and we obtain that the homomorphism $\mathcal{O}_{Y,y} \rightarrow \tilde{R}$ factors through R . It gives the desired morphism $h : \text{Spec}(R) \rightarrow Y$. \square

Note that Lemma 3.2.1 implies that it suffices to consider only the case when $k(i(\eta)) \xrightarrow{\sim} K$ in the valuative criteria (i.e. it suffices to take valuative diagrams corresponding to the elements of $\text{Spa}(Y, X)$), and then Lemma 3.2.2 and Remark 3.1.4(iv) imply that it even suffices to consider only the valuative diagrams corresponding to the elements of $\text{Val}_Y(X)$. It is also well known that in the valuative criteria one can restrict to the case when the image of η lies in a given dense subset which is closed under generalization (e.g. the generic point of an irreducible scheme), and such strengthening is the main issue of the following proposition.

Proposition 3.2.3. *Assume that $h : Z \rightarrow Y$ and $f : Y \rightarrow X$ are morphisms of qcqs schemes and consider the natural map $\bar{\psi} : \text{Spa}(Z, Y) \rightarrow \text{Spa}(Z, X)$.*

- (i) *f is separated if and only if $\bar{\psi}$ is injective.*
- (ii) *Assume that f is of finite type. Then f is proper if and only if $\bar{\psi}$ is bijective.*
- (iii) *If f and h are separated then $\bar{\psi}$ induces a map $\psi : \text{Val}_Z(Y) \rightarrow \text{Val}_Z(X)$, and $\bar{\psi}$ is bijective if and only if ψ is bijective.*

Proof. First we prove (iii). If $\mathbf{z} = (z, R, \phi_Y)$ is a point in $\text{Val}_Z(Y)$ then the morphism $z \rightarrow \text{Spec}(R) \times_Y Z$ is a closed immersion. But the target is a closed subscheme in $\text{Spec}(R) \times_X Z$ by separatedness of f , and hence $\bar{\psi}(\mathbf{z})$ is also a minimal valuation. Thus, $\bar{\psi}$ induces a map ψ between the subsets Val . Next we relate the fibers of ψ and $\bar{\psi}$. Consider any point $\mathbf{z} \in \text{Spa}(Z, X)$ and let $\mathbf{z}_0 \in \text{Val}_Z(X)$ be its minimal primary specialization. Then Lemma 3.2.2 implies that the sets $\bar{\psi}^{-1}(\mathbf{z})$ and $\psi^{-1}(\mathbf{z}_0)$ are naturally bijective, and this proves (iii).

We will deal with (i) and (ii) simultaneously. The direct implications follow from the standard valuative criteria. We will prove the opposite implications (which are

refined valuative criteria) by getting a contradiction. So, suppose that f is not separated in (i), or of finite type, separated and not proper in (ii) (if f is not separated in (ii) then $\bar{\psi}$ cannot be bijective by (i)). By the standard valuative criterion and Lemma 3.2.1, there exists an element $\mathbf{y} = (y, R_y, \phi_y) \in \text{Spa}(Y, X)$ such that the number of liftings of the morphism $\phi_y : \text{Spec}(R_y) \rightarrow X$ to Y is at least two in (i) or zero in (ii). Let x denote the center of R_y on X .

By [EGA I, 6.6.5], there exists a point $z \in Z$ for which $h(z)$ is a generalization of y , and so a homomorphism $\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{Z,z} \rightarrow k(z)$ arises. Let R' be any valuation ring of $k(z)$ which dominates the image of $\mathcal{O}_{Y,y}$. It gives rise to an element $(z, R', \phi') \in \text{Spa}(Z, Y)$ centered on y . Choose a valuation ring \tilde{R} of the residue field \tilde{K} of R' such that \tilde{R} dominates the valuation ring R_y of $k(y) \subset \tilde{K}$, and define a valuation ring R of $k(z)$ as the composition of R' and \tilde{R} . The compatible homomorphisms $\mathcal{O}_{X,x} \rightarrow \mathcal{O}_{Y,y} \rightarrow R'$ and $\mathcal{O}_{X,x} \rightarrow R_y \rightarrow \tilde{R}$ induce a homomorphism $\mathcal{O}_{X,x} \rightarrow R$, and we obtain the following commutative diagrams.

$$\begin{array}{ccc}
 \text{Spec}(k(z)) & \longrightarrow & Z \\
 \downarrow & & \downarrow \\
 \text{Spec}(\tilde{K}) & \longrightarrow & \text{Spec}(R') \xrightarrow{\phi'} Y \\
 \downarrow & & \downarrow \\
 \text{Spec}(\tilde{R}) & \longrightarrow & \text{Spec}(R) \xrightarrow{\phi_x} X
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{Spec}(\tilde{K}) & \longrightarrow & \text{Spec}(k(y)) \longrightarrow Y \\
 \downarrow & & \downarrow \\
 \text{Spec}(\tilde{R}) & \longrightarrow & \text{Spec}(R_y) \longrightarrow X
 \end{array}$$

Lemma 3.2.1 implies that there is a one-to-one correspondence between morphisms $\text{Spec}(R_y) \rightarrow Y$ and $\text{Spec}(\tilde{R}) \rightarrow Y$ compatible with the right diagram, and by Lemma 3.2.2, the latter morphisms are in one-to-one correspondence with the morphisms $\phi : \text{Spec}(R) \rightarrow Y$ compatible with the left diagram. So, there are at least two such ϕ 's in (i) and there is no such ϕ in (ii). Note that $\mathbf{z} = (z, R, \phi_x)$ is an element in $\text{Spa}(Z, X)$, and any morphism ϕ as above gives a preimage of \mathbf{z} in $\text{Spa}(Z, Y)$. We obtain that in the case (i), \mathbf{z} has at least two preimages and so $\bar{\psi}$ is not injective. The same argument would prove (ii) if we also know that, conversely, any preimage of \mathbf{z} in $\text{Spa}(Z, Y)$ comes from ϕ as above. In other words, we want to show that any lift of ϕ_x to $\tilde{\phi} : \text{Spec}(R) \rightarrow Y$ is compatible with the whole left diagram, and this actually reduces to compatibility of $\tilde{\phi}$ with ϕ' . Note that $Y \rightarrow X$ is separated by the already established case (i), and the valuative criterion of separatedness implies that the morphism ϕ' is uniquely determined by the morphisms $\text{Spec}(k(z)) \rightarrow Y$ and $\text{Spec}(R') \rightarrow X$. So, compatibility of $\tilde{\phi}$ with ϕ' is automatic. \square

3.3. Affinoid domains. Let $f' : Y' \rightarrow X'$ be another separated morphism of qcqs schemes and $g : f' \rightarrow f$ be a morphism. Recall that we defined in §3.1 a continuous map $\text{Spa}(g) : \bar{\mathfrak{X}}' \rightarrow \bar{\mathfrak{X}}$ which was shown to be injective if g_Y is an immersion and g_X is separated. However, our definition of a map $\text{Val}(g) : \mathfrak{X}' \rightarrow \mathfrak{X}$ was rather cumbersome because even if $\text{Spa}(g)$ is injective, it does not have to respect the subspaces Val in the spaces Spa . The following proposition gives a criterion when $\text{Spa}(g)$ does respect Val 's.

Proposition 3.3.1. *Suppose that g_Y is an open immersion and g_X is separated. Then $\mathrm{Spa}(g)(\mathfrak{X}') \subset \mathfrak{X}$ if and only if the locally closed immersion $(g_Y, f') : Y' \rightarrow Y \times_X X'$ is a closed immersion, in which case one actually has that $\mathfrak{X}' = \mathrm{Spa}(g)^{-1}(\mathfrak{X})$.*

Proof. Suppose that $h := (g_Y, f')$ is a closed immersion. Let $\mathbf{y}' = (y', R', \phi) \in \overline{\mathfrak{X}'}$ be a point with $\eta' = \mathrm{Spec}(k(y'))$ and $S' = \mathrm{Spec}(R')$, and let $\mathbf{y} = (y, R, \phi)$ be its image in $\overline{\mathfrak{X}}$. By Lemma 3.1.7(ii), \mathbf{y}' is minimal if and only if the natural morphism $\eta' \rightarrow Y' \times_{X'} S'$ is a closed immersion. By closedness of h , the latter happens if and only if the composition morphism $\eta' \rightarrow Y' \times_{X'} S' \rightarrow (Y \times_X X') \times_{X'} S' \xrightarrow{\sim} Y \times_X S'$ is a closed immersion. The latter happens if and only if \mathbf{y} is minimal because $k(y) \xrightarrow{\sim} k(y')$ and hence $R = R' \cap k(y) = R'$. Thus, under our assumption on h , minimality of \mathbf{y}' is equivalent to minimality of its image. This establishes the inverse implication in the proposition, and the complement.

It remains to show that if h is not a closed immersion then $\mathrm{Spa}(g)$ does not respect the subsets Val . Note that h is a locally closed immersion because g_Y is an open immersion, and assume that h is not a closed immersion. Set $Z = Y \times_X X'$ and find a Z -valuation $\mathbf{y}' = (y', R', \phi')$ of Y' such that the morphism $\phi' : \mathrm{Spec}(R') \rightarrow Z$ cannot be lifted to a morphism $\mathrm{Spec}(R') \rightarrow Y'$. Replacing \mathbf{y}' by its minimal primary specialization, we achieve that \mathbf{y}' is minimal and $R' \subsetneq k(y')$. Clearly \mathbf{y}' defines an X' -valuation $\mathbf{y} = (y', R', \phi)$ on Y' with $\phi = \mathrm{pr}_{X'} \circ \phi'$, and \mathbf{y} is minimal because any its non-trivial primary specialization corresponds to a lifting $\mathrm{Spec}(R'') \rightarrow Y'$ for some $R' \subsetneq R'' \subsetneq k(y)$ and such a lifting would induce a lifting $\mathrm{Spec}(R'') \rightarrow Z$ corresponding to a non-trivial primary specialization of \mathbf{y}' . Thus, $\mathbf{y} \in \mathfrak{X}'$, but $\mathrm{Spa}(g)(\mathbf{y})$ is not a minimal X -valuation on Y because the morphism $\mathrm{Spec}(R') \rightarrow X$ lifts to the morphism $\mathrm{pr}_Y \circ \phi : \mathrm{Spec}(R') \rightarrow Y$. \square

Let us assume that g_Y is an open immersion and g_X is separated and of finite type. We saw that if h is a closed immersion then \mathfrak{X}' is naturally identified with a quasi-compact open subset of \mathfrak{X} via $\mathrm{Spa}(g)$, and we say in this case that \mathfrak{X}' is an *open subdomain* of \mathfrak{X} . If, in addition, X' and Y' can be chosen to be affine then we say that \mathfrak{X}' is an *affinoid subdomain* of \mathfrak{X} . Note also that the situation described in the proposition appears in Deligne's proof of Nagata compactification theorem under the name of quasi-dominance. (Recall that by a quasi-dominance of Y over X' one means an open subscheme $Y' \subset Y$ and a morphism $Y' \rightarrow X'$ such that the morphism $Y' \rightarrow Y \times_X X'$ is a closed immersion, see [Con, §2].) The notion of quasi-dominance plays a central role in Deligne's proof. We list simple properties of open and affinoid subdomains in the following lemma and stress that it will be much more difficult to prove that open subdomains are preserved under taking finite unions (in a sense, this is a typical situation in algebraic geometry that preimages, intersections, projective limits, etc., are much easier for study than pushouts, images, direct limits, etc.).

Lemma 3.3.2. *Open subdomains are transitive and are preserved by finite intersections. Moreover, the intersection of open subdomains $\mathrm{Val}_{Y_i}(X_i)$ with $i \in \{1, 2\}$ is the open subdomain $\mathrm{Val}_{Y_1 \cap Y_2}(X_1 \times_X X_2)$. In particular, if X is separated and \mathfrak{X}_i 's are affinoid then \mathfrak{X}_{12} is affinoid.*

Proof. This follows from the analogous Lemma 3.1.1 concerning the spaces Spa . \square

The following remark will not be used in the sequel.

Remark 3.3.3. (i) Our definition of RZ spaces is a straightforward generalization of the classical one. It is also possible to define RZ spaces directly as follows: an affinoid space is a topological space $\mathfrak{X} = \text{Val}_B(A)$ provided with two sheaves of rings $\mathcal{O}_{\mathfrak{X}} \subset \mathcal{M}_{\mathfrak{X}}$ (which can be defined in a natural way), and general spaces are pasted from affinoid ones along affinoid subdomains.

(ii) The following example illustrates a difference between adic and Riemann-Zariski spaces. Let k be a field, $A = B = A' = k[T]$, $B' = k[T, T^{-1}]$ and $\mathfrak{X}, \mathfrak{X}', \bar{\mathfrak{X}}, \bar{\mathfrak{X}}'$ are as above. Then $\bar{\mathfrak{X}}'$ is a rational subdomain in $\bar{\mathfrak{X}}$ in the sense of [Hub2]. From other side, \mathfrak{X}' is not an affinoid domain in \mathfrak{X} . Note that actually $(\mathfrak{X}', \mathcal{O}_{\mathfrak{X}'}) \xrightarrow{\sim} (\mathfrak{X}, \mathcal{O}_{\mathfrak{X}}) \xrightarrow{\sim} X := \text{Spec}(A)$, but the sheaves $\mathcal{M}_{\mathfrak{X}}$ and $\mathcal{M}_{\mathfrak{X}'}$ are not isomorphic at the point $x \in X$ with $T = 0$. This can happen because the local (and even a valuation) ring $\mathcal{O}_{X,x}$ can be provided with two different structures of a semi-valuation ring by choosing semi-fraction rings $\mathcal{M}_{\mathfrak{X}',x} = k(T)$ or $\mathcal{M}_{\mathfrak{X},x} = \mathcal{O}_{X,x}$. (See also Remark 2.1.3(ii).)

Theorem 3.3.4. *The affinoid subdomains of \mathfrak{X} form a basis of its topology.*

Proof. It follows from Lemma 3.3.2 that we should prove that for any affine subset $\bar{\mathfrak{X}}_0 = \text{Spa}(B_0, A_0)$ in $\bar{\mathfrak{X}}$ and a point $\mathbf{y} = (y, R, \phi) \in \mathfrak{X} \cap \bar{\mathfrak{X}}_0$ there exists an affinoid subdomain $\text{Val}_{\bar{Y}}(\bar{X})$ containing \mathbf{y} and contained in $\bar{\mathfrak{X}}_0$. Moreover, we can assume that $X = \text{Spec}(A)$ is affine because \mathfrak{X} is covered by open subdomains of the form $\text{Val}_{Y'}(X')$, where $X' = \text{Spec}(A)$ is an open subscheme of X and $Y' = X' \times_X Y$. In order to construct $\text{Val}_{\bar{Y}}(\bar{X})$ as required we will extend diagram (2) to the following one, where $\bar{Y} = \text{Spec}(\bar{B})$ and $\bar{X} = \text{Spec}(\bar{A})$ will be finally defined in the end of the proof. Recall that $\mathcal{O}_{\mathbf{y}}$ is a semi-valuation ring with semi-fraction ring $\mathcal{O}_{Y,y}$ and such that $\mathcal{O}_{\mathbf{y}}/m_{\mathbf{y}} = R$.

$$\begin{array}{ccccccc} \text{Spec}(k(y)) & \longrightarrow & \text{Spec}(\mathcal{O}_{Y,y}) & \longrightarrow & \bar{Y} & \longrightarrow & Y \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(R) & \longrightarrow & \text{Spec}(\mathcal{O}_{\mathbf{y}}) & \longrightarrow & \bar{X} & \longrightarrow & X \end{array}$$

Since $\text{Spec}(R) \times_X Y$ is closed in $\text{Spec}(R) \times Y$ by separatedness of X , Lemma 3.1.7(ii) implies that the morphism $h : \text{Spec}(k(y)) \rightarrow \text{Spec}(R) \times Y$ is a closed immersion. To explain the strategy of the proof we remark that the morphism $\text{Spec}(\mathcal{O}_{Y,y}) \rightarrow \text{Spec}(\mathcal{O}_{\mathbf{y}}) \times Y$ is a closed immersion (actually it can be proved by the same argument as we use below), and our strategy will be to approximate $\mathcal{O}_{\mathbf{y}}$ and $\mathcal{O}_{Y,y}$ by A -rings \bar{A} and \bar{B} so that \bar{A} is finitely generated over A , $\bar{Y} = \text{Spec}(\bar{B})$ is a neighborhood of y and $\bar{Y} \rightarrow \bar{X} \times Y$ is a closed immersion.

It will be more convenient to work with affine schemes and Y is the only non-affine scheme in our consideration, so let us cover Y with open affine subschemes $Y_i = \text{Spec}(B_i), Z_j = \text{Spec}(C_j)$, where $1 \leq i \leq n, 1 \leq j \leq m, y \in Y_i$ and $y \notin Z_j$. Since $\text{Spec}(B_0)$ contains y by our assumptions, we also set $Y_0 = \text{Spec}(B_0)$. For each i , h factors through a closed immersion $\text{Spec}(k(y)) \rightarrow \text{Spec}(R) \times Y_i$, hence the images of R and B_i generate $k(y)$. Now, we will find a neighborhood $\bar{Y} = \text{Spec}(\bar{B})$ of y which is contained in all Y_i 's and satisfies the following condition: for each i , \bar{B} is a localization of the form $(B_i)_{f_i}$ and, the most important, we have that $f_i(y) \notin m_R$. Let us (until the end of this paragraph only) call *R-localization* for localization of an affine neighborhood $\text{Spec}(C)$ of y at an element f such that $f(y) \notin m_R$. Obviously,

R -localizations are transitive and we claim that the family of R -localizations of each Y_i form a basis of neighborhoods of y . Indeed, for any element $f \in B_i$ with $f(y) \neq 0$ we can find $g \in B_i$ with $f(y)g(y) \notin m_R$ (we use that $B_i(y)$ generates $k(y)$ over R , so it contains elements of arbitrary large valuation). Thus, $(B_i)_{fg}$ is an R -localization of B_i where f is inverted and we obtain that the maximal (infinite) R -localization of B_i is actually $\mathcal{O}_{Y,y}$. Now, set $\text{Spec}(B) = \bigcap_{i=1}^n Y_i$ and find R -localizations $Y'_i = \text{Spec}((B_i)_{g_i})$ contained in $\text{Spec}(B)$, and let $\bar{Y} = \text{Spec}(\bar{B})$ be an R -localization of $\text{Spec}(B)$ contained in all Y'_i . Then \bar{Y} is an R -localization of each Y'_i , hence an R -localization of each Y_i too. So, $\bar{B} = (B_i)_{f_i}$ is as required.

Let \bar{A} be the preimage of R under the character $\bar{B} \rightarrow k(y)$ corresponding to y . Clearly \bar{A} contains each element f_i^{-1} , hence the ring $\bar{B}(y) = B_i(y)[f_i^{-1}(y)]$ is generated by $\bar{A}(y)$ and $B_i(y)$. So, we obtain epimorphisms $\bar{A} \otimes B_i \rightarrow k(y)$, and then the homomorphisms $h_i : \bar{A} \otimes B_i \rightarrow \bar{B}$ are also surjective because \bar{A} contains the kernel p_y of $\bar{B} \rightarrow k(y)$. In particular, each morphism $\bar{Y} \rightarrow \bar{X} \times Y_i$ is a closed immersion. We claim that actually, $\alpha : \bar{Y} \rightarrow \bar{X} \times Y$ is a closed immersion, and to prove this we should check in addition that the morphisms $\alpha_j : \bar{Y} \times_Y Z_j \rightarrow \bar{X} \times Z_j$ with $1 \leq j \leq m$ are closed immersion. By separatedness of Y the source is affine, hence $\bar{Y} \times_Y Z_j = \text{Spec}(\bar{C}_j)$ where \bar{C}_j is generated by the images of $c_j : C_j \rightarrow \bar{C}_j$ and $b_j : \bar{B} \rightarrow \bar{C}_j$. Since our claim about α would follow if we prove that the homomorphisms $h'_j : \bar{A} \otimes C_j \rightarrow \bar{C}_j$ are surjective, it remains only to prove that for each j the image of h'_j contains the image of b_j . Since $y \in \bar{Y}$ and $y \notin Z_j$ we have that $b_j(p_y)\bar{C}_j = \bar{C}_j$, and hence the equality $\bar{C}_j = b_j(\bar{B})c_j(C_j)$ can be strengthened as $\bar{C}_j = b_j(p_y)c_j(C_j)$, i.e. \bar{C}_j is actually generated by $b_j(p_y)$ and $c_j(C_j)$. Since $p_y \in \bar{A}$ by the definition of \bar{A} , we obtain that h'_j is onto, as claimed.

Now, the morphism $\bar{Y} \rightarrow \bar{X}$ is almost as required: \bar{Y} is open in Y and α is a closed immersion. In addition, since $\mathbf{y} \subset \bar{\mathfrak{X}}_0$, the image of A_0 under the homomorphism $A_0 \rightarrow B_0 \rightarrow \bar{B} \rightarrow \bar{B}(y)$ is contained in R , and hence the image of A_0 in \bar{B} is actually contained in \bar{A} . So, it only remains to decrease the A -subalgebra $\bar{A} \subset \bar{B}$ so that $\bar{X} = \text{Spec}(\bar{A})$ becomes of finite type over X but all good properties are preserved: α is still a closed immersion, and \bar{A} contains the image of A_0 in \bar{B} . As we saw, α being a closed immersion is equivalent to surjectivity of the homomorphisms $h_i : \bar{A} \otimes B_i \rightarrow \bar{B}$ and $h'_j : \bar{A} \otimes C_j \rightarrow \bar{C}_j$. Since the homomorphisms $B_i \rightarrow \bar{B}$ and $C_j \rightarrow \bar{C}_j$ are of finite type, all we need for surjectivity of h_i 's and h'_j 's is a finite subset $S \subset \bar{A}$. So, replacing \bar{A} with its A_0 -subalgebra generated by S we obtain \bar{X} as required. Obviously, $\text{Val}_{\bar{Y}}(\bar{X})$ is an affinoid domain containing \mathbf{y} , and $\text{Val}_{\bar{Y}}(\bar{X})$ is contained in $\bar{\mathfrak{X}}_0$ because \bar{Y} is an open subscheme in Y_0 and the morphism $\bar{Y} \rightarrow X_0$ (obtained as $\bar{Y} \rightarrow Y_0 \rightarrow X_0$) factors through \bar{X} . \square

Corollary 3.3.5. *The space $\bar{\mathfrak{X}}$ is qcqs.*

Proof. Any open subdomain is quasi-compact by Proposition 3.1.10, and their intersection is quasi-compact by Lemma 3.3.2. Since open subdomains generate the topology of $\bar{\mathfrak{X}}$ by Theorem 3.3.4 we obtain the corollary. \square

Recall that we defined in Remark 3.1.9 the contraction $\pi_{\bar{\mathfrak{X}}} : \bar{\mathfrak{X}} \rightarrow \bar{\mathfrak{X}}$ and used it to define the maps $\text{Val}(g) : \bar{\mathfrak{X}}' \rightarrow \bar{\mathfrak{X}}$ for $g : f' \rightarrow f$.

Corollary 3.3.6. *The contraction $\pi_{\bar{\mathfrak{X}}}$ is continuous. In particular, the maps $\text{Val}(g)$ are continuous.*

Proof. Since open subdomains $\mathfrak{X}' = \text{Val}_{Y'}(X')$ form a basis of the topology of \mathfrak{X} by Theorem 3.3.4, it suffices to prove that the preimage of \mathfrak{X}' in $\text{Spa}(Y, X)$ is open. Since the minimality condition in $\text{Spa}(Y, X)$ and $\text{Spa}(Y', X')$ agree, $\pi^{-1}(\mathfrak{X}')$ coincides with the open affine subset $\text{Spa}(Y', X')$. \square

3.4. Y -blow ups of X . In this section we assume that f is affine. Then we will show that there exists a large family of projective Y -modifications of X having good functorial properties. Using these morphisms we will be able to describe the set $\text{Val}_Y(X)$ very concretely. Since the results of §3.4 are inspired in part by Raynaud's theory of formal models, we will sometimes indicate similarity between our results and Raynaud's theory by referencing to [BL].

Definition 3.4.1. A Y -modification $g_i : X_i \rightarrow X$ is called a Y -blow up of X if there exists a g_i -ample \mathcal{O}_{X_i} -module \mathcal{L} provided with a homomorphism $\varepsilon : \mathcal{O}_{X_i} \rightarrow \mathcal{L}$ such that $f_i^*(\varepsilon) : \mathcal{O}_Y \xrightarrow{\sim} f_i^*(\mathcal{L})$. We call ε a Y -trivialization of \mathcal{L} ; actually it is a section of \mathcal{L} that is invertible on the image of Y .

It will be more convenient to say X -ample instead of g_i -ample in the sequel.

Lemma 3.4.2. *The Y -blow ups satisfy the following properties.*

- (i) *Suppose that $X_j \rightarrow X_i$ and $X_i \rightarrow X$ are Y -modifications such that X_j is a Y -blow up of X . Then X_j is a Y -blow up of X_i .*
- (ii) *The family of Y -blow ups of X is filtered.*
- (iii) *The composition of Y -blow ups $g_{ij} : X_j \rightarrow X_i$ and $g_i : X_i \rightarrow X$ is a Y -blow up.*

Proof. The first statement is obvious because any X -ample \mathcal{O}_{X_j} -module \mathcal{L} is X_i -ample, and the notion of Y -trivialization of \mathcal{L} depends only on the morphism $f_j : Y \rightarrow X_j$.

(ii) Let X_i, X_j be two Y -blow ups of X . Find X -ample sheaves $\mathcal{L}_i, \mathcal{L}_j$ with Y -trivializations $\varepsilon_i, \varepsilon_j$. Then the X -proper scheme $X_{ij} = X_i \times_X X_j$ possesses an X -ample sheaf $\mathcal{L} = p_i^*(\mathcal{L}_i) \otimes p_j^*(\mathcal{L}_j)$, where p_i, p_j are the projections. The natural isomorphism $\mathcal{O}_{X_{ij}} \xrightarrow{\sim} \mathcal{O}_{X_i} \otimes \mathcal{O}_{X_j}$ followed by $f_i^*(\varepsilon_i) \otimes f_j^*(\varepsilon_j) : \mathcal{O}_{X_{ij}} \otimes \mathcal{O}_{X_{ij}} \rightarrow \mathcal{L}$ provides a Y -trivialization of \mathcal{L} . Consider the scheme-theoretic image X' of Y in X_{ij} , and let \mathcal{L}' and ε' be the pull backs of \mathcal{L} and ε . Then $(X', \mathcal{L}', \varepsilon')$ is a Y -blow up of X which dominates X_i and X_j .

(iii) Choose an X -ample \mathcal{O}_{X_i} -sheaf \mathcal{L}_i and an X_i -ample \mathcal{O}_{X_j} -sheaf \mathcal{L}_j with Y -trivializations ε_i and ε_j . By [EGA, II, 4.6.13(ii)], the sheaf $\mathcal{L}_j \otimes g_{ij}^*(\mathcal{L}_i^{\otimes n})$ is X -ample for sufficiently large n . It remains to notice that the composition of $\mathcal{O}_{X_j} \xrightarrow{\sim} \mathcal{O}_{X_j} \otimes \mathcal{O}_{X_j}^{\otimes n}$ with $\varepsilon_j \otimes g_{ij}^*(\varepsilon_i^{\otimes n})$ is a Y -trivialization. \square

We will need an explicit description of Y -blow ups. Let $\mathcal{E} \subset f_*(\mathcal{O}_Y)$ be a finitely generated \mathcal{O}_X -submodule containing the image of \mathcal{O}_X , and let $\mathcal{E}^n \subset f_*(\mathcal{O}_Y)$ denote the \mathcal{O}_X -modules which are powers of \mathcal{E} with respect to the natural multiplication on $f_*(\mathcal{O}_Y)$ (so \mathcal{E}^0 is the image of \mathcal{O}_X). We claim that $X_{\mathcal{E}} := \mathbf{Proj}(\bigoplus_{n=0}^{\infty} \mathcal{E}^n)$ is a Y -modification of X . Clearly, $X_{\mathcal{E}}$ is X -projective and there is a natural morphism $g_{\mathcal{E}} : Y = \mathbf{Spec}(f_*(\mathcal{O}_Y)) \rightarrow \mathbf{Spec}(\bigcup_{n=0}^{\infty} \mathcal{E}^n)$ where the union is taken inside $f_*(\mathcal{O}_Y)$. The target of $g_{\mathcal{E}}$ is the X -affine chart of $X_{\mathcal{E}}$ defined by non-vanishing of the section $s \in \Gamma(\mathcal{E})$ which comes from the unit section of \mathcal{O}_X , in particular, a map $Y \rightarrow X_{\mathcal{E}}$ naturally arises. In addition, the very ample sheaf $\mathcal{O}_{X_{\mathcal{E}}}(1)$ on $X_{\mathcal{E}}$ has a Y -trivialization $\mathcal{O}_{X_{\mathcal{E}}} \rightarrow \mathcal{O}_{X_{\mathcal{E}}}(1)$ induced by s . So, among all properties of Y -blow

ups it remains to check that $g_{\mathcal{E}}$ is schematically dominant. The latter can be checked locally over X , so assume that $X = \text{Spec}(A)$, $Y = \text{Spec}(B)$ and $E \subset B$ is an A -module containing 1. Then $X_E = \text{Proj}(\bigoplus_{n=0}^{\infty} E^n)$ is glued from affine charts $(X_E)_b$ given by non-vanishing of elements $b \in E$, so it suffices to show that the morphism $\alpha : Y \times_{X_E} (X_E)_b \rightarrow (X_E)_b$ is schematically dominant. Note that the source is the localization of Y at b , and so it is isomorphic to $\text{Spec}(B_b)$, and the target is $\text{Spec}(C)$ where C is the zeroth graded component of $(\bigoplus_{n=0}^{\infty} E^n)_b$. But $C = \text{inj} \lim_n b^{-n}(E^n/I_n)$, where I_n is the submodule of elements killed by a power of b , and the kernel of the homomorphism $E^n \hookrightarrow B \rightarrow B_b$ is I_n . Hence $b^{-n}(E^n/I_n) \hookrightarrow B_b$ and therefore $C \hookrightarrow B$. In particular, α is schematically dominant.

Lemma 3.4.3. *Any Y -blow up of X is isomorphic to some $X_{\mathcal{E}}$ as a Y -blow up of X .*

Proof. Let $g_i : X_i \rightarrow X$ be a Y -blow up. Find an X -ample \mathcal{O}_{X_i} -module \mathcal{L} with a Y -trivialization $\varepsilon : \mathcal{O}_{X_i} \rightarrow \mathcal{L}$. Then there is a closed immersion of X -schemes $h : X_i \rightarrow P := \mathbf{Proj}(\bigoplus_{n=0}^{\infty} (g_i)_* \mathcal{L}^{\otimes n})$ and the morphism $h \circ f_i : Y \rightarrow X_i \rightarrow P$ factors through the chart of P given by non-vanishing of the section $s \in \Gamma((g_i)_* \mathcal{L})$ corresponding to ε . The latter chart is of the form $\mathbf{Spec}(\mathcal{A})$ where \mathcal{A} is the zeroth graded component of the localization $(\bigoplus_{n=0}^{\infty} (g_i)_* \mathcal{L}^{\otimes n})_s$. Composing the \mathcal{O}_X -homomorphism $(g_i)_* \mathcal{L} \rightarrow \mathcal{A}$ that takes u to $s^{-1}u$ with the \mathcal{O}_X -homomorphism $\mathcal{A} \rightarrow f_*(\mathcal{O}_Y)$ corresponding to f_i we obtain a homomorphism $(g_i)_* \mathcal{L} \rightarrow f_*(\mathcal{O}_Y)$ that takes s to the unit section. Now we can define \mathcal{E} to be the image of $(g_i)_* \mathcal{L}$ in $f_*(\mathcal{O}_Y)$, and we claim that actually $X_i \xrightarrow{\sim} X_{\mathcal{E}}$ as a Y -modification of X . Indeed, the obvious epimorphism $\bigoplus_{n=0}^{\infty} (g_i)_* \mathcal{L}^{\otimes n} \rightarrow \bigoplus_{n=0}^{\infty} \mathcal{E}^n$ corresponds to a closed immersion $X_{\mathcal{E}} \rightarrow P$ which agrees with the morphisms $Y \rightarrow X_{\mathcal{E}}$ and $Y \rightarrow P$. Since, the first morphism is schematically dominant, $X_{\mathcal{E}}$ is the schematic image of Y in P , hence it must coincide with X_i as the closed subscheme of P . \square

Corollary 3.4.4. *Assume that X' is an open subscheme of X and $Y' = f^{-1}(X')$. Then any Y' -blow up $X'_i \rightarrow X'$ extends to a Y -blow up $X_i \rightarrow X$.*

Proof. Let $f' : Y' \rightarrow X'$ be the restriction of f , so $f'_*(\mathcal{O}_{Y'})$ is the restriction of $f_*(\mathcal{O}_Y)$ on X' . By the lemma, a Y' -blow up of X' is determined by a finitely generated $\mathcal{O}_{X'}$ -submodule $\mathcal{E}' \subset f'_*(\mathcal{O}_{Y'})$ containing the image of $\mathcal{O}_{X'}$. By [EGA I, 6.9.7], one can extend \mathcal{E}' to a finitely generated \mathcal{O}_X -submodule $\mathcal{E} \subset f_*(\mathcal{O}_Y)$. Replacing \mathcal{E} by $\mathcal{E} + \mathcal{O}_X$, if necessary, we can achieve that \mathcal{E} contains the image of \mathcal{O}_X . Now, \mathcal{E} defines a required extension of the blow up. \square

Remark 3.4.5. (i) Lemma 3.4.3 indicates that the notion of Y -blow up is in some sense a generalization of the notion of U -admissible blow up, where $i : U \hookrightarrow X$ is a schematically dense open subscheme, to the case of an arbitrary affine morphism $Y \rightarrow X$. Indeed, there is much similarity, but the notions are not equivalent in general: both U -admissible blow ups and U -blow ups are of the form $\text{Proj}(\bigoplus_{n=0}^{\infty} \mathcal{E}^n)$, but in the first case \mathcal{E} is an \mathcal{O}_X -submodule of \mathcal{O}_X which is trivial over U , and in the second one \mathcal{E} is an \mathcal{O}_X -submodule of $i_*(\mathcal{O}_U)$ that contains \mathcal{O}_X (so, it is trivial over U as well). The important case when these notions agree was pointed out by the referee: it follows from [EGA, II, 3.1.8(iii)] that U -admissible blow ups and U -blow ups agree when $X \setminus U$ is the zero set of an invertible sheaf of ideals.

(ii) Basic facts concerning compositions, extensions, etc., (see the above lemmas) hold for both families of U -modifications, but a slight advantage of U -blow ups is

that the proofs seem to be easier. For example, compare with [Con, 1.2] where one proves that U -admissible blow ups are preserved by compositions.

The following lemma is an analog of [BL, 4.4].

Lemma 3.4.6. *Given a quasi-compact open subset $\mathfrak{U} \subset \mathfrak{X} = \text{Val}_Y(X)$, there exists a Y -modification $X' \rightarrow X$ and an open subscheme $U \subset X'$ such that \mathfrak{U} is the preimage of U in \mathfrak{X} .*

Proof. If X_1, \dots, X_n form a finite open affine covering of X and $Y_i = f^{-1}(X_i)$ then $\mathfrak{X}_i = \text{Val}_{Y_i}(X_i)$ form an open covering of \mathfrak{X} by Lemma 3.1.8. It suffices to separately solve our problem for each \mathfrak{X}_i with $\mathfrak{U}_i := \mathfrak{U} \cap \mathfrak{X}_i$ because any Y_i -blow up of X_i extends to a Y -blow up of X , and Y -blow ups of X form a filtered family. Thus, we can assume that $X = \text{Spec}(A)$, and then $Y = \text{Spec}(B)$. We can furthermore assume that $\mathfrak{U} = \mathfrak{X} \cap \text{Spa}(B_b, A[a_1/b, \dots, a_n/b])$ with $a_i, b \in B$ because as we saw in the proof of Lemma 3.1.2, the sets $\text{Spa}(B_b, A[a_1/b, \dots, a_n/b])$ form a basis of the topology of $\text{Spa}(B, A)$. Now, the morphism $Y \rightarrow \text{Proj}(A[T_1, T_{a_1}, \dots, T_{a_n}, T_b])$ defined by $(1, a_1, \dots, a_n, b)$ determines a required Y -blow up $X' \rightarrow X$ with U given by the condition $T_b \neq 0$. \square

Corollary 3.4.7. *The map $\psi : \text{Val}_Y(X) \rightarrow \text{RZ}_Y(X)$ is a homeomorphism.*

Proof. Recall that ψ is surjective and continuous by Propositions 2.2.1 and 3.1.10, respectively. From other side, the lemma implies that ψ is injective and open. Indeed, any open quasi-compact $\mathfrak{U} \subset \mathfrak{X}$ is the full preimage of some $U \subset X'$ for a Y -modification $X' \rightarrow X$, hence $\psi(\mathfrak{U})$, which is the full preimage of U in $\text{RZ}_Y(X)$, is open. In addition, since any pair of different points of \mathfrak{X} is distinguished by some open quasi-compact set $\mathfrak{U} \subset \mathfrak{X}$, their images in an appropriate X' do not coincide. \square

We use the corollary to identify \mathfrak{X} with $\text{RZ}_Y(X)$ when f is decomposable. In particular, this provides \mathfrak{X} with a sheaf $\mathcal{O}_{\mathfrak{X}}$ of regular functions which was earlier defined on $\text{RZ}_Y(X)$, and for any point $\mathbf{x} \in \mathfrak{X}$, thanks to Proposition 2.2.1, the semi-valuation ring $\mathcal{O}_{\mathbf{x}}$ obtains a new interpretation as the stalk of $\mathcal{O}_{\mathfrak{X}}$ at \mathbf{x} . As another corollary of Lemma 3.4.6 we obtain the following version of Chow lemma.

Corollary 3.4.8. *Any Y -modification $\overline{X} \rightarrow X$ is dominated by a Y -blow up of X .*

Proof. Let $\overline{U}_1, \dots, \overline{U}_n$ be an affine covering of \overline{X} , and let Y_i and \mathfrak{U}_i denote the preimages of \overline{U}_i in Y and \mathfrak{X} , respectively. By Lemma 3.4.6, we can find a Y -blow up $X' \rightarrow X$ and a covering $\{U'_i\}$ of X' , whose preimage in \mathfrak{X} coincides with $\{\mathfrak{U}_i\}$. Note that the scheme-theoretic image X'' of Y in $\overline{X} \times_X X'$ is a Y -modification of both X' and \overline{X} . So, it suffices to show that X'' is a Y -blow up of X .

Since the preimages of \overline{U}_i and U'_i in \mathfrak{X} coincide, their preimages in X'' coincide too, and we will denote them as $U''_i \hookrightarrow X''$. Consider the induced Y -modification $h : X'' \rightarrow X'$ with restrictions $h_i : U''_i \rightarrow U'_i$. For any $1 \leq i \leq n$, the proper morphism h_i is affine because the morphism $\overline{U}_i \rightarrow X$ is affine and U''_i is closed in $U'_i \times_X \overline{U}_i$. Thus, h_i is finite, and therefore h is finite. We claim that finiteness of h implies that it is a Y -blow up (this claim is an analog of [BL, 4.5]). Indeed, $\mathcal{O}_{X''}$ is very ample relatively to h because h is affine, and the identity homomorphism gives its Y -trivialization. Thus, X'' is a Y -blow up of X by Lemma 3.4.2(iii). \square

3.5. Decomposable morphisms. In this section we will complete a basic description of the relative Riemann-Zariski space \mathfrak{X} associated with a separated morphism $f : Y \rightarrow X$ between qcqs schemes by proving that the finite union of open domains is an open domain, and any open domain in \mathfrak{X} is of the form $\text{Val}_{\bar{Y}}(\bar{X})$ where the morphism $\bar{Y} \rightarrow \bar{X}$ is affine and schematically dominant. The first claim actually means that any quasi-compact open subset is an open domain, i.e. admits a model by a morphism of schemes, and the second claim states that this model can be chosen to be affine. In particular, applying the second claim to \mathfrak{X} itself we obtain a bijection $\text{Val}_{\bar{Y}}(\bar{X}) \xrightarrow{\sim} \text{Val}_Y(X)$ with $\bar{Y} = Y$ and affine morphism $\bar{Y} \rightarrow \bar{X}$. But then \bar{X} is proper over X by the valuative criterion 3.2.3, and hence X admits a Y -modification \bar{X} such that the morphism $Y \rightarrow \bar{X}$ is affine. Thus, the morphism $f : Y \rightarrow X$ is decomposable and this gives a new proof of Theorem 1.1.3. In particular, one obtains new proofs of Nagata compactification and Thomason approximation theorems.

Theorem 3.5.1. *Let $f : Y \rightarrow X$ be a separated morphism between qcqs schemes and $\mathfrak{X} = \text{Val}_Y(X)$. Then*

- (i) *open domains in \mathfrak{X} are closed under finite unions,*
- (ii) *any open domain \mathfrak{X}' is of the form $\text{Val}_{\bar{Y}}(\bar{X})$, where the morphism $\bar{Y} \rightarrow \bar{X}$ is affine and schematically dominant.*

Proof. Note that any affinoid domain satisfies the assertion of (ii) (since schematical dominance is achieved by simply replacing \bar{X} with the schematic image of \bar{Y}), and by Theorem 3.3.4 and Corollary 3.3.5, \mathfrak{X}' admits a finite affinoid covering. Therefore, both (i) and (ii) would follow if we prove the following claim: the union of two domains satisfying the assertion of (ii) is an open domain that satisfies the assertion of (ii). So, we assume that $\mathfrak{X}' = \mathfrak{X}_1 \cup \mathfrak{X}_2$ where $\mathfrak{X}_i = \text{Val}_{Y_i}(X_i)$ with $i \in \{1, 2\}$ are open subdomains with affine morphisms $Y_i \rightarrow X_i$.

Set $\mathfrak{X}_{12} = \mathfrak{X}_1 \cap \mathfrak{X}_2$ and $Y_{12} = Y_1 \cap Y_2$. In the sequel, we will act as in Step 3 of the proof of Theorem 1.1.2, and the main difference is that we will use Y_i -blow ups instead of affine morphisms. For reader's convenience, we provide a commutative diagram containing the main objects which were and will be introduced.

$$\begin{array}{ccccc}
 Y_1 & \longleftarrow & Y_{12} & \hookrightarrow & Y_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathfrak{X}_1 & \longleftarrow & \mathfrak{X}_{12} & \hookrightarrow & \mathfrak{X}_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 Z_1 & \longleftarrow & Z_{12} & \hookrightarrow & Z_2 \\
 \downarrow & & \swarrow & \searrow & \downarrow \\
 X_1 & \longleftarrow & X'_1 & & X'_2 \hookrightarrow X_2
 \end{array}$$

Since Y_i 's are X_i -affine, Lemma 3.4.6 implies that we can replace X_i 's by their Y_i -blow ups such that each X_i contains an open subscheme X'_i , whose preimage in \mathfrak{X}_i coincides with \mathfrak{X}_{12} . Then the preimage of X'_i in Y is, obviously, Y_{12} . It can be impossible to glue X_i 's along X'_i 's, but by Lemma 3.1.8(ii), we at least know that $\text{Val}_{Y_{12}}(X'_i) \xrightarrow{\sim} \mathfrak{X}_{12}$ for $i = 1, 2$. Let T be the scheme-theoretic image of Y_{12} in

$X'_1 \times_X X'_2$; it is obviously separated over X'_i 's. Moreover, $\text{Val}_{Y_{12}}(T) \xrightarrow{\sim} \text{Val}_{Y_{12}}(X'_1) \cap \text{Val}_{Y_{12}}(X'_2) = \mathfrak{X}_{12}$ by Lemma 3.3.2, and, therefore, T is a Y_{12} -modification of X'_i 's by the valuative criterion 3.2.3.

By Corollary 3.4.8, we can find a Y_{12} -blow up $T' \rightarrow X'_1$, which dominates T . It still can happen that T' is not a Y_{12} -blow up of X'_2 , but it is dominated by a Y_{12} -blow up $Z_{12} \rightarrow X'_2$. Then $Z_{12} \rightarrow T'$ is a Y_{12} -blow up by Lemma 3.4.2(i), and hence $Z_{12} \rightarrow X'_1$ is a Y_{12} -blow up by Lemma 3.4.2(iii). By Lemma 3.4.4, we can extend the Y_{12} -blow ups $Z_{12} \rightarrow X'_i$ to Y_i -blow ups $Z_i \rightarrow X_i$. Then, the finite type X -schemes Z_i can be glued along the subschemes X -isomorphic to Z_{12} to a single X -scheme \overline{X} of finite type, and the schematically dominant affine morphisms $Y_i \rightarrow Z_i$ glue to a single schematically dominant affine morphism $\overline{Y} \rightarrow \overline{X}$. Note that $\text{Val}_{Y_i}(Z_i) = \mathfrak{X}_i$ is the preimage of Z_i in $\text{Val}_{\overline{Y}}(\overline{X})$, in particular, the latter is covered by its open subdomains \mathfrak{X}_i , $i \in \{1, 2\}$. Now, it remains to show that $\text{Val}_{\overline{Y}}(\overline{X})$ is an open subdomain in \mathfrak{X} , since this would immediately imply that $\text{Val}_{\overline{Y}}(\overline{X})$ is a required model of \mathfrak{X} . The morphism $\alpha : \overline{Y} \rightarrow \overline{X} \times_X Y$ is glued from the morphisms $\alpha_i : Y_i \rightarrow Z_i \times_X Y$ because Y_i is the preimage of Z_i in Y , but α_i 's are closed immersions by the construction. So, α is a closed immersion as well, and we are done. \square

Corollary 3.5.2. *The map $\eta : Y \rightarrow \mathfrak{X} := \text{RZ}_Y(X)$ is injective, each point $\mathbf{x} \in \text{RZ}_Y(X)$ possesses a unique minimal generalization y in $\eta(Y)$, $\mathcal{M}_{\mathfrak{X}, \mathbf{x}} \xrightarrow{\sim} \mathcal{O}_{Y, y}$, and the stalk $\mathcal{M}_{\mathfrak{X}, \mathbf{x}}$ is the semi-fraction ring of the semi-valuation ring $\mathcal{O}_{\mathfrak{X}, \mathbf{x}}$. In particular, $\mathcal{O}_{\mathfrak{X}}$ is a subsheaf of $\mathcal{M}_{\mathfrak{X}}$.*

Proof. By Theorem 3.5.1 and Corollary 3.4.7, we can identify \mathfrak{X} with $\text{Val}_Y(X)$. So, a point \mathbf{x} can be interpreted as an X -valuation (y, R, ϕ) on Y . Then it is clear that the map η sends $y \in Y$ to a trivial valuation $(y, k(y), f|_y)$ (with the obvious morphism $f|_y : \text{Spec}(k(y)) \rightarrow X$), and for an arbitrary $\mathbf{x} = (y, R, \phi)$ its minimal generalization in $\eta(Y)$ is $(y, k(y), f|_y)$. Uniqueness of minimal generalization implies that the stalk of $\mathcal{M}_{\mathfrak{X}} = \eta_*(\mathcal{O}_Y)$ at \mathbf{x} is simply $\mathcal{O}_{Y, y}$, so it remains to recall that the latter is the semi-fraction field of the semi-valuation ring $\mathcal{O}_{\mathfrak{X}}$ defined in §2.1, which coincides with the stalk $\mathcal{O}_{\mathfrak{X}, \mathbf{x}}$ by Proposition 2.2.1. \square

REFERENCES

- [BGR] Bosch, S.; Güntzer, U.; Remmert, R.: *Non-Archimedean analysis. A systematic approach to rigid analytic geometry*, Springer, Berlin-Heidelberg-New York, 1984.
- [BL] Bosch, S.; Lütkebohmert, W.: *Formal and rigid geometry. I. Rigid spaces.*, Math. Ann. **295** (1993), no. 2, 291–317.
- [BLR] Bosch, S.; Lütkebohmert, W.; Raynaud, M.: *Formal and rigid geometry. IV. The reduced fibre theorem.*, Inv. Math. **119** (1995), no. 2, 361–398.
- [CLO] Conrad, B.; Lieblich, M.; Olsson, M.: *Nagata compactification for algebraic spaces*, preprint, arXiv:[0910.5008].
- [Con] Conrad, B.: *Deligne's notes on Nagata compactifications*, J. Ramanujan Math. Soc. **22** (2007), no. 3, 205–257.
- [EGA] Dieudonné, J.; Grothendieck, A.: *Éléments de géométrie algébrique*, Publ. Math. IHES, **4, 8, 11, 17, 20, 24, 28, 32**, (1960-7).
- [EGA I] Dieudonné, J.; Grothendieck, A.: *Éléments de géométrie algébrique, I: Le langage des schémas*, second edition, Springer, Berlin, 1971.
- [FK] Fujiwara, K; Kato, F.: *Rigid geometry and applications*, Moduli spaces and arithmetic geometry, 327–386, Adv. Stud. Pure Math., 45, Math. Soc. Japan, Tokyo, 2006.
- [Har] Hartshorne, R.: *Algebraic geometry*, Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977. xvi+496 pp.

- [Hub1] Huber, R.: *Continuous valuations*, Math.Zeit. **212** (1993), 455–477.
- [Hub2] Huber, R.: *A generalization of formal schemes and rigid analytic spaces*, Math. Zeit. **217** (1994), 513–551.
- [Lüt] Lütkebohmert, W.: *On compactification of schemes*, Manuscripta Math. **80** (1993), 95–111.
- [Nag] Nagata, M.: *A generalization of the embedding problem*, J. Math. Kyoto **3** (1963), 89–102.
- [R1] Raoult, J.-C.: *Compactification des espaces algébriques normaux*, C.R. Acad. Sci. Paris Sér. A-B **273**, (1971), pp. A766–A767.
- [R2] Raoult, J.-C.: *Compactification des espaces algébriques*, C.R. Acad. Sci. Paris Sér. A-B **278**, (1974), pp. 867–869.
- [RG] Raynaud, M.; Gruson, L.: *Critères de platitude et de projectivité*, Inv. Math. **13** (1971), 1–89.
- [Rydh] Rydh, D.: *Compactification of Deligne-Mumford stacks*, preprint.
- [Tem1] Temkin, M.: *On local properties of non-Archimedean analytic spaces.*, Math. Ann. **318** (2000), no. 3, 585–607.
- [Tem2] Temkin, M.: *Stable modification of relative curves*, J. Alg. Geom. **19** (2010), 603–677.
- [TT] Thomason, R.; Trobaugh, T.: *Higher algebraic K-theory of schemes and of derived categories*, in The Grothendieck Festschrift, Vol. III, 247–436, Progress in Mathematics vol. 88, Birkhauser, Boston, 1990.

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIV'AT RAM, JERUSALEM, 91904, ISRAEL
E-mail address: `temkin@math.huji.ac.il`